Berberis vulgaris

COMMON NAME

European barbery

FAMILY

Berberidaceae

AUTHORITY

Berberis vulgaris L.

FLORA CATEGORY

Vascular - Exotic

STRUCTURAL CLASS

Trees & Shrubs - Dicotyledons

NVS CODE

BERVUL

CONSERVATION STATUS

Not applicable

DETAILED DESCRIPTION

Small deciduous shrub to about 2m. Spines generally 3-partite. Petiole to about 1 cm lamina up to 8 x 2.5 cm, but generally much smaller than this. Leaves with obvious veins and serrulate margins. Medium yellow flowers in racemes. Berries ripening to crimson-scarlet, oblong and up to $11 \, \text{mm}$ long.

SIMILAR TAXA

Similar in general appearance to B. wilsonae, but B. vulgaris has petiolate leaves, with obvious veination and serrulate margins.

FLOWERING

October, November

FLOWER COLOURS

Yellow

YEAR NATURALISED

1875

ORIGIN

Europe

ETYMOLOGY

berberis: From the Arabic name berberys

Reproduction

Seed and suckering.

Seed

prolific.

Dispersal

Birds, people

MORE INFORMATION

https://www.nzpcn.org.nz/flora/species/berberis-vulgaris/





Ahuriri Valley, Canterbury. Photographer: John Barkla, Licence: CC BY.



Ahuriri Valley, Canterbury. Photographer: John Barkla, Licence: CC BY.