

# Celmisia hieraciifolia var. oblonga

## COMMON NAME

mountain daisy

## SYNONYMS

None

## FAMILY

Asteraceae

## AUTHORITY

Celmisia hieraciifolia var. oblonga Kirk

## FLORA CATEGORY

Vascular – Native

## ENDEMIC TAXON

Yes

## ENDEMIC GENUS

No

## ENDEMIC FAMILY

No

## STRUCTURAL CLASS

Herbs - Dicotyledonous composites

## CHROMOSOME NUMBER

2n = 108

## CURRENT CONSERVATION STATUS

2017 | Data Deficient

## PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

## DISTRIBUTION

Endemic. New Zealand: North Island (from the Ruahine and Tararua Ranges south), South Island (south to North Canterbury and Westland).

## HABITAT

Montane to subalpine in grassland, herbfield, fellfield and debris slopes.

## DETAILED DESCRIPTION

Medium tufted herb, stock usually simple, pseudo-stem  $\pm$  40 mm long. **Lamina** coriaceous, c. 30–70  $\times$  4–15 mm; narrow-oblong to narrow-spathulate; upper surface glabrous, sometimes slightly viscid, pale green when fresh; lower densely clad in closely appressed satiny pale yellow to buff or almost white tomentum, nerves conspicuous in lower part; obtuse, sometimes apiculate; margins crenate to finely crenate-dentate, teeth small, sparse rarely almost absent, slightly narrowed to broad grooved short petiole or directly into glabrous sheath c. 20  $\times$  10 mm. **Scape** rather slender, 50–200 mm long, viscid with dense glandular hairs. **Bracts** linear to subulate, glandular-pubescent, lower up to 40 mm long. **Capitula** 20–40 mm diameter; involucre bracts  $\pm$  13 mm long, outer densely glandular-pubescent, linear-subulate, acute, midrib distinct, margins sometimes  $\pm$  floccose. **Ray-florets** numerous, c. 5 mm long, white, tube slender, limb narrow-linear. **Disk-florets** c. 7 mm long, tubular to narrow-funnelform, teeth narrow-triangular. **Achenes** c. 2–5 mm long, strongly ribbed, cylindrical to subfusiform; hairs usually few, weak. **Pappus-hairs** white to sordid-white, slender, up to c. 6 mm long, finely barbellate.



Mt Stokes January. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.



Mount Stokes, Marlborough. Photographer: Gillian M. Crowcroft, Licence: All rights reserved.

## FLOWERING

October–January

## FLOWER COLOURS

White, Yellow

## FRUITING

November–April

## PROPAGATION TECHNIQUE

Unknown. Probably best grown from fresh seed and like many *Celmisia* this species will probably dislike high humidity and drying out.

## ETYMOLOGY

**celmisia:** Apparently named after Kelmis, one of Idaean Dactyls, a group of skilled mythical beings associated with the Mother Goddess Rhea in Greek mythology. Kelmis, whose name means ‘casting’, was a blacksmith and childhood friend of Zeus, son of Rhea and later king of the gods. In Ovid’s ‘Metamorphoses’, Kelmis is described as offending Zeus who turned him into adamant so he was as hard as a tempered blade

**hieraciifolia:** Having foliage like Hieracium (genus name from the word for hawk)

**oblonga:** Oblong

## WHERE TO BUY

Not commercially available.

## NOTES ON TAXONOMY

The status of the three varieties of *C. hieraciifolia* needs critical investigation.

## ATTRIBUTION

Description adapted from Allan (1961)

## REFERENCES AND FURTHER READING

Allan HH. 1961. Flora of New Zealand, Volume I. Indigenous Tracheophyta: Psilopsida, Lycopsida, Filicopsida, Gymnospermae, Dicotyledones. Government Printer, Wellington, NZ. 1085 p.

## MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/celmisia-hieraciifolia-var-oblonga/>