# **Dryopteris filix-mas**

## **COMMON NAME**

male fern

#### **FAMILY**

Dryopteridaceae

#### **AUTHORITY**

Dryopteris filix-mas (L.) Schott

## **FLORA CATEGORY**

Vascular - Exotic

#### STRUCTURAL CLASS

**Ferns** 

## **NVS CODE**

**DRYFIL** 

#### **CONSERVATION STATUS**

Not applicable

#### **HABITAT**

Terrestrial. Streamsides, open scrub, damp forest.

### **DETAILED DESCRIPTION**

Stout rhizomatous fern. **Laminae bipinnate** up to 125 x 30 cm, dark green above, paler green below. **Primary pinnae** in 25–50 pairs, secondary pinnae in 15–30 pairs. **Sori** round, 1–5 pairs on each secondary pinnae.

#### **SIMILAR TAXA**

Dryopteris is closely related to Polystichium and Lastreopsis, but distinguished from Polystichum by the kidney-shaped indusia, and from Lastreopsis by its abundance of stipe-scales and absence of hairs. D. filix-mas is closely related to D. affinis but can be distinguished by the second pinnae being round-ended and margins toothed throughout.

## YEAR NATURALISED

1958

#### **ORIGIN**

Europe, Asia, N. America

#### **Reason For Introduction**

Ornamental

## Reproduction

Sexually from spores.

## MORE INFORMATION

https://www.nzpcn.org.nz/flora/species/dryopteris-filix-mas/





Dryopteris filix-mas. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.



Dunedin Town belt. Photographer: John Barkla, Licence: CC BY.