

Acaena dumicola

COMMON NAME

bidibid, piripiri

FAMILY

Rosaceae

AUTHORITY

Acaena dumicola B.H.Macmill.

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

NVS CODE

ACADUM

CHROMOSOME NUMBER

2n = 42

CURRENT CONSERVATION STATUS

2017 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

FLOWER COLOURS

Green

LIFE CYCLE

Spiny hypanthia are dispersed by attaching to fur, feathers and clothing and possibly also dispersed by wind and granivory (Thorsen et al., 2009).

ETYMOLOGY

acaena: From the Greek 'akanthos' thorn, referring to the spiny calyx that many species have

dumicola: In thickets or shrubs

ATTRIBUTION

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 2009 Vol. 11 No. 4 pp. 285-309

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/acaena-dumicola/>

