

# Aciphylla crosby-smithii

## FAMILY

Apiaceae

## AUTHORITY

*Aciphylla crosby-smithii* Petrie

## FLORA CATEGORY

Vascular – Native

## ENDEMIC TAXON

Yes

## ENDEMIC GENUS

No

## ENDEMIC FAMILY

No

## STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

## NVS CODE

ACICRO

## CURRENT CONSERVATION STATUS

2012 | At Risk – Naturally Uncommon | Qualifiers: RR

## PREVIOUS CONSERVATION STATUSES

2009 | At Risk – Naturally Uncommon

2004 | Range Restricted

## BRIEF DESCRIPTION

Robust elongate tufts of broad spiky leaves with white flowers in a ball on a long fleshy stalk in southern Fiordland

## DISTRIBUTION

Endemic. South Island; southern Fiordland (possibly also in western Otago and elsewhere in Fiordland)

## HABITAT

Exposed rocky slopes and short snow tussock herbfield. 1400-1600 m a.s.l.

## FEATURES

Herb forming stout tufts to 40 cm tall in flower. Rosettes several, crowded, +/- elongate, forming cushionlike mass to 60 cm diam. Leaves many, extending along base of flowering stem, stiff, 1-pinnate; sheath broad, membranous, 4 cm long x 1.5 cm wide, slowly narrowing to apex; stipules similar to pinnae, 35 mm long x 5 mm wide, narrowing to small mucro; leaf with pinnae in 2-3 pairs, petiole and internode (inter-pinnae spacing) short, linear-oblong, c. 3 cm long x 3-4 mm wide, midrib often red, margin slightly thickened, yellow, tapering to pungent mucro. Flowering stem grooved, c. 25 cm long x (2-)4-5 cm wide, upper nude part slender, lower part sheathed by leaves. Inflorescence a globose head of compound umbels; involucre bracts with thick sheaths up to 2 cm long and leaflike stipules and lamina. Primary umbels usually several, c. 2.5 cm diam., secondary umbels on almost filiform rays. Fruit of two mericarps; mericarps narrowly-elliptic to elliptic, 4.0-5.2 mm, with 5 even narrowly winged ribs, 4 smaller ribs on vittae, vittae largely obscured

## SIMILAR TAXA

Most similar to *A. spedenii* and *A. congesta*, but the leaves extending along the basal portion of the flowering stem is distinctive

## FLOWER COLOURS

Cream, White



Black Lake, Creek Saddle, Mt Burns.  
Photographer: John Smith-Dodsworth



Mt Burns, January. Photographer: John Smith-Dodsworth

## LIFE CYCLE

Winged schizocarps are dispersed primarily by wind (Thorsen et al., 2009).

## ETYMOLOGY

**aciphylla**: From the Latin acicula 'needle' and the Greek phyllum 'leaf', meaning needle-leaf.

## ATTRIBUTION

Description adapted from Allan (1961), Mark and Adams (1995), Webb and Simpson (2001) and Wilson (1996).

## REFERENCES AND FURTHER READING

- Allan, H.H. 1961. Flora of New Zealand. Government Printer, Wellington
- Mark, A.F; Adams, N.M. 1995. New Zealand alpine plants, 2nd Edition. Godwit Publishing, Auckland
- Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 2009 Vol. 11 No. 4 pp. 285-309
- Webb, C.J. & Simpson, M.J.A. 2001. Seeds of NZ gymnosperms and dicotyledons. Manuka Press, Christchurch

## MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/aciphylla-crosby-smithii/>