# Aciphylla pinnatifida

# **FAMILY**

Apiaceae

#### **AUTHORITY**

Aciphylla pinnatifida Petrie

#### **FLORA CATEGORY**

Vascular - Native

## **ENDEMIC TAXON**

Yes

#### **ENDEMIC GENUS**

No

#### **ENDEMIC FAMILY**

No

# STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

## **NVS CODE**

**ACIPIN** 

#### **CHROMOSOME NUMBER**

2n = 22

## **CURRENT CONSERVATION STATUS**

2017 | Not Threatened | Qualifiers: DP

# **PREVIOUS CONSERVATION STATUSES**

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

## WETLAND PLANT INDICATOR STATUS RATING

**OBL: Obligate Wetland** 

Almost always is a hydrophyte, rarely in uplands (non-wetlands).

## LIFE CYCLE

Winged schizocarps are dispersed primarily by wind (Thorsen et al., 2009).

## **ETYMOLOGY**

aciphylla: From the Latin acicula 'needle' and the Greek phyllum 'leaf', meaning needle-leaf.

pinnatifida: Pinnately divided, finely cut

# REFERENCES AND FURTHER READING

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 2009 Vol. 11 No. 4 pp. 285-309

## **MORE INFORMATION**

https://www.nzpcn.org.nz/flora/species/aciphylla-pinnatifida/





Mt Burns, January. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.



Hector Mountains. Photographer: John Barkla, Licence: CC BY.