

Anisotome imbricata var. imbricata

BIOSTATUS

Native – Endemic taxon

CURRENT CONSERVATION STATUS

2023 | Not Threatened

[Jump to previous conservation statuses](#)

CATEGORY

Vascular

STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

FLOWER COLOURS

Yellow

GENUS

Anisotome

FAMILY

Apiaceae

AUTHORITY

Anisotome imbricata (Hook.f.) Cockayne var. imbricata

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

LIFE CYCLE AND DISPERSAL

Winged mericarps are dispersed by wind (Thorsen et al., 2009).

WETLAND PLANT INDICATOR STATUS RATING

FACU: Facultative Upland

Occasionally is a hydrophyte but usually occurs in uplands (non-wetlands).

ETYMOLOGY

anisotome: Unequal sided

imbricata: From the Latin imbricatus 'tiled', refers to overlapping features such as leaves

NVS CODE

ANIIVI

PREVIOUS CONSERVATION STATUSES

2017 | Not Threatened

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

[Jump to current conservation status](#)



Old Woman Range. Photographer: John Barkla, Licence: CC BY.



Harris Mts. Photographer: John Barkla, Licence: CC BY.

REGIONAL CONSERVATION STATUSES

Otago: 2025 | Regionally Not Threatened | Qualifiers: TL Help

The regional threat classification system leverages off the national assessments in the NZTCS, providing information relevant for the regional context. Otago conservation status information is sourced from the "Conservation Status of Indigenous Vascular Plants in Otago, 2025" Jarvie S et al. (2025) report.

REFERENCES AND FURTHER READING

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309.

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/anisotome-imbricata-var-imbricata/>

PDF DATE

25 May 2026