Anisotome pilifera

FAMILY

Apiaceae

AUTHORITY

Anisotome pilifera (Hook.f.) Cockayne & Laing

FLORA CATEGORY

Vascular - Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

Νo

STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

NVS CODE

ANIPIL

CHROMOSOME NUMBER

2n = 22

CURRENT CONSERVATION STATUS

2017 | At Risk - Declining | Qualifiers: DP, PD

PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

FLOWER COLOURS

White

LIFE CYCLE

Winged mericarps are dispersed by wind (Thorsen et al., 2009).

ETYMOLOGY

anisotome: Unequal sided

pilifera: Hair-bearing; having soft short hairs

REFERENCES AND FURTHER READING

Thorsen MJ, Dickinson KJM, Seddon PJ. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285–309. https://doi.org/10.1016/j.ppees.2009.06.001.

MORE INFORMATION

https://www.nzpcn.org.nz/flora/species/anisotome-pilifera/





Hooker valley, December. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.



Mt Cook. Photographer: John Barkla, Licence: CC BY.