# Archeria racemosa

## **FAMILY**

Ericaceae

#### **AUTHORITY**

Archeria racemosa Hook.f.

#### **FLORA CATEGORY**

Vascular - Native

## **ENDEMIC TAXON**

Yes

#### **ENDEMIC GENUS**

Nο

#### **ENDEMIC FAMILY**

No

#### STRUCTURAL CLASS

Trees & Shrubs - Dicotyledons

#### **NVS CODE**

**ARCRAC** 

## **CHROMOSOME NUMBER**

2n = 24

#### **CURRENT CONSERVATION STATUS**

2017 | Not Threatened | Qualifiers: DP

#### **PREVIOUS CONSERVATION STATUSES**

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

## **BRIEF DESCRIPTION**

Bushy shrub of upland middle North Island forests. Leaves harsh, small, widest at the middle before narrowing to a pointed tip, with green lines underneath. Flowers reddish, in small spikes. Fruit a woody capsule with an obvious stalk at the tip, splits to release small dry seeds.

### **FLOWER COLOURS**

Red/Pink

# LIFE CYCLE

Seeds are dispersed by wind (Thorsen et al., 2009)

## **ETYMOLOGY**

archeria: After Archer, a Tasmanian botanical artist

racemosa: Raceme bearing

## WHERE TO BUY

Not commercially available

## REFERENCES AND FURTHER READING

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 2009 Vol. 11 No. 4 pp. 285-309

#### **MORE INFORMATION**





Mt Kaitarakihi, December. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.




Table Mt, Thames. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.

