

Azorella nitens

SYNONYMS

Schizeilema nitens (Petrie) Domin

FAMILY

Apiaceae

AUTHORITY

Azorella nitens Petrie

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

NVS CODE

SCHNIT

CHROMOSOME NUMBER

2n = 48

CURRENT CONSERVATION STATUS

2017 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

DISTRIBUTION

Endemic. North and South Islands from the Central Volcanic Plateau south.

HABITAT

Lowland to montane, margins of lakes, ponds, streams, and wet river terraces.

WETLAND PLANT INDICATOR STATUS RATING

FACW: Facultative Wetland

Usually is a hydrophyte but occasionally found in uplands (non-wetlands).

FLOWERING

December - January

FLOWER COLOURS

Cream, Violet/Purple

FRUITING

December - February

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/azorella-nitens/>



Flowering, in montane bog, upper Angus Burn, Waitutu, Southland. Photographer: Colin C. Ogle, Date taken: 01/12/1984, Licence: CC BY-NC.



Flowering in montane bog, upper Angus Burn, Waitutu, Southland. Photographer: Colin C. Ogle, Date taken: 01/12/1984, Licence: CC BY-NC.