

Blindia contecta

COMMON NAMES

moss

BIOSTATUS

Native

CURRENT CONSERVATION STATUS

2025 | At Risk – Uncommon | Qualifiers: DPS, DPT, RR

[Jump to previous conservation statuses](#)

CATEGORY

Non-vascular

STRUCTURAL CLASS

Mosses

DETAILED DESCRIPTION

Autoicous, robust, subaquatic moss forming loose, somewhat dull to lustrous tufts. Plants golden-green to yellow-brown above, olive-green to black below. Stems stiff, rigidly and sparsely branched, with outer stem cells thick-walled. Leaves rigid, loosely erect to slightly secund, occasionally with apices curved, sometimes contorted to helicoid when dry, stiffly erect to spreading, upper portions slightly reflexed when moist, 3.0-8.0 mm long; lamina long subulate from an ovate to oblong-ovate base with subulae hemispheric in transverse section. about 2x lamina length, acute to narrowly obtuse, some leaves deciduous, not decurrent; margins entire. Nerve more or less 100-200 microns wide, variable in width, but always strong and distinct. Upper cells 8-20 x 5-8 microns, rounded-quadrate to short-rectangular, thick-walled, covering the excurrent costa. Lower cells 30-140 x 5-10 microns, elongate to rectangular, thick-walled, non-porose, straight. Alar cells 20-65 x 10-35 microns, variable, slightly enlarged and hyaline to thicker-walled, orange, and inflated, sometimes auriculate, not reaching nerve. Perichaetial leaves similar to vegetative, more or less wider at base. perigonia lateral below perichaetia. Seta 1.0-2.0 mm long, thick, stout, straight not twisted. Capsules with urns 0.5-1.1 mm long, hemispheric to obovate when dry and moist, brown, deeply immersed. Exothecial cells 15-35 microns wide at mid-urn, decreasing in size towards rim, isodiametric to rounded, thick-walled, rim cells in 5-10 rows, oblate to quadrate, with thick, sinuous walls. Stomates not seen. Peristome of 16, lanceolate teeth, hyaline above, broken in upper portions in old capsules and blunt. Operculum conic rostrate, the beak more or less erect, long. Spores 32-40 microns.

DISTRIBUTION

Indigenous. Mainly subantarctic. recorded from Iles de Kerguelen, Heard, Campbell and the Auckland Islands. Also reported from South America (Chile) but those records are not substantiated by specimens

SUBSTRATE DETAILS

Saxicolous on wet rocks, usually beside waterfalls.

THREATS

Not Threatened. Listed because it qualifies as Range Restricted within the New Zealand part of its range.

GENUS

Blindia

FAMILY

Seligeriaceae

AUTHORITY

Blindia contecta (Hook.f. & Wilson) Mull.Hal.

SYNONYMS

Weissia contecta Hook.f. et Wilson

ENDEMIC TAXON

No



ENDEMIC GENUS

No

ENDEMIC FAMILY

No

FRUITING

Although fruit has been seen insufficient information exists to provide any details on the timing of fruiting

PREVIOUS CONSERVATION STATUSES

2014 | At Risk – Naturally Uncommon | Qualifiers: RR

2010 | At Risk – Naturally Uncommon | Qualifiers: RR

2005 | Range Restricted | Qualifiers: S?O

[Jump to current conservation status](#)

REFERENCES AND FURTHER READING

Bartlett, J.K.; Vitt, D. H. 1986: A survey of the species in the genus *Blindia* (Bryopsida, Seligeriaceae). New Zealand Journal of Botany 24: 203-246.

ATTRIBUTION

Fact Sheet Prepared for NZPCN by: P.J. de Lange (6 November 2007). Description adapted from Bartlett & Vitt (1986).

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/blindia-contecta/>

PDF DATE

25 May 2026