

Bulbinella angustifolia

SYNONYMS

Bulbinella hookeri (Hook.) Cheeseman var. *angustifolia* Cockayne et Laing

FAMILY

Asphodelaceae

AUTHORITY

Bulbinella angustifolia (Cockayne et Laing) L.B.Moore

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Herbs - Monocots

NVS CODE

BULANG

CHROMOSOME NUMBER

2n = 14

CURRENT CONSERVATION STATUS

2017 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

WETLAND PLANT INDICATOR STATUS RATING

FAC: Facultative

Commonly occurs as either a hydrophyte or non-hydrophyte (non-wetlands).

FLOWER COLOURS

Yellow

LIFE CYCLE

Winged seeds are dispersed by wind (Thorsen et al., 2009).

ETYMOLOGY

bulbinella: Little bulb

angustifolia: From the Latin *angustus* 'narrow, constricted' and *folius* 'leaf', meaning narrow-leaved

REFERENCES AND FURTHER READING

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 2009 Vol. 11 No. 4 pp. 285-309



Old Man range, December. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.



Old Woman Range. Photographer: John Barkla, Licence: CC BY.

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/bulbinella-angustifolia/>