

Calliergidium austro-stramineum

COMMON NAME

Moss

SYNONYMS

Hypnum austro-stramineum Müll.Hal.; *Warnstorfia laculosa* (Müll. Hal.)
Ochyra et Matter; *Hypnum laculosum* Müll. Hal.;

FAMILY

Amblystegiaceae

AUTHORITY

Calliergidium austro-stramineum (Müll.Hal.) E.B.Bartram

FLORA CATEGORY

Non-vascular – Native

ENDEMIC TAXON

No

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

CURRENT CONSERVATION STATUS

2009 | Data Deficient | Qualifiers: OL, SO

PREVIOUS CONSERVATION STATUS

2004 | Threatened – Nationally Critical

DISTRIBUTION

Indigenous. New Zealand: South Island (Cape Foulwind, near Westport).
Mainly found in Antarctica and subantarctic islands.

HABITAT

In coastal wetlands

THREATS

First recognised from New Zealand in 1984 when it was found near Cape Foulwind. Despite searching it has not been there again and it is possibly that local changes in farming and farm management in that area may have wiped it out (Glenny et al. 2011).

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/calliergidium-austro-stramineum/>



Calliergidium austro-stramineum.
Photographer: Bill Malcolm, Licence: All rights reserved.



Calliergidium austro-stramineum.
Photographer: Bill Malcolm, Licence: All rights reserved.