

Carex breviculmis

COMMON NAME

grassland sedge

SYNONYMS

None

FAMILY

Cyperaceae

AUTHORITY

Carex breviculmis R.Br.

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

No

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Sedges

NVS CODE

CARBRE

CHROMOSOME NUMBER

2n = c.64

CURRENT CONSERVATION STATUS

2012 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2009 | Not Threatened

2004 | Not Threatened

DISTRIBUTION

Indigenous, North and South Islands. Also Australia, New Guinea, Lord Howe and Norfolk Islands

HABITAT

Coastal to montane. Usually in open grassland, gum land scrub, clay pans, on rock stacks, and talus slopes and other similar sparsely vegetated sites.

FEATURES

Shortly rhizomatous; tufted sedge usually forming low-growing, close-packed, bright green patches. Culms hidden among leaves, usually 10-20 mm long; basal sheaths dull brown. Leaves much > culms, 1.5-3.0 mm wide, grass-like, recurved, channelled, margins exceedingly finely and closely scabrid almost throughout. Spikes 2-5, approximate, pale green; terminal spike male, usually sessile; remaining spikes female, occasionally with a few male flowers at the top, 6-9 mm long, ± pedunculate, clustered round base of male spike; subtending bracts leaf-like, very narrow-linear, the uppermost almost filiform, margins finely scabrid. Glumes > utricles, ovate, pale green, almost white, membranous, midrib green, stiff, thickened, produced to a stout, finely hispid awn. Utricles c.2.5 × 1.0 mm, biconvex or subtrigonus, fusiform, pale yellow-green, faintly many-nerved, pubescent all over; beak slightly narrowed, pale green, c.0.5 mm long, orifice ± truncate; stipe c.0.3 mm long, often much contracted. Stigmas 3. Nut c.1.5 mm long, obtusely trigonus, oblong, obovoid, light brown, surmounted by a minute, persistent, dilated style-base.



Whitianga, March. Photographer: John Smith-Dodsworth



Long Bay, Coromandel. Sept. Photographer: John Smith-Dodsworth

SIMILAR TAXA

Carex breviculmis is one of the earliest flowering New Zealand species and is immediately recognised by its palen-green, faintly nerved, pubescent utricles.

FLOWERING

August - December

FRUITING

October - May

LIFE CYCLE

Nuts surrounded by inflated utricles are dispersed by granivory and wind (Thorsen et al., 2009).

PROPAGATION TECHNIQUE

Easily grown from fresh seed and by division of established plants. In ideal conditions this species sometimes naturalises in urban areas but it is never invasive. Tolerant of a wide range of conditions except permanently waterlogged soils. However does best in full sun in a free draining soil.

ETYMOLOGY

carex: Latin name for a species of sedge, now applied to the whole group.

ATTRIBUTION

Fact Sheet prepared by P.J. de Lange (110 August 2006). Description adapted from Moore and Edgar (1970)

REFERENCES AND FURTHER READING

Moore, L.B.; Edgar, E. 1970: Flora of New Zealand. Vol. II. Government Printer, Wellington.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora.

Perspectives in Plant Ecology, Evolution and Systematics 2009 Vol. 11 No. 4 pp. 285-309

NZPCN FACT SHEET CITATION

Please cite as: de Lange, P.J. (Year at time of access): *Carex breviculmis* Fact Sheet (content continuously updated). New Zealand Plant Conservation Network. <https://www.nzpcn.org.nz/flora/species/carex-breviculmis/> (Date website was queried)

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/carex-breviculmis/>