

Carex pumila

COMMON NAME

sand sedge

SYNONYMS

Carex littorea Labill.; *Carex pumila* Thunb. subsp. *littorea* (Labill.) Kük.

FAMILY

Cyperaceae

AUTHORITY

Carex pumila Thunb.

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

No

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Sedges

NVS CODE

CARPUM

CHROMOSOME NUMBER

2n = 82

CURRENT CONSERVATION STATUS

2012 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2009 | Not Threatened

2004 | Not Threatened

DISTRIBUTION

Indigenous. New Zealand: North, South and Chatham Islands. Uncommon in parts of the South Island. Also recorded from Australia, Lord Howe Island, Chile, China, Japan and Korea.

HABITAT

Mostly coastal, rarely extending inland. A species of mobile sand dunes, sand flats and dune slacks (swales). Sometimes found fringing the sandy margins of coastal rivers and lagoons. Occasionally found as an urban lawn weed, especially in coastal settlements.



Waikawau bay, November. Photographer: John Smith-Dodsworth



Wainuiomata River mouth. Photographer: Jeremy Rolfe

FEATURES

Deep blue-green to glaucous, tufted sedge; tufts coarse, arising from a long, wiry creeping rhizome of c.2 mm diameter Culms mostly buried in sand, 50–300 mm long, terete, smooth, cream or light green, almost entirely enclosed by light brown or cream, occasionally red-brown leaf-sheaths. Leaves > culms, up to 400 mm long, 1.5–0 mm. wide, channelled, rigid, glaucous, curved and tapering to a fine point, margins mostly smooth. Spikes 3–8, ± approximate; terminal spike male, often long-pedunculate, very slender, often with 1–3 very small, occasionally partly female, spikes at the base; remaining spikes female, often male at the top, 10–35 x c.10 mm. Glumes c.½ length of utricles, rarely only slightly < utricles, ovate, acute, red-brown, with broad colourless hyaline margins, midrib very pale brown, thickened, usually produced to a short awn. Utricles 6.0–7.5 x 2.0–3.5 mm, biconvex to subtrigonus, ovoid, light brown, thick, corky, turgid, smooth or faintly nerved, narrowed to a bifid beak, 1.5–2.0 mm long, orifice membranous, crura faintly scabrid at tip. Stigmas 3. Nut 2.5–4.0 x 1.5–2.5 mm., trigonus, obovoid, light brown, shortly mucronate.

SIMILAR TAXA

None. The long, creeping, wiry rhizome, channeled, smooth-edged glaucous leaves and distinctly corky utricles are especially diagnostic.

FLOWERING

October - December

FRUITING

December - June

LIFE CYCLE

Spongy utricles dispersed by water and wind (Thorsen et al., 2009).

PROPAGATION TECHNIQUE

Easily grown from fresh seed and by division of established plants. Will grow in most soils and conditions but does best in a freely draining medium in full sun. Some wild populations rarely set seed, and some selection of wild forms is merited. An excellent and highly under-rated sand binder.

ETYMOLOGY

carex: Latin name for a species of sedge, now applied to the whole group.

pumila: Small

ATTRIBUTION

Fact Sheet prepared for NZPCN by P.J. de Lange (110 August 2006). Description adapted from Moore and Edgar (1970)

REFERENCES AND FURTHER READING

Moore, L.B.; Edgar, E. 1970: Flora of New Zealand. Vol. II. Government Printer, Wellington.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora.

Perspectives in Plant Ecology, Evolution and Systematics 2009 Vol. 11 No. 4 pp. 285-309

NZPCN FACT SHEET CITATION

Please cite as: de Lange, P.J. (Year at time of access): Carex pumila Fact Sheet (content continuously updated). New Zealand Plant Conservation Network. <https://www.nzpcn.org.nz/flora/species/carex-pumila/> (Date website was queried)

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/carex-pumila/>