

# Celmisia cockayneana

## COMMON NAMES

Cockayne's mountain daisy

## BIOSTATUS

Native – Endemic taxon

## CURRENT CONSERVATION STATUS

2023 | At Risk – Naturally Uncommon | Qualifiers: Sp, DPS, DPT

[Jump to previous conservation statuses](#)

## CATEGORY

Vascular

## STRUCTURAL CLASS

Herbs - Dicotyledonous composites

## FLOWER COLOURS

White, Yellow

## DETAILED DESCRIPTION

Stout subshrub; stems woody, c.100 mm diameter; branches stout, usually rather close-set, erect or ascending, clad in leaf-remains; living leaves rosulate at tips of branchlets. Lamina coriaceous,  $\pm 40\text{--}100 \times 10\text{--}15$  mm; narrowly obovate-spathulate to obovate-oblong, subacute to obtuse, usually minutely apiculate; upper surface glabrous, veins  $\pm$  distinct; lower surface clad in dense appressed white satiny tomentum, midrib dark, prominent; margins sinuate, distantly furnished with glandular denticles, narrowed cuneately to petiole then widening to nearly glabrous sheath c. 10 mm long. Scape rather slender to rather stout, 145-210 mm long, rather densely clad in viscid glandular hairs; bracts usually many; lower up to 30 mm long, leaf-like,  $\pm$  lanceolate, apiculate; upper linear-oblong. Capitula 20-40 mm diameter; involucre bracts numerous, lower bractiform,  $\pm 12$  mm long, narrow lanceolate, acuminate, densely clad in glandular hairs; upper linear-subulate. Ray-florets  $\pm 15$  mm long, white, limb narrow-oblong; disk-florets tubular, c. 5-6 mm long, yellow. Achenes narrow-cylindric,  $\pm$  compressed. 1.0-1.5 mm long; ribs very densely clad in short silky hairs. Pappus-hairs up to 6 mm long, off-white, hardly barbellate.

## SIMILAR TAXA

Perhaps closest to *Celmisia haastii* from which it is best distinguished by its longer leaves (400-100  $\times$  10-15 mm cf. 30-80  $\times$  10-250 mm), shorter sheath (10 mm cf. 20-30 mm) and extremely hairy rather than glabrous seeds. *Celmisia cockayneana* is endemic to Marlborough, while *C. haastii* ranges from about North Canterbury south.

## DISTRIBUTION

South Island. Endemic: Marlborough

## HABITAT

Montane to subalpine. Rupestral, in rock-crevices in shaded sites

## THREATS


A naturally uncommon species that does not appear to be actively threatened

## GENUS

Celmisia



Marlborough. Photographer: Melissa Hutchison, Date taken: 01/03/2021, Licence: CC BY-NC.



Habit; scape and bracts; involucre bracts; Seaward Kaikoura Range. Photographer: Jane Gosden, Date taken: 02/01/2014, Licence: CC BY-NC-SA.

## FAMILY

Asteraceae

## AUTHORITY

*Celmisia cockayneana* Petrie

## SYNONYMS

None

## ENDEMIC TAXON

Yes

## ENDEMIC GENUS

No

## ENDEMIC FAMILY

No

## FLOWERING

October - January

## FRUITING

November - March

## LIFE CYCLE AND DISPERSAL

Pappate cypselae are dispersed by wind (Thorsen et al., 2009).

## PROPAGATION TECHNIQUE

Unknown. Probably best grown from fresh seed and like many *Celmisia* this species will probably dislike high humidity and drying out.

## ETYMOLOGY

**celmisia:** Apparently named after Kelmis, one of Idaean Dactyls, a group of skilled mythical beings associated with the Mother Goddess Rhea in Greek mythology. Kelmis, whose name means 'casting', was a blacksmith and childhood friend of Zeus, son of Rhea and later king of the gods. In Ovid's 'Metamorphoses', Kelmis is described as offending Zeus who turned him into adamant so he was as hard as a tempered blade

**cockayneana:** Named after Leonard Cockayne FRS (7 April 1855 - 8 July 1934) who is regarded as New Zealand's greatest botanist and a founder of modern science in New Zealand

## MANAAKI WHENUA ONLINE INTERACTIVE KEY

[Key to \*Celmisia\* in New Zealand](#)

## NVS CODE

CELCOC

## CHROMOSOME NUMBER

2n = 108

## PREVIOUS CONSERVATION STATUSES

2017 | At Risk – Naturally Uncommon | Qualifiers: DP, Sp

2012 | At Risk – Naturally Uncommon | Qualifiers: Sp

2009 | At Risk – Naturally Uncommon

2004 | Range Restricted

[Jump to current conservation status](#)

## REFERENCES AND FURTHER READING

Allan H.H. 1961: Flora of New Zealand. Vol. I, Government Printer, Wellington.

Thorsen M.J.; Dickinson K.J.M.; Seddon P.J. 2009: Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11(4): 285-309.

## **ATTRIBUTION**

Description adapted from Allan (1961).

Some of this factsheet information is derived from [Flora of New Zealand Online](#) and is used under a [Creative Commons Attribution 3.0 New Zealand](#) licence.

## **MORE INFORMATION**

<https://www.nzpcn.org.nz/flora/species/celmisia-cockayneana/>

## **PDF DATE**

27 May 2026