

Celmisia parva

COMMON NAMES

mountain daisy

BIOSTATUS

Native – Endemic taxon

CURRENT CONSERVATION STATUS

2023 | Not Threatened | Qualifiers: DPS, DPT

[Jump to previous conservation statuses](#)

CATEGORY

Vascular

STRUCTURAL CLASS

Herbs - Dicotyledonous composites

FLOWER COLOURS

White

DETAILED DESCRIPTION

Small branching herb hugging ground in small patches; leaves spreading, rosulate at tips of branchlets. Lamina submembranous, $\pm 10\text{--}30 \times 3\text{--}10$ mm; linear- to oblong-lanceolate to narrow-oblong; upper surface glabrous or nearly so, midrib and usually main veins evident; lower surface densely clad in appressed soft to satiny white hairs, midrib usually distinct; apex subacute, apiculate; margins slightly recurved, minutely distantly denticulate, cuneately narrowed to slender petiole up to 20 mm long; sheath membranous, $\pm =$ lamina. Scape almost filiform, glabrous or with a few spreading hairs, $\pm 40\text{--}100$ mm long; bracts almost filiform, with widened bases, few (sometimes absent), lowermost up to c. 10 mm long. Capitula $\pm 10\text{--}15$ mm diameter; involucral bracts linear-subulate, acute to acuminate, apiculate, scarious, midrib distinct. Rays-florets up to c. 8 mm. long, white, linear, teeth very narrow-triangular; disk-florets 4-5 mm long, $\pm$ glandular at base, teeth triangular. Achenes narrow-cylindric, 1-2 mm long, glabrous or nearly so (in some forms with stiff hairs on obscure ribs). Pappus-hairs up to 4-5 mm long, very slender, sordid-white, very minutely barbellate

SIMILAR TAXA

Readily recognised by the slender, much branched, creeping and freely rooting stems. Plants form compact patches or mats and the leaves are arranged in small, rosulate clusters at the tips of the branchlets. *Celmisia parva* is mostly likely to be confused with *C. bellidoides* and *C. thomsonii* which share a similar growth habit. It is distinguished from both species by the leaf undersides which are densely clad in thick, white tomentum. The leaf undersides of *Celmisia bellidoides* and *C. thomsonii* are either glabrous or sparsely hairy (with the hairs distinctly bristle-like)

DISTRIBUTION

Endemic. South Island: westerly from North-West Nelson to about the Paparoa Range

HABITAT


Lowland to subalpine. Inhabiting poorly drained ground in shrubland, pakihi, grassland, herbfield, and around rock outcrops

GENUS

Celmisia

FAMILY

Asteraceae



Mt Herbert, Herbert Range, Kahurangi, 1400m elevation. Photographer: Rowan Hindmarsh-Walls, Date taken: 23/02/2020, Licence: CC BY-NC.



Leaf underside, Gunner Downs, Kahurangi, 1100m elevation. Photographer: Rowan Hindmarsh-Walls, Date taken: 02/05/2019, Licence: CC BY-NC.

AUTHORITY

Celmisia parva Kirk

SYNONYMS

None

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

FLOWERING

November - January

FRUITING

January - April

LIFE CYCLE AND DISPERSAL

Pappate cypselae are dispersed by wind (Thorsen et al., 2009).

PROPAGATION TECHNIQUE

Easily grown in a shaded site, planted within a permanently moist, free draining, acidic soil. Dislikes humidity and will not tolerate drying out. Best grown from fresh seed which should be sown immediately or stratified in a fridge or freezer for 1-3 months

WHERE TO BUY

Not Commercially Available

ETYMOLOGY

celmisia: Apparently named after Kelmis, one of Idaean Dactyls, a group of skilled mythical beings associated with the Mother Goddess Rhea in Greek mythology. Kelmis, whose name means 'casting', was a blacksmith and childhood friend of Zeus, son of Rhea and later king of the gods. In Ovid's 'Metamorphoses', Kelmis is described as offending Zeus who turned him into adamant so he was as hard as a tempered blade

MANAAKI WHENUA ONLINE INTERACTIVE KEY

[Key to Celmisia in New Zealand](#)

NVS CODE

CELPAR

CHROMOSOME NUMBER

2n = 108

PREVIOUS CONSERVATION STATUSES

2017 | Not Threatened | Qualifiers: DP

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

[Jump to current conservation status](#)

REFERENCES AND FURTHER READING

Allan HH. 1961. Flora of New Zealand, Volume I. Indigenous Tracheophyta: Psilopsida, Lycopsidea, Filicopsida, Gymnospermae, Dicotyledones. Government Printer, Wellington, NZ. 1085 p.

Thorsen MJ, Dickinson KJM, Seddon PJ. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285–309. <https://doi.org/10.1016/j.ppees.2009.06.001>.

ATTRIBUTION

Description adapted from Allan (1961)

Some of this factsheet information is derived from [Flora of New Zealand Online](#) and is used under a [Creative Commons Attribution 3.0 New Zealand](#) licence.

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/celmisia-parva/>

PDF DATE

09 July 2026