

Clianthus maximus

COMMON NAME

kākābeak, kōwhai ngutu-kākā, kākā beak

SYNONYMS

Clianthus puniceus var. *maximus* (Colenso) Kirk

FAMILY

Fabaceae

AUTHORITY

Clianthus maximus Colenso

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

Yes

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Trees & Shrubs - Dicotyledons

NVS CODE

CLIMAX

CHROMOSOME NUMBER

$2n = c.32$

CURRENT CONSERVATION STATUS

2017 | Threatened – Nationally Critical | Qualifiers: CD, RF, Sp

PREVIOUS CONSERVATION STATUSES

2012 | Threatened – Nationally Critical | Qualifiers: CD, RF

2009 | Threatened – Nationally Critical | Qualifiers: RF, CD

2004 | Threatened – Nationally Endangered

BRIEF DESCRIPTION

Rare (common in cultivation) small bushy shrub with drooping clusters of red sharp-tipped flowers. Leaves with many pairs of glossy leaflets arranged along a central stalk. Flowers 80mm long. Fruit a green pea-like pod that splits releasing the numerous hard small blotched seeds.

DISTRIBUTION

Endemic. North Island. Formerly on Great Barrier Island (Aotea Island). Still present in scattered populations from the East Coast of the North Island from Te Araroa south to northern Hawke's Bay and Te Urewera in the west.

HABITAT

Like the closely related *C. puniceus* (G.Don) Sol. ex Lindl. this species prefers early to mid successional shrubland habitats dominated by flax (*Phormium cookianum* Le Jolis, and *P. tenax* J.R.Forst et G.Forst) and tutu (*Coriaria arborea* Lindsay) in coastal, lowland and montane habitats. Often found along the tops and bases of unstable cliff faces or rock falls. Some habitats may not be natural, as this species, was said to have been grown by Maori, and many inland associations occur in the vicinity of former pa, kainga, gardens or canoe haul outs.



Roadside population ex Te Araroa.
Photographer: Bec Stanley, Licence: CC BY-SA.



Roadside population ex Te Araroa.
Photographer: Bec Stanley, Licence: CC BY-SA.

DETAILED DESCRIPTION

Shrub 1.5-6 m tall. Wood soft, stems “watery” easily broken. Branchlets semi-erect to weakly ascending, often decurved. Leaves 15-25 cm long, imparipinnate, with 15-30 pairs of subsessile leaflets. Leaflets, dark green, upper surface shiny (very glossy) 150-300 mm, linear-oblong, apex retuse or rounded. Inflorescences racemose, 15-30-flowered, located in leaf axils near branch apices. Flowers 80 mm, dark scarlet. Standard ovate-acuminate, 60 mm, dark scarlet, with a dark maroon (almost black blotch) and usually lacking stripes (these if present indistinct, often dotted; wings 30 mm long, lanceolate-falcate; keel 60 mm long, falcate-acuminate, dark scarlet. Pods long persistent, 80 mm, at first green and turgid, drying black and splitting open for entire length. Seeds numerous, c.1-1.5 mm diam, grey various striped or blotched with black, embedded in wispy grey, floccose hairs.

SIMILAR TAXA

Closely related to *C. puniceus* which has mat-green foliage, whose upper leaf surface is often slightly glaucous in colour. *C. maximus* leaves are dark green, and the upper leaf surface very glossy. *C. puniceus* flowers are slightly smaller, the standard is conspicuously striped white, and the spur blotched or striped white as well. *C. maximus* flowers are larger, very dark red, blotched dark purple-black near the base (very rarely with a few faint white stripes) and the spur is uniformly dark red.

FLOWERING

May flower throughout the year. However plants are mostly found in flower between August and January

FLOWER COLOURS

Black, Red/Pink

FRUITING

Seed pods may be present at any time of the year

LIFE CYCLE

Seeds are wind dispersed (Thorsen et al., 2009).

PROPAGATION TECHNIQUE

Easily grown from seed, semi-hardwood cuttings, and stem layerings. Plants tend to be short-lived in cultivation (2-4 years), and benefit from hard pruning after flowering. Kaka beak is vulnerable to a range of common garden pests which include slugs and snails, it can be severely defoliated, by these animals, and young plants may be killed completely. Caterpillars, mites - which cause witches brooms, and various fungal diseases will also kill plants. To combat these problems grow plants in fertile, well drained, sunny sites free from surrounding shrubs. Despite its northerly distribution, kaka beak often does best in Southland, and is very tolerant of snowfall, and light frosts.

THREATS

Though more widespread than *C. puniceus* this species is now at a very serious risk of extinction. Only 153 mature plants are known from the wild and at all sites they are threatened by a diverse range of introduced browsing animals, diseases, and natural senescence. Many populations occur in low scrub where they are threatened by fire, weed control operations, natural succession, and the unstable, erosion prone nature of the habitats in which they grow.

Overview of recovery work [here](#).

ETYMOLOGY

clianthus: From Greek ‘kleios’ glory and ‘anthos’ flower, meaning glory flower

WHERE TO BUY

Now common in cultivation and widely sold. But prior to the early 1990s this species was virtually unknown in cultivation, most if not all horticultural stock then available being referable to *C. puniceus* s.s.

EXTRA INFORMATION

[New Zealand Geographic article](#) about kākābeak restoration work.

ATTRIBUTION

Fact sheet prepared for NZPCN by P.J. de Lange 1 October 2003. Description adapted from Heenan (2000).

REFERENCES AND FURTHER READING

Heenan, P.B. 2000: *Clianthus* (Fabaceae) in New Zealand: a reappraisal of Colenso's taxonomy. *New Zealand Journal of Botany* 38(3): 361-371

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

NZPCN FACT SHEET CITATION

Please cite as: de Lange, P.J. (Year at time of access): *Clianthus maximus* Fact Sheet (content continuously updated). New Zealand Plant Conservation Network. <https://www.nzpcn.org.nz/flora/species/clianthus-maximus/> (Date website was queried)

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/clianthus-maximus/>