

Coprosma rubra

SYNONYMS

Coprosma rubra var. *pendula* (Colenso) Kirk

FAMILY

Rubiaceae

AUTHORITY

Coprosma rubra Petrie

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Trees & Shrubs - Dicotyledons

NVS CODE

COPRUB

CHROMOSOME NUMBER

2n = 44

CURRENT CONSERVATION STATUS

2012 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2009 | Data Deficient

2004 | Not Threatened

BRIEF DESCRIPTION

Rare small bushy small-leaved wide-angled shrub with reddish bark and fuzzy twigs. Leaves thin, usually 10–15mm long but up to 2.5cm long, gradually narrowing to a winged stalk that has tiny hairs (lens needed), small black tip on stem between leaf bases, leaf margin sometimes hairy. Fruit yellowish white.

DISTRIBUTION

Endemic. North and South Islands: Mostly eastern. Sporadic in Northland around the upper Wairoa River and Pipwai, more common from the Hawkes Bay and Taihape south but often absent or very uncommon over large parts of its range.

HABITAT

Lowland to montane. Usually in riparian forest and shrubland, especially on alluvial soils or those derived from calcareous parent materials.



At Mataroa, Taihape, May. Photographer: John Smith-Dodsworth



Mataroa, Taihape, May. Photographer: John Smith-Dodsworth

FEATURES

Shrub up to 4 m tall; branches rather slender, filiramulate, weakly divaricating. Bark reddish brown; branchlets finely pubescent. Leaves membranous; petioles 4-6 mm. long more or less pubescent-ciliolate, winged. Stipules small, triangular, pubescent, usually terminated by single denticle. Lamina glabrous or nearly so, broad-ovate to oblong, obtuse, sometimes mucronulate, narrowed to subtruncate base; 10-25 × 6-15 mm; margins sometimes ciliolate. Reticulations usually evident on both surfaces, at least when leaf fresh. Male flowers 6-1 on short branchlets; calyx 0; corolla funnelform, lobes about = tube, ovate, acute. Female flowers solitary or 2-3 together on short branchlets; calyx-teeth linear-triangular, obtuse, c.³/₄ length of corolla-tube, corolla lobes > tube, narrow-linear. Drupe yellowish white, oblong, 4-6 mm long.

SIMILAR TAXA

Coprosma rubra is superficially similar to and most often confused with *C. rotundifolia*. *Coprosma rubra* differs from the much more widespread *C. rotundifolia* by its smaller leaves (10-18 x 6-13 mm cf. 15-25 x 10-20 mm in *C. rotundifolia*, female flowers whose calyces have conspicuous linear rather than inconspicuous minute teeth, and oblong, yellowish-white to white rather than red or white globose drupes. Both species may be found growing together.

FLOWERING

September - January

FLOWER COLOURS

Green

FRUITING

February - August

LIFE CYCLE

Fleshy drupes are dispersed by frugivory (Thorsen et al., 2009).

PROPAGATION TECHNIQUE

Easy from fresh seed and semi-hardwood cuttings. An attractive shrub to small tree that does well in dappled light. Should be planted in a free draining but moist, fertile soil.

THREATS

Previously listed as Data Deficient because in some parts of its range it appears to be in decline and in other areas it is extremely uncommon.

ETYMOLOGY

coprosma: From the Greek kopros 'dung' and osme 'smell', referring to the foul smell of the species, literally 'dung smell'

rubra: Red

WHERE TO BUY

Not commercially available

ATTRIBUTION

Description adapted from Allan (1961).

REFERENCES AND FURTHER READING

Allan, H.H. 1961: *Flora of New Zealand. Volume I*. Government Printer, Wellington.

Thorsen, M.J.; Dickinson, K.J.M.; Seddon, P.J. 2009: Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/coprosma-rubra/>