

Cyperus ustulatus

COMMON NAME

coastal cutty grass, giant umbrella sedge, cyperus

SYNONYMS

Mariscus ustulatus (A.Rich.) C.B.Clarke, *Cyperus ustulatus* f. *grandispiculosus* Carse ex Kük.

FAMILY

Cyperaceae

AUTHORITY

Cyperus ustulatus A.Rich.

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Sedges

CURRENT CONSERVATION STATUS

2012 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2009 | Not Threatened

2004 | Not Threatened

DISTRIBUTION

Abundant in the North Island and northern South Island, west to Fiordland, and not threatened. Naturally uncommon at its eastern South Island limit, where it is known only from Tai Tapu, Motukarara, Banks Peninsula and the Rakaia River mouth. Also on the Chatham Islands, where it is not very common.

HABITAT

Coastal to lowland sites in open ground. Tolerant of a wide range of habitats and conditions but evidently preferring wetland margins, seepages, streamsides, lagoon and estuary margins.

FEATURES

Robust, 0.6–2.0 m tall, with leaves crowded at base of culms. Culms triquetrous, glabrous, striated. Leaves 0.6–1.2 m long, 8–15 mm wide, lamina coriaceous, strongly keeled, multitubular with numerous septa prominent on the abaxial side, margins and keel very sharply and minutely serrulate; sheath brown. Inflorescence a terminal umbel 40–140 mm long, of 6–12 unequal rays; rays usually unbranched, rarely with secondary branches at base; involucre of numerous leaf-like bracts very much > inflorescence Spikelets 8–13 mm long, numerous, dark brown or yellow-brown, crowded on each ray into a dense spike 35–70 mm. long. Glumes 5–20 in each spikelet, ovate-oblong, obtuse or mucronate, hard, smooth and shining, keeled, red-brown with white nerves, 2(-3) lowermost and 1–2 uppermost glumes smaller, empty, the remainder fertile. Stamens with persistent filaments. Nut 1.5–2 × c.0.5 mm., c.½ length of glume, linear-oblong, trigonous, brown.



Cyperus ustulatus f. *ustulatus* plant, Coromandel, February. Photographer: John Smith-Dodsworth



Cyperus ustulatus f. *ustulatus* inflorescences, Coromandel, February. Photographer: John Smith-Dodsworth

SIMILAR TAXA

Distinguished from *C. insularis* by its light green leaves (*C. insularis* has grey-green leaves and involucre bracts). It has a brown adaxial midrib whereas *C. insularis* is grey-green.

Its leaf sheaths are brown to red-brown and are rigid and hard (whereas in *C. insularis* they are light pink to purple-pink and flexible and soft).

The culm in *C. ustulatus* is green, distal 5-15 mm is always brown to red-brown and long persistent after flowering. The culm of *C. insularis* is green, rarely pale brown in distal part and is upright at flowering but weakens and collapses at seed fall after flowering. There are other floral characters which also distinguish these species, for these and other details see Heenan & de Lange (N.Z.J.Bot. 43: 351-359 (2005) - link provided below).

FLOWERING

July - December

FRUITING

July - April

LIFE CYCLE

Nuts are dispersed by water, granivory and attachment (Thorsen et al., 2009).

PROPAGATION TECHNIQUE

Easily grown from fresh seed, and often self sows in gardens. A quite attractive plant now prove popular in cultivation. However it should be planted with caution, the leaf, keel and culm margins are very sharp and can cause very deep cuts.

ETYMOLOGY

cyperus: From the ancient Greek name for sedge, kypeiros

ustulatus: Burned or scorched

TAXONOMIC NOTE

In many populations seed is not formed due to the endemic smut fungus *Bauerago (Ustilago) gardneri* (Vankey) Vankey which infects the spikes. Infected spikes can be recognised by the abundance of pollen-like yellow powder—the fruiting bodies of the smut fungus—within the spikelets. Plants bearing this smut were formerly thought to be a genetic form and were named *Cyperus ustulatus* f. *grandispiculosus* Carse ex Kük.

WHERE TO BUY

Over the last decade this species has become very popular and is now widely sold by most nurseries either as *Cyperus ustulatus* or *Mariscus ustulatus*.

ATTRIBUTION

Description adapted from Moore and Edgar (1970)

REFERENCES AND FURTHER READING

Heenan and de Lange (2005). N.Z.J.Bot. 43: 351-359: *Cyperus insularis* (Cyperaceae), a new species of sedge from northern New Zealand.

Moore, L.B.; Edgar, E. 1970: Flora of New Zealand. Vol. II. Government Printer, Wellington.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora.

Perspectives in Plant Ecology, Evolution and Systematics 11: 285-309

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/cyperus-ustulatus/>