

# Erioderma leylandii

## BIOSTATUS

Native

## CURRENT CONSERVATION STATUS

2018 | At Risk – Naturally Uncommon | Qualifiers: SO, Sp

## CATEGORY

Lichen

## STRUCTURAL CLASS

Lichens - Foliose

## LICHEN SUBSTRATE

Corticolous (bark, wood)

## SUBSTRATE DETAILS

Corticolous (twigs and branches)

## GENUS

Erioderma

## FAMILY

Pannariaceae

## AUTHORITY

Erioderma leylandii (Taylor) Müll.Arg.

## SYNONYMS

Erioderma glaucescens sensu D.J.Galloway (1985)

## ENDEMIC TAXON

No

## ENDEMIC GENUS

No

## ENDEMIC FAMILY

No

## ETYMOLOGY

**erioderma**: woolly skin

## REFERENCES AND FURTHER READING

Galloway D.J. 1985: *Flora of New Zealand: Lichens*. Wellington: PD Hasselberg, Government Printer. 662 pp.

Galloway D.J. 2007: *Flora of New Zealand: Lichens, including lichen-forming and lichenicolous fungi*. 2nd edition. Lincoln, Manaaki Whenua Press. 2261 pp.

Taylor M., Bieleski R.L. and Allan H.H. 2002: Meanings and origins of botanical names of New Zealand plants. *Auckland Botanical Society Bulletin* 26.

## MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/erioderma-leylandii/>

## PDF DATE

25 May 2026



Corticolous on manuka, Wainui Auckland.  
Photographer: Marley Ford, Date taken:  
23/02/2022, Licence: CC BY-NC.



Corticolous on manuka, Wainui Auckland.  
Photographer: Marley Ford, Date taken:  
23/02/2022, Licence: CC BY-NC.