# **Exocarpos bidwillii**

# **FAMILY**

Santalaceae

#### **AUTHORITY**

Exocarpos bidwillii Hook.f.

#### **FLORA CATEGORY**

Vascular - Native

# **ENDEMIC TAXON**

Yes

#### **ENDEMIC GENUS**

Nο

#### **ENDEMIC FAMILY**

Νo

#### STRUCTURAL CLASS

Trees & Shrubs - Dicotyledons

#### **NVS CODE**

**EXOBID** 

## **CURRENT CONSERVATION STATUS**

2017 | Not Threatened

# **PREVIOUS CONSERVATION STATUSES**

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

# **BRIEF DESCRIPTION**

Bushy small shrub with many yellowish leafless twigs inhabiting mountain areas of the South Island. Twigs with small dark triangular scales and pale grooves. New twigs narrower, orange. Flowers small, inconspicuous, reddish. Fruit reddish, oval, with hard seed at tip.

# LIFE CYCLE

Arillate nutlets are dispersed by frugivory (Thorsen et al., 2009).

#### **ETYMOLOGY**

exocarpos: Outside fruit

bidwillii: Named after the botanist - John Carne Bidwill (born 1815 and died 16 March 1853)

# REFERENCES AND FURTHER READING

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 11: 285-309

#### **MORE INFORMATION**

https://www.nzpcn.org.nz/flora/species/exocarpos-bidwillii/





Cobb ridge, January. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.



Cobb ridge, January. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.