

Heterodermia casarettiana

BIOSTATUS

Native

CURRENT CONSERVATION STATUS

2018 | At Risk – Naturally Uncommon | Qualifiers: DP, SO

CATEGORY

Lichen

STRUCTURAL CLASS

Lichens - Foliose

LICHEN SUBSTRATE

Corticolous (bark, wood)

SUBSTRATE DETAILS

Corticolous

GENUS

Heterodermia

FAMILY

Physciaceae

AUTHORITY

Heterodermia casarettianum (A.Massal.)

SYNONYMS

Anaptychia casarettiana A.Massal., Polyblastidium casarettianum (A.Massal.) Kalb.

ENDEMIC TAXON

No

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

ETYMOLOGY

heterodermia: From the Greek heteros (other, different) and derma (a skin or hide), in reference to the presence or absence of a lower cortex

REFERENCES AND FURTHER READING

Galloway D.J. 2007: *Flora of New Zealand: Lichens, including lichen-forming and lichenicolous fungi*. 2nd edition. Lincoln, Manaaki Whenua Press. 2261 pp.

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/heterodermia-casarettiana/>

PDF DATE

27 May 2026



Corticolous lichen on margins of alluvial forest remnant, Long Bush Gisborne. Photographer: Marley Ford, Date taken: 24/01/2022, Licence: CC BY-NC.



Corticolous lichen on margins of alluvial forest remnant, Long Bush Gisborne. Photographer: Marley Ford, Date taken: 24/01/2022, Licence: CC BY-NC.