

Homalanthus polyandrus

COMMON NAMES

Kermadec poplar

BIOSTATUS

Native – Endemic taxon

CURRENT CONSERVATION STATUS

2023 | At Risk – Naturally Uncommon | Qualifiers: IE

[Jump to previous conservation statuses](#)

CATEGORY

Vascular

STRUCTURAL CLASS

Trees & Shrubs - Dicotyledons

SIMPLIFIED DESCRIPTION

Broad small tree bearing thin wide triangular leaves inhabiting the Kermadec Islands. Twigs bleed white sap. Leaves 5-10cm long, as wide as long, on long stalk to 10cm long. Flowers in a green long spike. Fruit purple with white lines that split to reveal wrinkled yellow seeds.

DETAILED DESCRIPTION

Small, glabrous tree up to 10 m tall. Branches and branchlets slender, rather brittle, terete in cross-section, initially slightly ribbed, leaf-scars prominent. Leaves adaxially dark or bright-green to yellow-green above, sometimes suffused with red (especially new growth), abaxially subglaucous, membranous, stipulate, stipules } 20 mm long, caducous; juvenile lamina up to 300 mm diameter, broadly ovate, truncate or cordate at base, rarely perfoliate, adult lamina 30-200 diameter, usually rather broad-ovate to subrhombic, abruptly narrowed to acute or acuminate apex, cuneate to truncate at base; petioles > or } = to blade. Inflorescences terminal on new growth, racemose. Racemes slender up to 200 mm long, usually with a few solitary long-pedicellate ♂ flowers below or at base and many solitary short-pedicellate ♀ flowers above (racemes occasionally unisexual); each flower subtended by a caducous, glandular bract and 2 prominent glands at the base of the pedicel Male flowers bearing c.30-35 close-set stamens; perianth segments 1. Female flowers tri-locular with 3 styles, stigma capitate. Fruit } smooth, coriaceous, 3-angled, reddish, 10-12 mm diameter. Seed elliptic, oblong-elliptic or more irregular, terete or somewhat compressed, (2.3-)3.0 - 4.8 mm long, almost entirely covered with a yellowish aril.

SIMILAR TAXA

None in the wild. However has often been confused with the naturalised and very weedy *Homalanthus populifolius* Graham. From that species it differs by the male flowers which are solitary rather than clusters along the raceme, and female flowers which are tri rather than bi-locular and which have 3 rather than 2 styles.

DISTRIBUTION

Endemic. Kermadec Island group, Raoul and Macauley Islands

HABITAT

A successional species of coastal scrub, forest and upland forest, where it usually forms the understorey or is a sporadic emergent. Sometimes, as on old slip scars, it may form the main canopy.



Macauley Island. Photographer: John Barkla, Licence: CC BY.



Homalanthus polyandrus. Photographer: Bec Stanley, Licence: CC BY-SA.

THREATS

Formerly seriously threatened with extinction due to feral goats and regarded as extinct on Macauley Island, this species has made a spectacular recovery following the successful goat eradication on Raoul Island (the last goat was shot there in 1986). *Homalanthus* is now widespread and no longer regarded as threatened. In 2006 it was rediscovered on Macauley Island - though whether this was the original tree seen by Cheeseman or a new arrival is unclear. The species is now listed only because it is still a naturally uncommon, range restricted endemic.

GENUS

Homalanthus

FAMILY

Euphorbiaceae

AUTHORITY

Homalanthus polyandrus (Hook.f. ex Müll.Arg.) G.Nicholson

SYNONYMS

Homalanthus polyandrus (Hook.f.) Cheeseman comb. et nom. superf.; *Carumbium polyandrum* Hook.f. ex Müll.Arg.; *Carumbium polyandrum* Hook.f. nom. inval.

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

FLOWERING

Throughout the year

FRUITING

Throughout the year

LIFE CYCLE AND DISPERSAL

Arrilate seeds are dispersed by frugivory (Thorsen et al., 2009).

PROPAGATION TECHNIQUE

Easily grown from fresh seed and cuttings, and often naturalises where it has been planted. However, extremely cold sensitive and will not tolerate even a slight frost. Best grown in a warm, sheltered site.

NVS CODE

HOMPOL

CHROMOSOME NUMBER

$2n = 64$

PREVIOUS CONSERVATION STATUSES

2017 | At Risk – Naturally Uncommon | Qualifiers: IE

2012 | At Risk – Naturally Uncommon | Qualifiers: IE, RR

2009 | At Risk – Naturally Uncommon | Qualifiers: OL

2004 | Range Restricted

[Jump to current conservation status](#)

REFERENCES AND FURTHER READING

Gardner, R. 1999. *Homalanthus* (Euphorbiaceae) in New Zealand and its fruit. Auckland Botanical Society Journal, 54: 6-7

Müller Argoviensis, J. 1864 (3. September): Neue Euphorbiaceen des Herbarium Hooker in Kew, auszugsweise vorläufig mitgeteilt aus dem Manuscript für De Candolles Prodrömus. *Flora, oder Allgemeine Botanische Zeitung* 47(28): 433.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

ATTRIBUTION

For the authority citation see Mueller (1864)

NZPCN FACT SHEET CITATION

Please cite as: de Lange, P.J. (Year at time of access): *Homalanthus polyandrus* Fact Sheet (content continuously updated). New Zealand Plant Conservation Network.

<https://www.nzpcn.org.nz/flora/species/homalanthus-polyandrus/> (Date website was queried)

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/homalanthus-polyandrus/>

PDF DATE

07 June 2026