

Inula conyzae

COMMON NAMES

Ploughman's spikenard

BIOSTATUS

Exotic

CONSERVATION STATUS

Not applicable

CATEGORY

Vascular

STRUCTURAL CLASS

Herbs - Dicotyledonous composites

FLOWER COLOURS

Yellow

DETAILED DESCRIPTION

Erect, shortly rhizomatous perennial up to c. 1m tall. Stems densely hairy, branched above to form infl. Lower cauline leaves sparsely to moderately hairy on upper surface, tomentose on lower, elliptic, long-cuneate, acute, finely denticulate, up to c. 15 x 14 cm; upper lvs similar but smaller and short-cuneate or obtuse at base. Capita 8-12 mm diam., many in corymbs. Outer involucral bracts sparsely hairy and ciliate, herbaceous at apex, subulate, 2-5 mm long; inner bracts sparsely hairy and ciliate, membranous, linear-lanceolate, 8-12 mm long. Ray florets numerous; ligules inconspicuous, yellow, c. 1 mm long. Disc yellow. Mature achenes not seen, sparsely hairy. (Webb et al 1988)

GENUS

Inula

FAMILY

Asteraceae

AUTHORITY

Inula conyzae (Griess.) Meikle

ENDEMIC FAMILY

No

FLOWERING

January, February

YEAR NATURALISED

1988

ORIGIN

Europe to W. Asia, N. Africa

ETYMOLOGY

inula: Believed to be a corruption of helenium, *Inula helenium* (Elecampane) being the *Inula campana* of medieval Latin (Johnson and Smith, 1986).

REFERENCES AND FURTHER READING

Johnson, A. T. and Smith, H. A (1986). *Plant Names Simplified: Their pronunciation, derivation and meaning.* Landsman Bookshop Ltd: Buckenhill, UK.

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/inula-conyzae/>



