

# Kelleria croizatii

## BIOSTATUS

Native – Endemic taxon

## CURRENT CONSERVATION STATUS

2023 | Not Threatened

[Jump to previous conservation statuses](#)

## CATEGORY

Vascular

## STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

## FLOWER COLOURS

White

## GENUS

Kelleria

## FAMILY

Thymelaeaceae

## AUTHORITY

Kelleria croizatii Heads

## ENDEMIC TAXON

Yes

## ENDEMIC GENUS

No

## ENDEMIC FAMILY

No

## LIFE CYCLE AND DISPERSAL

Achenes are possibly wind dispersed (Thorsen et al., 2009).

## NVS CODE

KELCRO

## CHROMOSOME NUMBER

2n = 18

## PREVIOUS CONSERVATION STATUSES

2017 | Not Threatened

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

[Jump to current conservation status](#)

## REGIONAL CONSERVATION STATUSES

Otago: 2025 | Regionally At Risk – Regionally Naturally Uncommon | Qualifiers: DPR, DPS, DPT, NS, NStr, Sp Help  
The regional threat classification system leverages off the national assessments in the NZTCS, providing information relevant for the regional context. Otago conservation status information is sourced from the "[Conservation Status of Indigenous Vascular Plants in Otago, 2025](#)" Jarvie S et al. (2025) report.

## REFERENCES AND FURTHER READING

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 11: 285-309



Temple basin, January. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.



Mount Burns, Fiordland. Photographer: Anne Humburg, Date taken: 13/12/2008, Licence: All rights reserved.

**MORE INFORMATION**

<https://www.nzpcn.org.nz/flora/species/kelleria-croizatii/>

**PDF DATE**

25 May 2026