

Koeleria antarctica

BIOSTATUS

Native – Endemic taxon

CURRENT CONSERVATION STATUS

2023 | At Risk – Declining | Qualifiers: Sp

[Jump to previous conservation statuses](#)



CATEGORY

Vascular

STRUCTURAL CLASS

Grasses

DETAILED DESCRIPTION

Densely tufted, dull green to grey-green grass up to 400 mm tall, with rather rigid leaves usually reaching or sometimes overtopping the dense spike-like panicles. **Branching** extravaginal at plant base, sometimes intravaginal above. **Leaf-sheath** to 40 mm, very minutely pubescent or with extremely minute appressed hairs between ribs. **Ligule** 0.2–0.6 mm, truncate, erose often sparsely minutely ciliate, undersides sometimes minutely prickly-toothed. **Leaf-blade** 35–220 × 1.5–4 mm, flat or inrolled, undersides mostly smooth, often scabrid near the long-narrow tapering apex; upper surface ribbed with sparse to dense minute hairs and prickly-teeth on ribs, hairs slightly longer near ligule; margins minutely prickly-toothed. **Culm** 40–300 mm long, internodes glabrous, occasionally bearing a few minute prickly-teeth below panicle. **Panicle** 20–150 × 6–25 mm, compact, oblong, very dense, rarely with some lower branches and pedicels smooth or with sparse minute prickly-teeth or rarely minute hairs. **Spikelets** 4.5–7.0 mm, greyish green or brownish amber. **Glumes** unequal, hyaline, keel thickened with sparse long prickly-teeth on upper $\frac{1}{2}$; lower   length of upper, linear-lanceolate, upper slightly < spikelet, elliptic; margins with prickly-teeth near acuminate to almost mucronate apex. **Lemmas** 4.0–6.2 mm, bidentate to shortly bicuspid, papillose, prickly-toothed above and on keel; awn 3.5–6.0, initially straight becoming recurved, insertion in upper $\frac{1}{4}$ of lemma. **Palea** with minute prickly-teeth on keels and usually on margins. **Callus** hairs to 0.5 mm. **Rachilla hairs** to 2.5 mm. **Lodicules** c. 1 mm, glabrous. **Anthers** 1.0–1.7 mm. **Ovary** 0.8 mm; stigma-styles to 1.3 mm. **Seed** 2.5–3.0 × 0.6 mm.

SIMILAR TAXA

Distinguished from the 12 other indigenous species by the dense, compact, oblong panicle (rarely with spreading lower branches) and by the stiff, rigidly erect, flat grey-green to dull green leaves with the undersides glabrous and the upper minutely, though distinctly scabrid.

DISTRIBUTION

Endemic. North Island south from the southern Egmont Coast, on Kapiti Island and western and southern Wellington Coasts to Cape Palliser. In the South Island at Cape Farewell and along the outer Marlborough Sounds and adjacent islands.

HABITAT

Coastal, on gravel, sand and bluffs.

THREATS

Still quite common in parts of its range but there has been a range contraction around Wellington due to the spread of weeds into its habitat, and also because of greywacke quarrying along the south coast.



In cultivation, ex Makara. Dec 2004.
Photographer: Barbara Mitcalfe, Licence: CC BY-NC.



In cultivation, ex Makara. Dec 2004.
Photographer: Barbara Mitcalfe, Licence: CC BY-NC.

GENUS

Koeleria

FAMILY

Poaceae

AUTHORITY

Koeleria antarctica (G.Forst.) Barberá, Quintanar, Soreng et P.M.Peterson

SYNONYMS

Aira antarctica G.Forst., *Avena antarctica* (G.Forst.) Roem. et Schult., *Trisetum saxeticolum* Cockayne et Allan;
Trisetum antarcticum (G.Forst.) Trin.

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

FLOWERING

November–February

FRUITING

December–May

PROPAGATION TECHNIQUE

Easy from fresh seed and the division of whole plants. An attractive grass for an open spot in well drained soil. Once established rather drought tolerant.

WHERE TO BUY

Not commercially available

ETYMOLOGY

koeleria: Named after George Ludwig Koeler (1765-1807), 18th century German botanist. Author of a work on the grasses of Germany and France, *Descriptio graminum in Gallia et Germania* (1802).

antarctica: Antarctic

MANAAKI WHENUA ONLINE INTERACTIVE KEY

[Key to the grasses of New Zealand](#)

NVS CODE

TRIAN T

CHROMOSOME NUMBER

2n = 28

PREVIOUS CONSERVATION STATUSES

2017 | At Risk – Declining | Qualifiers: Sp

2012 | At Risk – Declining | Qualifiers: Sp

2009 | At Risk – Declining

2004 | Gradual Decline

[Jump to current conservation status](#)

REFERENCES AND FURTHER READING

Edgar E, Connor HE. 2000. Flora of New Zealand. Vol. V. Grasses. Manaaki Whenua Press, Christchurch, NZ. 650 p.

ATTRIBUTION

Description modified from Edgar and Connor (2000).

Some of this factsheet information is derived from [Flora of New Zealand Online](#) and is used under a [Creative Commons Attribution 3.0 New Zealand](#) licence.

NZPCN FACT SHEET CITATION

Please cite as: de Lange, P.J. (Year at time of access): *Koeleria antarctica* Fact Sheet (content continuously updated). New Zealand Plant Conservation Network. <https://www.nzpcn.org.nz/flora/species/koeleria-antarctica/> (Date website was queried)

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/koeleria-antarctica/>

PDF DATE

25 May 2026