

Leptopteris hymenophylloides

COMMON NAME

crêpe fern, single crêpe fern, heruheru

SYNONYMS

Todea hymenophylloides A.Rich.; *Todea marginata* Colenso; *Todea pellucida* Carmich. ex Grev. et Hook.; *Leptopteris marginata* (Colenso) C.Chr.; *Osmunda hymenophylloides* (A.Rich.) J.B.Armstr.

FAMILY

Osmundaceae

AUTHORITY

Leptopteris hymenophylloides (A.Rich.) C.Presl

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Ferns

NVS CODE

LEPHYM

CHROMOSOME NUMBER

2n = 44

CURRENT CONSERVATION STATUS

2012 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2009 | Not Threatened

2004 | Not Threatened

DISTRIBUTION

Endemic. New Zealand: North, South, Stewart and Chatham Islands from North Cape (Whiriwhiri Stream) south. Common throughout, though more abundant in the northern part of its range

HABITAT

Lowland to montane forest. Rarely in gumland scrub and coastal forest. Usually found along stream sides, and on damp banks, occasional on forested ridge lines. Once established this species is remarkably tolerant of drought and high light conditions and so it can be found growing as a persistent relict in disturbed forest or in areas cleared by wind throw.

FEATURES

Trunks up to 0.1 m tall. Stipes 0.15-0.5 m long, pale brown, sparsely hairy, with ear like lobes at base. Frond delicate, membranous, translucent, laminae ± deltoid, 3-pinnate, 0.2-1.0m long, 150-350 mm wide, dark green to light emerald green, sparsely hairy, veins free. Primary pinnae in 20-30 pairs, widely spaced, basal ones 60-120 mm long. Ultimate segments linear, flattened in plane of frond. Sporangia scattered on underside of pinnae (not in discrete sori), though tending to be more abundant toward frond centre. Description modified from Brownsey & Smith-Dodsworth 2000.



Wairere falls. Photographer: John Smith-Dodsworth



Wairere falls. Photographer: John Smith-Dodsworth

SIMILAR TAXA

Leptopteris hymenophylloides differs from the closely related *L. superba* by its longer stipe, triangular frond, longer and broader pinnae, and by its ultimate lamina segments flattened in one plane. Where both species meet they commonly hybridise to produce the hybrid known as *Leptopteris ×intermedia* (André) Brownsey.

FLOWERING

N.A.

FLOWER COLOURS

No flowers

FRUITING

N.A.

LIFE CYCLE

Minute spores are wind dispersed (Thorsen et al., 2009).

PROPAGATION TECHNIQUE

Easily grown in a damp shaded site, in a free draining but moist, humus enriched soil.

ETYMOLOGY

leptopteris: Thin fern; from the Greek leptos and pteris

hymenophylloides: Resembling *Hymenophyllum*

ATTRIBUTION

Fact sheet prepared for NZPCN by P.J. de Lange 10 March 2011. Description modified from Brownsey & Smith-Dodsworth 2000.

REFERENCES AND FURTHER READING

Brownsey, P.J.; Smith-Dodsworth, J.C. 2000: *New Zealand Ferns and Allied Plants*. Auckland, David Bateman
Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

NZPCN FACT SHEET CITATION

Please cite as: de Lange, P.J. (Year at time of access): *Leptopteris hymenophylloides* Fact Sheet (content continuously updated). New Zealand Plant Conservation Network.

<https://www.nzpcn.org.nz/flora/species/leptopteris-hymenophylloides/> (Date website was queried)

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/leptopteris-hymenophylloides/>