

# Lindsaea trichomanoides

## SYNONYMS

*Lindsaea cuneata* (G. Forst.) C.Chr.; *Lindsaea cuneata* var. *lessonii* (Bory) C.Chr.; *Lindsaea trichomanoides* var. *lessoni* (Bory) Hook.f.; *Lindsaea lessonii* Bory in Duperrey; *Schizoloma trichomanoides* (Dryand.) Kuhn; *Adiantum cuneatum* G.Forst.; *Adiantum trichomanoides* (Dryand.) Poir.

## FAMILY

Lindsaeaceae

## AUTHORITY

*Lindsaea trichomanoides* Dryand.

## FLORA CATEGORY

Vascular – Native

## ENDEMIC TAXON

No

## ENDEMIC GENUS

No

## ENDEMIC FAMILY

No

## STRUCTURAL CLASS

Ferns

## NVS CODE

LINTRI

## CHROMOSOME NUMBER

2n = 84, 86

## CURRENT CONSERVATION STATUS

2017 | Not Threatened | Qualifiers: SO

## PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

## DISTRIBUTION

Indigenous. New Zealand: North Island (common from Te Paki south and west throughout, less common in the east), South Island (confined to the Marlborough Sounds and West Coast), Stewart Island/Rakiura. Also Australia.

## HABITAT

Coastal to montane. Usually in closed forest (sometimes in scrub) often growing in relatively open sites on the forest floor, in leaf litter or amongst *Carex* spp. (especially *C. banksiana*, *C. subviridis* and *C. lectissima*).



Haywards Scenic Reserve, Lower Hutt.  
Photographer: Jeremy R. Rolfe, Date taken:  
26/12/2004, Licence: CC BY.



Stokes Valley, Lower Hutt. Photographer:  
Jeremy R. Rolfe, Date taken: 14/07/2001,  
Licence: CC BY.

## DETAILED DESCRIPTION

Terrestrial, tufted fern. **Rhizome** short- to long-creeping, to 1.5 mm diameter; scales narrowly triangular, reddish brown. **Fronds** monomorphic, erect and stiff. **Stipes** clustered or more widely spaced, shorter to longer than lamina, glossy, reddish brown to dark brown. **Lamina** 100–280 × 30–68 mm, 1–2-pinnate-pinnatifid, lanceolate to triangular-oblong, herbaceous. **Pinnae** oblong-deltoid or oblong-lanceolate, variously dissected, 4–10 mm long; pinnules flabellate-spathulate, asymmetric, often erose. **Upper pinnae and pinnules** gradually reduced and confluent. **Veins** free, simple to 3× forked. **Sori** mostly on 1–4 vein endings; indusium 0.3–0.5 mm wide, subentire. **Spores** trilete. (Description from Kramer & McCarthy (1998)).

## SIMILAR TAXA

Easily distinguished from the two other New Zealand species of *Lindsaea* by the terrestrial habit (not rheophytic like *L. viridis*), by the broader, bipinnate fronds with round ended ultimate segments.

## FLOWERING

Not applicable—spore producing

## FLOWER COLOURS

No flowers

## FRUITING

Not applicable—spore producing

## PROPAGATION TECHNIQUE

Difficult—should not be removed from the wild

## ETYMOLOGY

**lindsaea**: Named after John Lindsay, 19th century British surgeon who discovered fern spore

**trichomanoides**: Fern-like

## Where To Buy

Not commercially available.

## ATTRIBUTION

Fact sheet prepared for NZPCN by P.J. de Lange (Updated 4 May 2011)

## REFERENCES AND FURTHER READING

Kramer KU, McCarthy PM. 1998. Lindsaeaceae. *Flora of Australia* 48, *Ferns Gymnosperms and allied groups*: 228–240. ABRIS/CSIRO Victoria, Australia.

## NZPCN FACT SHEET CITATION

Please cite as: de Lange, P.J. (Year at time of access): *Lindsaea trichomanoides* Fact Sheet (content continuously updated). New Zealand Plant Conservation Network.

<https://www.nzpcn.org.nz/flora/species/lindsaea-trichomanoides/> (Date website was queried)

## MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/lindsaea-trichomanoides/>