

# Myosotis angustata

## BIOSTATUS

Native – Endemic taxon

## CURRENT CONSERVATION STATUS

2023 | Threatened – Nationally Critical | Qualifiers: DPT, RR

[Jump to previous conservation statuses](#)

## CATEGORY

Vascular

## STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

## FLOWER COLOURS

White

## DETAILED DESCRIPTION

Tufted, perennial, grey-green, rosette forming herb. Rosette-leaves 20 x 4 mm, linear-spathulate, apex sub acute. Leaf-hairs coarse, of two types, small and long, these usually appressed, overlapping. Lateral branches few, ascending to erect, up to 150 mm long in fruiting specimens, internodes < leaves. Stem-leaves 15-20 x 2-4 mm, linear to narrow-oblong, apex sub acute, hairs as for rosette-leaves. Inflorescence simple, sometimes branched, compact terminal cyme, 10-20 mm long when in flower, elongating to 50 mm when in fruit. Calyx 5-8 mm, lobes cut to half calyx length, sub acute; hairs on lobes long, straight and numerous, towards base changing to long and hooked with an underlayer of short, straight, appressed hairs. Corolla white, 7 mm diam., tube 5-6(-8-10) mm, cylindric, corolla lobes 2.5 x 2.5 mm, broad-oblong; filaments fixed between small scales, about length of anthers.

## SIMILAR TAXA

Could be confused with *M. traversii* Hook.f. (var. *traversii* and var. *cantabrica* L.B.Moore). These have their anthers partly included within the corolla-tube, while the hairs on the undersurfaces of the rosette leaves are retrorse, and the long hairs of the calyces are all or almost all hooked. *M. traversii* var. *traversii* is known from scree in Kahurangi National Park.

## DISTRIBUTION

Endemic. South Island, North West Nelson, Kahurangi National Park, Mt Arthur.

## HABITAT

Alpine. On shaded marble outcrops and associated talus.

## THREATS

Apparently confined to single location in Kahurangi National Park. At this location although the species appears to be genuinely uncommon, indications are that the population is stable. The species qualifies for Nationally Critical because the total area of occupancy is < 1 ha.

## GENUS

*Myosotis*

## FAMILY

Boraginaceae

## AUTHORITY

*Myosotis angustata* Cheeseman

## SYNONYMS

None



A *Myosotis angustata* plant. Photographer: Shannel Courtney, Licence: CC BY-NC.

## ENDEMIC TAXON

Yes

## ENDEMIC GENUS

No

## ENDEMIC FAMILY

No

## FLOWERING

December - February

## FRUITING

January - April

## PROPAGATION TECHNIQUE

Difficult. Should not be removed from the wild.

## CULTIVATION

A few plants are held by specialist growers. Very difficult to maintain in cultivation.

## ETYMOLOGY

**myosotis:** Mouse-eared

**angustata:** From the Latin angustus 'narrowed'

## PREVIOUS CONSERVATION STATUSES

2017 | Threatened – Nationally Critical | Qualifiers: DP, RR

2012 | Threatened – Nationally Critical | Qualifiers: RR, St

2009 | Threatened – Nationally Critical | Qualifiers: St, OL

2004 | Threatened – Nationally Critical

[Jump to current conservation status](#)

## REFERENCES AND FURTHER READING

Allan, H.H. 1961: Flora of New Zealand. Vol. I, Government Printer, Wellington

## ATTRIBUTION

Fact Sheet prepared for NZPCN by P.J. de Lange 1 February 2008. Description based on Allan (1961).

Some of this factsheet information is derived from [Flora of New Zealand Online](#) and is used under a [Creative Commons Attribution 3.0 New Zealand licence](#).

## MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/myosotis-angustata/>

## PDF DATE

25 May 2026