

Pimelea sericeovillosa subsp. sericeovillosa

COMMON NAMES

pimelea

BIOSTATUS

Native – Endemic taxon

CURRENT CONSERVATION STATUS

2023 | At Risk – Declining | Qualifiers: DPS, DPT

[Jump to previous conservation statuses](#)

CATEGORY

Vascular

STRUCTURAL CLASS

Trees & Shrubs - Dicotyledons

SIMPLIFIED DESCRIPTION

Low growing sprawling shrub with hairy twigs bearing pairs of hairy leaves; hairy white flowers and orange fruit; inhabiting the northern South Island. Leaf dimensions not known, hairs curved, denser on underside.

FLOWER COLOURS

White

DETAILED DESCRIPTION

A gynodioecious, low, much-branched, loose cushion-forming shrublet to 50 × 250 mm. with brown, clustered, appressed, leafy, densely villous young branchlets, and with older, gnarled, leafless stems often visible (these often leafless and glabrous, dark brown to black, often partly buried by windblown silt or sand). **Internode** length 0.3–0.6 mm. **Branching** mainly sympodial and radiating from a stout main stem up to 15 mm in diameter. **Node buttresses** lunate, dark brown, masked by hairs on young stems, not prominent on leafless branchlets. **Leaves** decussate, ascending, imbricate, sessile or with very short petioles (0.2 mm); lamina medium to pale green, elliptic to oblong, 2.2–4.0 × 1.0–1.3 mm, adaxially concave, mid-vein not evident, abaxial surface very densely covered with curled (sometimes straight), yellowish or dull-white, moderately long hairs; adaxial surface less densely hairy, sometimes glabrate (the youngest leaves have more or less dense adaxial vestiture), obtuse, base cuneate, stomata on both adaxial and abaxial surfaces. **Inflorescences** terminal, with 1 or 2, sometimes 3, flowers. **Involucral bracts** 4, the same size as, or slightly wider than adjacent leaves (2.3 × 1.5 mm). **Receptacle** usually with abundant long hairs. **Flowers** 1–2 per inflorescence, white, on very short (0.1 mm) pedicels, very hairy outside, inside hairless; female tube 2.5 mm long, ovary portion 2 mm, calyx lobes 1.0–1.2 × 0.5 mm; hermaphrodite tube 3–4 mm long, ovary portion 2 mm, calyx lobes 1.5 × 0.8–1.0 mm. **Anther** dehiscence introrse. **Ovary** with dense short hairs on summit, less dense to half way down. **Fruits** ovoid, fleshy, pale orange 2.5–3.0 × 1.8–2.0 mm, seeds narrow-ovoid 2.0–2.2 × 1.0–1.3 mm.

SIMILAR TAXA


Pimelea sericeovillosa subsp. *sericeovillosa* is distinguished from the other two subspecies by its loosely and openly branched cushion-forming habit, dull-green colouration, leaf indumentum comprised of curled (rarely straight hairs), and by its restriction to Marlborough and North Canterbury, where it inhabits mountain tops (only occasionally extending down to valley floors).

DISTRIBUTION

Endemic. South Island: Marlborough and North Canterbury.

HABITAT

Subalpine to alpine. Usually in open fell-field on mountain tops, ridge crests and in open stonefields. Occasionally found at lower levels in short short-tussock grassland.



Rachael Range, Molesworth. Photographer: Gillian M. Crowcroft, Licence: All rights reserved.

THREATS

Burrows (2011) rates the status of all three *Pimelea sericeovillosa* subspecies as “precarious to varying degrees”. However, no data is provided to show this, nor is the New Zealand Threat Classification system (NZTCS) (Townsend et al. 2008) used for his assessments; rather Burrows consistently confuses the NZTCS with listings provided by Hitchmough et al. (2007). Burrows (2011) also uses the ranking of Molloy et al. (2001), which is superseded by Townsend et al. (2008). *Pimelea sericeovillosa* is declining (de Lange et al. 2018), although the causes of decline are not specified in the NZ Threat Classification System database (accessed December 2023). The paucity of flowering may be a factor.

GENUS

Pimelea

FAMILY

Thymelaeaceae

AUTHORITY

Pimelea sericeovillosa Hook.f. subsp. *sericeovillosa*

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

FLOWERING

September–January

FRUITING

December–May

PROPAGATION TECHNIQUE

Easily grown from semi-hardwood cuttings but difficult to maintain in cultivation. Prefers a moist free-draining soil, planted in full sun. Dislikes humidity, shade and poor drainage.

WHERE TO BUY

Not commercially available

ETYMOLOGY

pimelea: from Greek pimelē, meaning “lard” or “soft fat,” presumably referring to the oily seeds or fleshy cotyledons.

NVS CODE

PIMSSS

CHROMOSOME NUMBER

2n = 36

PREVIOUS CONSERVATION STATUSES

2017 | At Risk – Declining | Qualifiers: DP

2012 | At Risk – Declining | Qualifiers: DP

2009 | Data Deficient

2004 | Not Threatened

[Jump to current conservation status](#)

REFERENCES AND FURTHER READING

- Burrows, C.J. 2011: Genus *Pimelea* (Thymelaeaceae) in New Zealand 5. The taxonomic treatment of five endemic species with both adaxial and abaxial leaf hair. *New Zealand Journal of Botany* 49: 367-412.
- de Lange, P.J.; Norton, D.A.; Courtney, S.P.; Heenan, P.B.; Barkla, J.W.; Cameron, E.K.; Hitchmough, R.; Townsend, A.J. 2009: Threatened and uncommon plants of New Zealand (2008 revision). *New Zealand Journal of Botany* 47: 61-96.
- de Lange PJ, Rolfe JR, Barkla JW, Courtney SP, Champion PD, Perrie LR, Beadel SM, Ford KA, Breitwieser I, Schonberger I, Hindmarsh-Walls R, Heenan PB, Ladley K. 2018. Conservation status of New Zealand indigenous vascular plants, 2017. *New Zealand Threat Classification Series* 22. Department of Conservation, Wellington, NZ. 82 p. <https://www.doc.govt.nz/globalassets/documents/science-and-technical/nztcs22entire.pdf>
- Hitchmough R, Bull L, Cromarty P (compilers) 2007: New Zealand Threat Classification System lists - 2005. Wellington: Department of Conservation, Scientific Publishing.
- Molloy J, Bell B, Clout M, de Lange P, Gibbs G, Given D, Norton D, Smith N, Stephens T. 2002: Classifying species according to threat of extinction – a system for New Zealand. *Threatened Species Occasional Publication* 22. Department of Conservation, Wellington, NZ. 26 p. <https://www.doc.govt.nz/globalassets/documents/science-and-technical/tsop22.pdf>.
- NZTCS. 2013 - present. *Pimelea sericeovillosa* Hook.f. subsp. *sericeovillosa* in NZ Threat Classification System database, <https://nztcs.org.nz>. Accessed 30 December 2023.
- Townsend AJ, de Lange PJ, Norton DA, Molloy J, Miskelly C, Duffy C. 2008: New Zealand Threat Classification manual. Department of Conservation, Wellington, NZ. 35 p. <https://www.doc.govt.nz/globalassets/documents/science-and-technical/sap244.pdf>

ATTRIBUTION

Fact sheet prepared for NZPCN by P.J. de Lange (29 September 2011). Description adapted from Burrows (2011).

NZPCN FACT SHEET CITATION

Please cite as: de Lange, P.J. (Year at time of access): *Pimelea sericeovillosa* subsp. *sericeovillosa* Fact Sheet (content continuously updated). New Zealand Plant Conservation Network. <https://www.nzpcn.org.nz/flora/species/pimelea-sericeovillosa-subsp-sericeovillosa/> (Date website was queried)

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/pimelea-sericeovillosa-subsp-sericeovillosa/>

PDF DATE

27 May 2026