

Pittosporum eugenioides

COMMON NAME

Tarata, lemonwood

SYNONYMS

Pittosporum elegans Raoul, *P. microcarpum* Putt.

FAMILY

Pittosporaceae

AUTHORITY

Pittosporum eugenioides A.Cunn.

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Dicotyledonous Trees & Shrubs

NVS CODE

PITEUG

CHROMOSOME NUMBER

$2n = 24$

CURRENT CONSERVATION STATUS

2012 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2009 | Not Threatened

2004 | Not Threatened

BRIEF DESCRIPTION

Tree bearing light green wavy-edge oval leaves and with a contrasting pale green central vein, dense sprays of yellow flowers and small dry fruits. Leaf buds covered in dark-edged scales. Fruit pointed, 5-6mm long which splits into two to show a papery layer covering black sticky seeds.

DISTRIBUTION

Endemic. Common in the North and South Islands.

HABITAT

Common tree of regenerating and mature forest in coastal to montane situations.



Female flowers. Oct 2006. Photographer: Peter de Lange



Female flowers. Oct 2006. Photographer: Peter de Lange

FEATURES

Gynodioecious tree up to 12 m tall but usually much less. Trunk 0.6-1 m diam, stout, clad in persistent pale-grey bark, branches numerous, erect then spreading. Leaf buds sticky, resinous. Leaves borne on slender petioles 10-20 mm long, alternate, 50-100(-150) x 25-40 mm, yellow-green, green, more or less blotched and mottled with paler green or yellow-green (sometimes white), somewhat leathery, glossy, smelling strongly when crushed of ivy or resin, elliptic to elliptic-oblong, apex acute to subacute; leaf margin undulate (very rarely not so), midrib pale green. Inflorescences terminal, numerous, subcorymbose compound umbels. Flowers pale yellow to yellow, very fragrant. Peduncles 10-20 mm, pedicels 5 mm, both sparsely hairy. Sepals 2 mm, ovate to narrow-ovate, pale caducous. Petals 5, 5-7 mm long, narrow-oblong. Capsules 2-valved (rarely 3), 5-6 mm, ovoid to elliptic, caducous, seeds immersed in dark yellow viscid pulp, whole structure covered in long persistent papery endocarp.

SIMILAR TAXA

Well marked from all other indigenous and exotic *Pittosporum* spp. in New Zealand, by the yellow-green, mottled lanceolate leaves with undulating margins, and pale-yellow to yellow flowers arranged in subcorymbose compound umbels.

FLOWERING

October - December

FLOWER COLOURS

Yellow

FRUITING

October - January

PROPAGATION TECHNIQUE

Easy from fresh seed. Can be grown from semi-hardwood cuttings.

ETYMOLOGY

pittosporum: Pitch seed

eugenioides: Like *Eugenia*, a species of myrtle

WHERE TO BUY

Commonly cultivated and available from most garden centres, and then often as a variegated form rather than the pure plant. Occasionally seen for sale in European and English garden centres.

ATTRIBUTION

Fact sheet prepared for NZPCN by P.J. de Lange 30 August 2006. Description adapted from Cooper (1956).

REFERENCES AND FURTHER READING

Cooper, R.C. 1956: The Australian and New Zealand species of *Pittosporum*. *Annals of the Missouri Botanical Garden* 43: 87-188

Gardner, R. 1999. Notes towards an excursion Flora. *Pittosporum eugenioides* as a wild plant. *Auckland Botanical Society Journal*, 54, 1

CITATION

Please cite as: de Lange, P.J. (Year at time of access): *Pittosporum eugenioides* Fact Sheet (content continuously updated). New Zealand Plant Conservation Network.

<https://www.nzpcn.org.nz/flora/species/pittosporum-eugenioides/> (Date website was queried)

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/pittosporum-eugenioides/>