

# Potamogeton crispus

## COMMON NAME

curly pondweed

## FAMILY

Potamogetonaceae

## AUTHORITY

Potamogeton crispus L.

## FLORA CATEGORY

Vascular – Exotic

## STRUCTURAL CLASS

Herbs - Monocots

## NVS CODE

POTCRI

## CONSERVATION STATUS

Not applicable

## BRIEF DESCRIPTION

Submerged aquatic plant that has green to red/brown leaves arranged alternately on stems. the leaves are ca 10mm in width and up to 80 mm long with distinct wavy edges (hence the name 'crispus'). Centre veins of the leaves are normally reddish.

## DISTRIBUTION

Widely naturalised throughout New Zealand, rare in West Coast Region

## HABITAT

Moderately fast flowing to still water bodies.

## WETLAND PLANT INDICATOR STATUS RATING

OBL: Obligate Wetland

Almost always is a hydrophyte, rarely in uplands (non-wetlands).

## DETAILED DESCRIPTION


Entirely submerged, bottom-rooting, normally perennial. Slender rhizomes and roots. *P. crispus* has long branched stems, that feels slightly flattened (as opposed to round or cylindrical), with submerged leaves that are arranged alternately on the stem. The leaves are green to reddish brown, slightly translucent, about 10mm in width, and up to 80mm long. Leaf margins are markedly wavy and crimped, with fine teeth near the tip. Centre veins of the leaves are normally reddish. The inflorescence spike has few flowers. Flowers small, green, on spikes up to 1 cm long, 4 flat dark seeds. The fruit (achenes) are large (4mm) and thick, with a curved beak and a toothed (rounded toothed) dorsal keel. Turions or winter buds of hard scales formed by shortened, thickened leaves with broad and strongly toothed bases.

## SIMILAR TAXA

Potamogeton ochreatus and *P. cheesemanii*. *P. ochreatus* has a straight edged leaf and a blunt leaf tip, whereas curled pondweed has a rounded leaf tip with crimped edges. It can be distinguished from *P. cheesemanii* by the submerged leaves, which are narrower and much more crimped in curled pondweed.

## FLOWERING

November, December, January



Fruit of *Potamogeton crispus*. Photographer: Rohan Wells, Date taken: 22/05/2012, Licence: All rights reserved.



Leaves of *Potamogeton crispus*. Photographer: Rohan Wells, Date taken: 19/07/2012, Licence: All rights reserved.

## FLOWER COLOURS

Cream, Green

## FRUITING

Autumn

## LIFE CYCLE

Spread by seeds, stem fragments, rhizomes, or turions. Birds disperse seed, fragments, rhizomes and turions dispersed by contaminated machinery and water movement.

## YEAR NATURALISED

1940

## ORIGIN

Native to Europe, Asia, Africa and Australia.

## REASON FOR INTRODUCTION

Ornamental pond and aquarium plant

## CONTROL TECHNIQUES

Not usually controlled in New Zealand, but may be controlled manually, or mechanically.

## TOLERANCES

Tolerates brackish, flowing or still water, and grows to 10 m depth.

## ETYMOLOGY

**potamogeton**: River dweller

## ATTRIBUTION

Factsheet prepared by Paul Champion and Deborah Hofstra (NIWA).

## REFERENCES AND FURTHER READING

Champion et al (2012). Freshwater Pests of New Zealand. NIWA publication.

<http://www.niwa.co.nz/freshwater-and-estuaries/management-tools/identification-guides-and-fact-sheets/freshwater-pest-species>.

Johnson PN, Brooke PA (1989). Wetland plants in New Zealand. DSIR Field Guide, DSIR Publishing, Wellington. 319pp.

Coffey BT, Clayton JS (1988). New Zealand water plants: a guide to plants found in New Zealand freshwaters. Ruakura Agricultural Centre. 65pp.

Aston, H (1977). Aquatic plants of Australia. Melbourne University Press, 367pp.

## MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/potamogeton-crispus/>