

Pseudotsuga menziesii

COMMON NAME

Douglas fir, Oregon pine

FAMILY

Pinaceae

AUTHORITY

Pseudotsuga menziesii (Mirb.) Franco

FLORA CATEGORY

Vascular – Exotic

STRUCTURAL CLASS

Trees & Shrubs - Gymnosperms

NVS CODE

PSEMEN

CONSERVATION STATUS

Not applicable

HABITAT

Terrestrial. A plant of lowland, montane and subalpine habitats growing in sites with low-moderate fertility (Timmins & MacKenzie 1995). A plant of scrub and forest margin communities, shrublands, tussockland and light wells in forest (Timmins & MacKenzie 1995).

WETLAND PLANT INDICATOR STATUS RATING

FACU: Facultative Upland

Occasionally is a hydrophyte but usually occurs in uplands (non-wetlands).

FEATURES

Very large resinous evergreen tree. Bark thick, reddish-brown beneath, rough and furrowed when mature. Branches in irregular whorls, horizontal; branchlets usually drooping. Shoots light to dark brown, ridged, with short hairs. Winter buds to 1 cm long, shining purplish-brown, narrow and sharp. Leaves 15–38 x 1–2 mm, in 2 ranks, needle-like, whitish beneath, edges often rolled, orange-scented when crushed. Male cones 12–20 mm long, catkin-like. Female cones 5–10 cm long, cylindrical, papery, downward-pointing; seed scales large and broad; bract scales longer and thinner than seed scales, 3-pointed, centre point longest.

SIMILAR TAXA

Pseudotsuga can be separated from *Tsuga* as it lacks the persistent leaf base. *Pseudotsuga macrocarpa* is known to be cultivated in New Zealand and has cones 10 – 18 cm long. *P. sinensis* & *P. japonica* is also cultivated in specialist collections; both are from Asia .

FLOWER COLOURS

No flowers

LIFE CYCLE

Perennial. Seedlings require the presence of a mycorrhizal fungi with seedlings establishing most readily on well lit sheltered sites where there is no competition from other vegetation. The plant seldom spreads onto land where vegetation cover is dense. Seed is produced at a rate of 20 000 seeds per mature tree annually (Timmins & MacKenzie 1995).

Seed is dispersed in autumn and winter by wind and gravity and remains viable for a few weeks to many years.



Pseudotsuga menziesii. Photographer: John Smith-Dodsworth



Pseudotsuga menziesii. Photographer: John Smith-Dodsworth

PROPAGATION TECHNIQUE

Seed

YEAR NATURALISED

1925

ORIGIN

N. America

REASON FOR INTRODUCTION

Forestry.

TOLERANCES

The plant is tolerant to shade, more so than other conifers and intolerant of drought. The plant is frost tolerant when it is higher than 2m. Physical damage results in regrowth if green foliage remains intact, recovers from browsing (Timmins & MacKenzie 1995). The plant recovers after fire if there is an adjacent seed source.

EXTRA INFORMATION

The National Wilding Conifer Control Programme team at Biosecurity New Zealand, a branch of Ministry for Primary Industries, has produced this wilding conifer quick ID guide.

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/pseudotsuga-menziesii/>