

Rhopalostylis baueri

COMMON NAMES

Kermadec nīkau (N.Z. only)

BIOSTATUS

Native

CURRENT CONSERVATION STATUS

2023 | At Risk – Naturally Uncommon | Qualifiers: RR, SO

[Jump to previous conservation statuses](#)

CATEGORY

Vascular

STRUCTURAL CLASS

Trees & Shrubs - Monocotyledons

SIMPLIFIED DESCRIPTION

Palm to 15 m tall with a ringed trunk and 3 m-long leaves inhabiting Raoul Island in the Kermadec Islands. Leaves with multiple long narrow leaflets (to 1 m long) closely-spaced along the central stem. Flowers pinkish, in multiple spikes at the top of trunk. Fruit red.

FLOWER COLOURS

Red/Pink, White

DETAILED DESCRIPTION

Trunk up to 15 m tall, and 0.35 m wide, green to dark green fading to grey with age between rather closely spaced leaf-scars; crown-shaft to 0.80 m long, smooth and green, slightly bulging. **Leaf** to 3 m long, usually dark green; leaflets closely set, ascending sharply, up to 1 m long and 40 mm wide. **Spathes** c. 380 × 150 mm, between pink and yellow, smooth, falling as first flowers open. Inflorescence: Ultimate branches of inflorescence to 300 mm long, c. 15 mm diameter with buds on, at first pale cream-coloured; flower-buds tightly packed. **Sepals** short, the longest one in male rarely = petals. **Petals** white or pink. **Fruit** c. 13 × 12 mm, subglobose to globose brick-red. **Seed** almost spherical, tightly invested in pale fawn endocarp which is marked by several vascular strands curving ± obliquely from hilum and only a few running longitudinally; hilum a broad band from the slightly flattened chalazal area to a pronounced papilla beside the micropyle.

SIMILAR TAXA

Rhopalostylis sapida is a very variable species. It is best distinguished from *R. baueri* by its elliptic-oblong, smaller (mostly up to 7 mm diameter) fruit. While many New Zealand populations of *R. sapida* have narrower leaflets, some, especially those in the Hauraki Gulf and on the Chatham Islands have equally broad leaflets and fruit dimensions comparable to *R. baueri*. *Rhopalostylis sapida* is absent from the indigenous range of *R. baueri*.

DISTRIBUTION

Indigenous. In New Zealand known only from Raoul Island (Kermadec Islands group). Also on Norfolk Island, its type locality.

HABITAT

Abundant in both dry and wet forest types on Raoul Island where it sometimes is the main canopy dominant.



Raoul Island. Photographer: Bec Stanley, Licence: CC BY-SA.



Rhopalostylis baueri. Photographer: John Barkla, Licence: CC BY.

THREATS

In the New Zealand part of its range it does not face any specific threats. However, it is known only from Raoul Island, and so qualifies as At Risk – Naturally Uncommon. On Norfolk, while common it occupies a very reduced range in the centre of that island. There the species is threatened to some extent by rats which eat the fruit and germinating seedlings.

GENUS

Rhopalostylis

FAMILY

Areaceae

AUTHORITY

Rhopalostylis baueri (Seem.) H.Wendl. et Drude

SYNONYMS

Areca sapida Endl., Kentia sapida (Endl.) Mart., Kentia baueri Seem., Eora baueri (H.Wendl. et Drude) O.F.Cook., Rhopalostylis cheesemanii Beccari, R. baueri var. cheesemanii (Beccari) Sykes

ENDEMIC TAXON

No

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

FLOWERING

December–January

FRUITING

December–January

PROPAGATION TECHNIQUE

Easily grown from fresh seed. Seed should be soaked in water to remove flesh and then sown over a damp peat/coarse sand mix and left in a shaded spot (ideally in a mister) and ignored. Fruit may take up to a year to germinate. Frost tender. Plants resent root disturbance so they should be planted in a sheltered semi-shaded site and then left alone.

CULTIVATION

Occasionally sold by retail plant nurseries and most specialist native plant nurseries, usually under the names *R. cheesemanii* and *R. baueri* var. *cheesemanii*. Most material sold here as *R. baueri* is from Norfolk Island.

ETYMOLOGY

rhopalostylis: Club style

NVS CODE

RHOBAU

CHROMOSOME NUMBER

2n = 32

PREVIOUS CONSERVATION STATUSES

2017 | At Risk – Naturally Uncommon | Qualifiers: RR, SO

2012 | At Risk – Naturally Uncommon | Qualifiers: RR, SO

2009 | At Risk – Naturally Uncommon | Qualifiers: TO

2004 | Range Restricted

[Jump to current conservation status](#)

REFERENCES AND FURTHER READING

Moore LB, Edgar E. 1970. Flora of New Zealand, Volume II. Indigenous Tracheophyta: Monocotyledones except Gramineae. Government Printer, Wellington, NZ. 354 p.

ATTRIBUTION

Description adapted from Moore and Edgar (1970).

Some of this factsheet information is derived from [Flora of New Zealand Online](#) and is used under a [Creative Commons Attribution 3.0 New Zealand](#) licence.

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/rhopalostylis-baueri/>

PDF DATE

06 June 2026