

Rytidosperma corinum

COMMON NAME

bristle grass

SYNONYMS

None

FAMILY

Poaceae

AUTHORITY

Rytidosperma corinum Connor et Edgar

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Grasses

NVS CODE

RYTCOR

CHROMOSOME NUMBER

2n = 48

CURRENT CONSERVATION STATUS

2012 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2009 | Not Threatened

2004 | Not Threatened

DISTRIBUTION

Endemic. North Island, southern Kaweka Range only. South Island on Banks Peninsula, central and south Canterbury and Otago

HABITAT

A species of dry, exposed rocky ground, and cliff faces in montane to subalpine zones. Most commonly found growing in deep fissures and cracks within fractured greywacke cliff faces.

FLOWER COLOURS

Red/Pink

PROPAGATION TECHNIQUE

Easily grown from fresh seed and division of whole plants. Needs a sunny free draining soil to flourish. Best grown in a pot. Dislikes humidity and prolonged wet conditions.

ETYMOLOGY

rytidosperma: Wrinkled seed



WHERE TO BUY

Not commercially available.

TAXONOMIC NOTES

Previously considered a South Island endemic this species has just (2004) been recognised in the North Island from the southern Kaweka Range.

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/rytidosperma-corinum/>