

# Schoenus pauciflorus

## COMMON NAME

bog rush, sedge tussock

## SYNONYMS

*Chaetospora pauciflora* Hook.f.

## FAMILY

Cyperaceae

## AUTHORITY

*Schoenus pauciflorus* (Hook.f.) Hook.f.

## FLORA CATEGORY

Vascular – Native

## ENDEMIC TAXON

Yes

## ENDEMIC GENUS

No

## ENDEMIC FAMILY

No

## STRUCTURAL CLASS

Sedges

## NVS CODE

SCHPAU

## CHROMOSOME NUMBER

2n= 28, 56

## CURRENT CONSERVATION STATUS

2012 | Not Threatened

## PREVIOUS CONSERVATION STATUSES

2009 | Not Threatened

2004 | Not Threatened

## DISTRIBUTION

Endemic. North, South, Stewart, Chatham and Auckland Islands.

Uncommon north of Rotorua.

## HABITAT

Coastal to alpine (up to 1800 m a.s.l.). However, mostly montane to alpine in northern two-thirds of its range. Common in damp seepages along cliff faces, in swamps, in seepages within forest, within mires and around lake tarn and stream sides. Sometimes colonises poorly drained pasture.



Hooker valley, January. Photographer: John Smith-Dodsworth



Mount Torlesse. Photographer: Melissa Hutchison

## FEATURES

Rush-like sedge up to 1 m tall. Rather variable with respect to colour and stature, ranging from stout dark red plants to flaccid bright green specimens. Rhizome short, hard and lignaceous, up to 4 mm diameter. Culms 0.1-1.0 m tall, 0.5-1.8 mm diameter, densely tufted, caespitose, longitudinally striate, grey-green, wine red to bright green. Leaves reduced to numerous, dark red-purple (rarely green) basal sheaths, the uppermost 40-140 mm long, the mucro much elongated with toothed margins. Panicle 15-30 mm long, more or less elongated, bearing 1-9 spikelets at the tips of erect, slightly scabrid branchlets, the whole subtended by a stiff bract overtopping the panicle. Spikelets 5 mm long, 2-4-flowered, lanceolate. Glumes 4-6, lanceolate, 2-3 lowermost smaller, empty, membranous and colourless or occasionally brown, upper glumes darker brown with pale centres, margins hyaline, without cilia. Hypogynous bristles 6, filiform, almost equal in length to style, scabrid, persistent. Stamens 3. Style-branches 3, style often persistent. Nut 2.0-2.5 mm long, slightly less than 1 mm wide, elliptic-oblong, greenish brown to red-brown, lighter brown at the angles, glossy, smooth.

## SIMILAR TAXA

Distinguished from the other rush-like *Schoenus* species by the more or less compact panicle which is subtended by a very long bract which consistently overtops the panicle, and also by the brown rather white or pale cream nuts.

## FLOWERING

September - April

## FRUITING

November - June

## PROPAGATION TECHNIQUE

Easily grown from rooted pieces and the division of whole plants. Can be grown from fresh seed but seed is often slow to germinate. Rather variable and as a horticultural subject it would require some selection of the diversity of forms present in the wild.

## ETYMOLOGY

**schoenus:** Rush

**pauciflorus:** Few flowers

## WHERE TO BUY

Not commercially available.

## NOTES ON TAXONOMY

There is good cytological and molecular evidence that *S. pauciflorus* comprises at least two species.

## ATTRIBUTION

Description adapted from Moore and Edgar (1970).

## REFERENCES AND FURTHER READING

Moore, L.B.; Edgar, E. 1970: Flora of New Zealand. Vol. II. Government Printer, Wellington.

## MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/schoenus-pauciflorus/>