

Tmesipteris tannensis

COMMON NAME

fork fern

SYNONYMS

Lycopodium tannense Spreng.; *Tmesipteris fowerakeri* H.N.Barber,
Tmesipteris forsteri sensu A.Cunn. nom. inv.,

FAMILY

Psilotaceae

AUTHORITY

Tmesipteris tannensis (Spreng.) Bernh.

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Ferns

NVS CODE

TMETAN

CHROMOSOME NUMBER

2n = 208

CURRENT CONSERVATION STATUS

2012 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2009 | Not Threatened

2004 | Not Threatened

DISTRIBUTION

Endemic. New Zealand, North, South, Stewart, Chatham and Auckland Islands.

HABITAT

Coastal to subalpine. Terrestrial or epiphytic on a wide range of hosts and often sympatric with *Tmesipteris elongata* (less frequently with *T. lanceolata* and *T. sigmatifolia*). Less common in coastal and lowland areas in the far north where it is mostly known from higher altitude forest. However, steadily becoming more common from about Whangarei south.



Tararua Forest Park. Photographer: Jeremy Rolfe



Tararua Forest Park. Photographer: Jeremy Rolfe

FEATURES

Rhizome: dichotomously branched, brittle, 2.0-3.5 mm diameter. Aerial shoot: developing over one to many years, but eventually terminating in a small appendage 0.1-0.5× the length of the largest leaves, simple, erect, suberect, or pendulous, 50-1200 mm long, triangular in cross-section, leaves and sporophylls spirally arranged. Leaves coriaceous, brittle, one surface deep glossy green, occasionally with a few stomata towards the far end, other surface dull green covered with stomata; shape variable often on same shoot, oblong, lanceolate, falcate, or ovate, 6-30 mm long × 2.5-9.0 mm broad; apex of leaf very variable often on the same plant, acute, obtuse to truncate, mucronate; mucro 1-2 mm long. Sporophylls: developed in regular or irregular zones or throughout most of the shoot except for the lowermost part, equal to or slightly shorter than the leaves; 5-7 per 10 mm of shoot. Synangium: 4.0-8.0 × 1.5-2.5 mm at point of attachment, biconic, persistent. Spores: yellow, released in a mass, anisopolar, bilateral, monoete, foveolate, concavo-convex, 67-92 × 27-45 microns broad (longitudinal plane).

SIMILAR TAXA

Easily distinguished from all other named New Zealand species of fork fern by the biconic synangia. An apparently unnamed species which may be endemic to Banks Peninsula is rather similar, differing mainly by its octoploid rather than tetraploid chromosome number and less distinctly biconic synangia. This form is in the process of being formally described and named.

FLOWERING

Not applicable - spore producing

FLOWER COLOURS

No flowers

FRUITING

Not applicable - spore producing

PROPAGATION TECHNIQUE

Difficult - should not be removed from the wild.

ETYMOLOGY

tmesipteris: From the Greek tmesis (cutting) and pteris (fern), alluding to the forked appendages on fertile fronds

tannensis: Growing at Tanna, New Hebrides, where the species was first collected

WHERE TO BUY

Not commercially available.

ATTRIBUTION

Fact sheet prepared for NZPCN by P.J. de Lange (June 2009). Description adapted from Chinnock (1975).

REFERENCES AND FURTHER READING

Chinnock, R.J. 1975: The New Zealand Species of *Tmesipteris* (Psilotaceae). *New Zealand Journal of Botany* 1: 743-768.

NZPCN FACT SHEET CITATION

Please cite as: de Lange, P.J. (Year at time of access): *Tmesipteris tannensis* Fact Sheet (content continuously updated). New Zealand Plant Conservation Network. <https://www.nzpcn.org.nz/flora/species/tmesipteris-tannensis/> (Date website was queried)

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/tmesipteris-tannensis/>