

Veronica chionohebe

SYNONYMS

Chionohebe thomsonii var. *glabra* (Cheeseman) Briggs et Ehrend.;
Veronica chionohebe Garn.-Jones; *Veronica thomsonii* var. *glabra*
Cheeseman, *Chionohebe glabra* (Cheeseman) Heads

FAMILY

Plantaginaceae

AUTHORITY

Veronica chionohebe Garn.-Jones

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

NVS CODE

CHIGLA

CHROMOSOME NUMBER

2n = 42

CURRENT CONSERVATION STATUS

2017 | At Risk – Naturally Uncommon | Qualifiers: RR, Sp

PREVIOUS CONSERVATION STATUSES

2012 | At Risk – Naturally Uncommon | Qualifiers: RR, Sp

2009 | At Risk – Naturally Uncommon | Qualifiers: DP

2004 | Range Restricted

FLOWER COLOURS

White

LIFE CYCLE

Seeds are dispersed by ballistic projection, wind and water (Thorsen et al., 2009).

ETYMOLOGY

veronica: Named after Saint Veronica, who gave Jesus her veil to wipe his brow as he carried the cross through Jerusalem, perhaps because the common name of this plant is 'speedwell'. The name Veronica is often believed to derive from the Latin *vera* 'truth' and *iconica* 'image', but it is actually derived from the Macedonian name Berenice which means 'bearer of victory'.

REFERENCES AND FURTHER READING

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/veronica-chionohebe/>



Old Man Range. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.



Chionohebe glabra. Photographer: Phil Garnock-Jones, Licence: CC BY-NC.