

Veronica dilatata

SYNONYMS

Hebe dilatata G.Simpson et J.S.Thomson, *Hebe crawii* Heads

FAMILY

Plantaginaceae

AUTHORITY

Veronica dilatata (G.Simpson et J.S.Thomson) Garn.-Jones

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Trees & Shrubs - Dicotyledons

CHROMOSOME NUMBER

2n = 120

CURRENT CONSERVATION STATUS

2017 | At Risk – Naturally Uncommon | Qualifiers: Sp

PREVIOUS CONSERVATION STATUSES

2012 | At Risk – Naturally Uncommon | Qualifiers: Sp

2009 | At Risk – Naturally Uncommon

2004 | Sparse

BRIEF DESCRIPTION

Low growing sprawling blueish-green shrub forming patches to several metres wide inhabiting mountains of northern Southland and the Takitimu. Twigs erect, reddish. Leaves dished, oval, with pointed tip. Leaf bud with small gap between base of leaves. Flowers in spike to 4cm long, sometimes branched.

DISTRIBUTION

Endemic. South Island, confined to the southern South Island where it is locally known from the Umbrella, Garvie, Eyre and Takitimu ranges. It has also been recorded once from the Blue Mountains near Tapanui.

HABITAT

A species of alpine habitats (usually > 1000 m a.s.l.) which is usually found in rocky ground, in and around boulders, at the toe of scree slopes or in adjoining tussock grassland, seepages and bogs.



Hebe dilatata, Hector Mountains. Photographer: John Barkla, Licence: CC BY.



Hebe dilatata - cultivated. Photographer: John Barkla, Licence: CC BY.

DETAILED DESCRIPTION

Spreading (either openly, densely or even mat-like) low or erect shrub up to 0.4 m tall. Branches decumbent to suberect, old stems dark brown to black; branchlets green, red or brown, bifariously pubescent; internodes 1-18 mm, leaf scars evident. Leaf bud distinct; sinus small and narrowly acute. Petiole 0.5-4 mm, hairy. Leaves 4-25 x 4-11 mm, glossy or dull green to glaucescent on both surfaces, obovate, spatulate or elliptic, coriaceous, concave, erect to recurved, arranged decussately or somewhat distichously; apex obtuse to rounded. Inflorescences with 5-52 flowers, lateral unbranched or tripartite (when unbranched with lowermost bracts enlarged and sterile); 5-42 mm, usually longer than or equal to subtending leaves. Peduncle 10-90 mm; rachis 5-23 mm. Flowers gynodioecious. Pedicels 0.5-2 mm, calyx 2.4-3.8 mm, 3-5-lobed, with anterior lobes united from one third to all the way to apex; lobes ovate, subacute, obtuse, or emarginate. Corolla tube 1.2-2.5 x 1.8-2.5 mm, funnelform, glabrous. Stamen filaments 3.8-4.2 mm, anthers magenta 1.9-2.2 mm. Ovary 0.7-1.5 mm; style 3.4-4.2 mm. Capsules 3.4-4.5 x 2.1-2.8 mm, obtuse to subacute, glabrescent. Seeds 0.9-1.1 x 0.6-1 mm, straw-yellow to pale brown, broad ovoid to sub-discoid.

SIMILAR TAXA

A very distinctive species whose closest relative seems to be *Veronica societatis*, which is a northern South Island species known at present from just one site on Mt Murchison above the Buller River. It is a sparingly branched, whip-like shrub with a rather different chromosome number ($2n = 42$ cf. $2n = 120$). In the field it could be confused with *V. cockayneana*. *Veronica cockayneana* differs from *V. dilatata* by its usually more erect bushy habit, and leaves which are glossy above and glaucous below. The inflorescences of *V. dilatata* differ in that they are all usually branched (with at least 1-2 branches near the base), or if unbranched, the basal bracts are rather more conspicuous than usual and sterile. Within *V. dilatata* a distinctive, more upright bushy form from the Takitimu Range has been separated as *Hebe crawii* Heads, mainly due to its erect habit and usually tripartite inflorescences. It is now included within the normal range of variation exhibited by *V. dilatata* but many botanists still prefer to regard it as a distinct species.

FLOWERING

November - March

FLOWER COLOURS

Violet/Purple, White

FRUITING

November - April

LIFE CYCLE

Seeds are wind dispersed (Thorsen et al., 2009).

PROPAGATION TECHNIQUE

Easily grown from semi-hardwood cuttings and layered, rooted pieces. Fresh seed (as in most hebes) probably germinates well. An excellent plant for a sunny, open spot in a rock garden or in a tub/planter on a patio. As a high alpine it rarely flowers in the northern part of New Zealand. Nevertheless it has an attractive form and should be more widely grown.

THREATS

A naturally uncommon, wide ranging species which is never very common at any particular place.

ETYMOLOGY

veronica: Named after Saint Veronica, who gave Jesus her veil to wipe his brow as he carried the cross through Jerusalem, perhaps because the common name of this plant is 'speedwell'. The name Veronica is often believed to derive from the Latin *vera* 'truth' and *iconica* 'image', but it is actually derived from the Macedonian name Berenice which means 'bearer of victory'.

dilatata: Expanded

WHERE TO BUY

Not commercially available.

ATTRIBUTION

Fact Sheet Prepared by P.J. de Lange (1 November 2009). Description based on Bayly & Kellow (2006)

REFERENCES AND FURTHER READING

Bayly M.; Kellow A. 2006: An Illustrated Guide to New Zealand Hebes. Te Papa Press: Wellington
Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 11: 285-309

NZPCN FACT SHEET CITATION

Please cite as: de Lange, P.J. (Year at time of access): *Veronica dilatata* Fact Sheet (content continuously updated). New Zealand Plant Conservation Network. <https://www.nzpcn.org.nz/flora/species/veronica-dilatata/> (Date website was queried)

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/veronica-dilatata/>