


INDIGENOUS HIGHER PLANTS (LYCOPODS, FERNS, GYMNOSPERMS,
FLOWERING PLANTS) OF KAWEKA RANGE AND ADJACENT AREAS
1000-5657 FT/300-1725 M


Highest point: Kaweka, 5657 ft/1725m, grid ref. N123 829650

A.P.Druce, Wellington, August 1992

(Based on visits 1957-1989)

 boundary of area studied

FOR LIST OF PLANTS FROM
KANEKA AND BLACK BIRCH
RANGES ABOVE 3000 FT
(900M) SEE LIST 72



GYMNOSPERM TREES AND SHRUBS

- Dacrycarpus dacrydioides*
Dacrydium cupressinum
Halocarpus bidwillii
Lepidothamnus laxifolius
L. intermedius × *L. laxifolius*
Phyllocladus alpinus s.s.
Podocarpus hallii
P. nivalis
P. totara s.s.
P. hallii × *P. nivalis*
Prumnopitys ferruginea
P. taxifolia
- Nestegis cunninghamii*
N. lanceolata
N. montana (Cunningham 1974)
Nothofagus fusca
N. menziesii
N. solandri var. *solandri*
N. s. var. *differticoides*
N. fusca × *N. solandri*
Pennantia corymbosa
Pittosporum eugenioides
P. ralphii
P. tenuifolium ssp. *colensoi*
P. t. ssp. *tenuifolium*
P. t. ssp. *colensoi* ⇒ ssp *tenuifolium*
P. ralphii × *P. tenuifolium*
Plagianthus regius var. *regius*
Pseudopanax arboreus var. *arboreus*
P. colensoi s.s.
P. crassifolius
P. edgarleyi
P. simplex (incl. *P. s.* var. *sinclairii*)
Schefflera digitata
Sophora microphyllus
S. tetrapetala
Streblus heterophylla

DICOT TREES

- Alectryon excelsus* var. *excelsus*
Aristotelia serrata
Carpodetus serratus
Elaeocarpus hookerianus
Fuchsia excorticata
Griselinia littoralis
Hedycarya arborea
Hohertia populnea var. *lanceolata*
Lophomyrtus obcordata
Mastopipera excelsum var. *excelsum*
Melicrytus lanceolatus (incl. *M. l.* var. *laticornis*)
M. tamiflorus ssp. *tamiflorus*
Mida salicifolia (Cunningham 1974)
Myoporum laetum var. *laetum*
Myrsinace australis
M. salicina

DICOT SHRUBS

- Alepis flavida*
Alseuosmia pusilla
Aristotelia fruticosa var. *microphylla*
A. f. × *A. serrata*
Brachyglottis bidwillii var. *bidwillii*
B. repanda s.s.
B. rotundifolia var. (*B. eleagnifolia*)
B. bidwillii × *B. rotundifolia*

<i>Carmichaelia arborea</i> var. (<i>C.</i> <i>odorata</i>)	<i>Conaria arborea</i>
<i>C. enysii</i> var. <i>orbiculata</i> (Cunningham 1974, as <i>C. orbiculata</i>)	<i>C. kingiana</i>
<i>C. sp.</i> (subgenus <i>Carmichaelia</i>)	<i>C. plumosa</i>
<i>Cassinia vauvilliersii</i>	<i>C. pteridoides</i>
<i>Coprosma acerosa</i> ssp.	<i>C. arborea</i> × <i>C. kingiana</i>
<i>C. cheesemanii</i> s.s.	<i>C. a.</i> × <i>C. pteridoides</i>
<i>C. colensoi</i> s.s.	<i>C. kingiana</i> × <i>C. pteridoides</i>
<i>C. crassifolia</i>	<i>Corokia cotoneaster</i> var. <i>cotoneaster</i>
<i>C. depressa</i>	<i>Cyathodes dealbata</i> (incl. <i>C. pumila</i>)
<i>C. foetidissima</i>	<i>C. empetrifolia</i>
<i>C. grandifolia</i>	<i>C. sp.</i> (<i>C. juniperina</i> agg.) (lvs vs. ≤ 1cm × ≤ 1mm)
<i>C. linariifolia</i>	<i>Dracophyllum recurvum</i>
<i>C. lucida</i> s.s.	<i>D. subulatum</i>
<i>C. microcarpa</i>	<i>D. recurvum</i> × <i>D. subulatum</i>
<i>C. perpusilla</i> ssp. <i>perpusilla</i>	<i>Epactis alpina</i>
<i>C. propinqua</i> ssp. <i>propinqua</i>	<i>Fuchsia excorticata</i> × <i>F. perscandens</i>
<i>C. pseudocuneata</i> ssp. <i>pseudo-</i> <i>cuneata</i>	<i>Gaultheria antipoda</i>
<i>C. thamnoides</i> (incl. <i>C. polymorpha</i>)	<i>G. colensoi</i>
<i>C. rigida</i>	<i>G. macrostigma</i>
<i>C. robusta</i>	<i>G. sp.</i> (unnamed; included in <i>G.</i> <i>depressa</i> , as var. <i>novae-zelandiae</i> , by Franklin 1962)
<i>C. rotundifolia</i>	<i>G. antipoda</i> × <i>G. colensoi</i>
<i>C. rubra</i>	<i>G. a.</i> × <i>G. macrostigma</i>
<i>C. rugosa</i> ssp. <i>rugosa</i>	<i>G. a.</i> × <i>G. sp.</i> (unnamed)
<i>C. tenuifolia</i>	<i>G. colensoi</i> × <i>G. sp.</i> (unnamed)
<i>C. sp.</i> (i) (unnamed; included in <i>C.</i> <i>parviflora</i> by Oliver, 1935, and others)	<i>Geniostoma rupestre</i> var. <i>ligostig-</i> <i>folium</i> (Cunningham 1974)
(undersurface not minutely hairy as is usual in <i>C. parviflora</i> ; droplets vs. unpigmented, occ. purplish red)	<i>Habe colensoi</i> (incl. <i>H. c.</i> var. <i>hillii</i>)
	<i>H. macrocarpa</i> var. (<i>H. corriganii</i>)
	<i>H. odora</i> s.s.
<i>C. sp.</i> (ii) (unnamed; aff. <i>C. cheesemanii</i>)	<i>H. stricta</i> var. <i>stricta</i>
(lvs broader than in <i>C. cheesemanii</i>)	<i>H. tetragona</i> var. <i>tetragona</i>
<i>C. propinqua</i> × <i>C. robusta</i>	<i>H. venusta</i> (incl. <i>H. brachysiphon</i>)
<i>C. robusta</i> × <i>C. rugosa</i>	<i>H. sp.</i> (i) (unnamed; included in <i>H.</i> <i>stricta</i> , as var. <i>egmontiana</i> and as var. <i>lata</i> , by Moore 1961)
<i>C. r.</i> × <i>C. tenuifolia</i>	

- H. sp.(ii) (Veronica squalida) var.*
H. macrocarpa X *H. venusta* (?)
Helichrysum aggregatum var.
aggregatum
H. depressa
Ileostylus micranthus
Korthalsella lindsayi
Kunzea ericoides agg.
Leptospermum scoparium
Leucopogon fasciculatus var.
 (juv. lvs not much broader than
 adult lvs)
L. fraseri s.s.
Melicope simplex
Melicotus alpinus s.s.
M. micranthus (incl. *M. m.* var. *micto-*
phylla)
Myrsine divaricata s.s.
M. nummularia
Neomyrtus pedunculata
Olearia arborescens
O. furfuracea var.
O. hectorii var. (lvs smaller than
 in var. *hectorii*)
O. ilicifolia
O. nummulariifolia var. *nummul-*
O. arborescens X *O. furfuracea*
O. a. X *Celmisia spectabilis*
Pentachondra pumila
Petaxilla tetrapetala
Pimelea buxifolia
P. longifolia var. *longifolia*
P. microphylla
P. prostrata
P. suteri var. (*P. oreophila*)
- Pittosporum divaricatum* (Cunningham
 1974)
P. rigidum var. *rigidum*
Pomaderris ericifolia
Pseudopanax anomalous
Pseudowintera axillaris
P. colorata
Solanum aviculare var. *aviculare*
Tecridium patrifolium (Cunningham
 1974)
Tupeia antarctica
Urtica ferox
- [MONOCOT LIANES]**
- Ripogonum scandens* (Cunningham
 1974)
- [DICOT LIANES AND RELATED PLANTS]**
- Calystegia tuguriorum*
Clematis afoliata (Longfellow Ra.)
C. foetida
C. forsterti
C. paniculata
C. quadriflora
Metrosideros colensoi (incl. *M. c.*
 var. *pendens*)
M. diffusa
Muehlenbeckia australis
M. axillaris
M. complexa (Cunningham 1974)
M. australis X *M. axillaris*
Parsonsia capsularis var.
P. heterophylla
P. capsularis X *P. heterophylla*
Passiflora tetrandra (Cunningham
 1974)
Rubus australis
R. cissoides var. (caulats narrower
 than in var. *cissoides*)

R. schmideloides var. *schmideloides*

R. squarrosum

PSILOPSIDS AND LYCOPODS

Tmesipteris tannensis (Cunningham 1974)

Lycopodium australianum

L. fastigiatum

L. scariosum

L. varium (incl. *L. billardieri* and
L. novae-zelandicum)

L. volubile

B. discolor

B. fluviale agg.

B. membranaceum

B. minus

B. nigrum (Cunningham 1974)

B. penna-martini

B. procerrum

B. vulcanicum

B. sp.(i) (unnamed) (common sp.; lower
pinnae reduced in length)

B. sp.(ii) (unnamed) (mountain sp.;
lower pinnae not reduced)

B. sp.(iii) (unnamed) (cliff sp.; lower pinna
not reduced) ("The Green Bay Species
of Crookes 1963")

B. procerrum X *B. sp.(i)*

Botrychium austale

B. bifforme

Cheilanthes humilis

Ctenopteris heterophylla

Cyathea colensoi

C. dealbata

C. medullaris

C. smithii

Dicksonia fibrosa

D. lanata s.s.

D. squarrosa

Gleichenia dicarpa

Grammitis billardieri

G. magellanica ssp. *nothofageti*

G. patagonica

G. poeppigiana

Histropteris incisa

Hymenophyllum bivalve

H. cupressiforme

H. demissum

H. dilatatum

FERNS

Adiantum cunninghamii

A. diaphanum (Cunningham 1974)

Asplenium bulbiferum

A. colensoi

A. flabellifolium

A. flaccidum

A. gracillimum

A. hookerianum

A. lyallii

A. oblongifolium

A. polyodon

A. richardii

A. terrestris ssp. *terrestris*

A. sp. (*A. trichomanes* agg.)

A. bulbiferum X *A. flaccidum*

A. b. X *A. hookerianum*

A. flaccidum X *A. gracillimum*

A. f. X *A. hookerianum*

A. gracillimum X *A. hookerianum*

Azolla filiculoides

Blechnum chambersii

B. colensoi

H. flabellatum
H. minimum
H. multifidum
H. peltatum
H. pulcherrimum
H. parvum
H. revolutum
H. sanguinolentum agg.
H. scabrum
Hypolepis ambigua
H. millefolium
H. rufobarbata
Leptolepia novae-zelandiae
Leptopteris hymenophylloides
L. superba
L. hymenophylloides x *L. superba*
Ophioglossum cordaceum agg.
Paesia scaberula
Pellaea rotundifolia
Phymatosorus pustulatus
Pleurozorus tatifolius
Pneumatopteris pennigera
Polystichum richardii
P. vestitum
Pteridium esculentum
Pteris tremula
Pyrosia eleagnifolia
Rumohra adiantiformis
Sticherus cunninghamii
Trichomanes venosum

ORCHIDS

Acianthus sinclairii
Adenochilus gracilis
Aporostylis bifolia
Caladenia carneoides agg.
C. lyallii

Chiloglottis cornuta
Corybas cryptanthus
C. mactanthis
C. oblongus
C. rivularis s.s.
C. trilobus s.s.
Earia mucronata s.s.
Gastrodia cunninghamii
G. minor
G. sesamoides
Genoplesium nudum
Microlis unifolia
Orthoceras novae-zelandiae
Prasophyllum edensoi
Pterostylis allobula
P. foliata
P. furcata
P. humilis
P. patens
P. venosa
*P. sp. (unnamed; aff. *P. montana*)*
Spiranthes novae-zelandiae
Thelymitra cyanea
T. decora
T. formosa
T. hatchii
T. longifolia
T. pauciflora

GRASSES

Agrostis dyeri s.s.
A. muelleriana
A. muscosa
A. personata
Amphibromus fluitans
Chionochloa conspicua (Cunningham 1974)
C. pallens ssp. *pallens*

- C. rubra* ssp. *rubra*
C. pallens X *C. rubra*
Cortaderia fulvida
Deschampsia caespitosa var.
macrantha
D. novae-zelandiae
D. tenella
Deyeuxia aucklandica
D. avenoides (incl. *D. a.* var. *brachy-antha*)
D. sp. (*D. quadriflora* agg.) (lvs narrow)
Dichelachne crinita
Echinochogon ovatus
Elymus multiflorus
E. hardyioides
E. sp. (aff. *E. tectorius*) (plants vs. glaucous; lamina vs. flat, up to 5mm wide; awns 4-6cm long) var. (culms erect)
E. multiflorus X *E. sp.* (?)
Festuca novae-zelandiae
F. sp. (unnamed; aff. *F. rubra*)
Hierochloe tectorum
H. redolens
Koeleria cheesemanii
Lachnagrostis filiformis s.s.
L. striata
L. sp.(i) (unnamed; included in *L. filiformis*, as var. *semiglabra*, by Zofov 1965)
L. sp.(ii) (unnamed) (tall, slender plants, with narrow lvs; glumes up to 4mm long; awn short or absent)
L. filiformis X *L. striata* (?)
Microlaena cavenacea
M. stipoides
- Poa anceps* ssp. *anceps*
*P. breviglumi*s
P. cita (both narrow-lvd and broad lvs forms present)
P. colensoi
P. imbecilla
P. lindsayi
P. pusilla
P. sp. (unnamed; aff. *P. novae-zelandiae*) (scree plants with *Rytidosperma buchananii* narrow lvs)
R. clavatum
R. gracile
R. pulchrum
R. setifolium
R. viride
Trisetum sp. (i) (unnamed; aff. *T. antarcticum*) (lf sheath densely hairy; lf blade flat, up to 5mm wide, margins hairy)
T. sp.(ii) (unnamed) (plants slender; lf sheath usually glabrous; lf blade flat, up to 2.5mm wide, margins smooth)
T. sp.(iii) (unnamed; aff. *T. antarcticum*) (plants robust; lf sheath glabrous in upper part, densely hairy in lower part; lf blade flat, up to 7mm wide, margins scabrid)
- [SEDGES]**
- Baumea rubiginosa*
B. tenax
Carex acicularis
C. breviculmis
C. buchananii
C. cirthosa

- C. comans*
C. coriacea
C. dipsacea var. (*C. tahoata*)
C. dissita ss.
C. echinata var. (*C. stellulata* var.
 australis)
C. fasciculata
C. flagellifera
C. forsteri
C. gaudichaudiana
C. inversa
C. maonica
C. petriei
C. pyrenaica var. *cephalotes*
 (Cunningham 1974)
C. secta ss.
C. sinclairii
C. solandri
C. testacea ss.
C. virgata
Carphe alpina
Eleocharis acuta
E. gracilis
E. sphacelata
Gahnia pauciflora
Isolepis aucklandica
I. catigenis
I. crassiuncula
I. distigmatosa
I. fluitans
I. habra
I. pottsii
I. reticulatis
I. subtilissima
Lepidosperma australe
Machaerina sinclairii
Motolotia affinis
- Oreobolus pectinatus*
Schoenus maschalinus
S. pauciflorus ss.
*S. sp. (unnamed; aff. *S. pauciflorus*)*
 ("robust form" of Edgar 1970, p. 74)
Uncinia banksii
U. caespitosa
U. clavata
U. distans
U. ferruginea
U. fuscovaginata var. *fuscovaginata*
U. gracilenta
U. laxiflora
U. rubra
U. rupestris (incl. *U. angustifolia*)
U. scabra
U. silvestris (Cunningham 1974)
U. uncinata
U. vitidis
U. zotovii
*U. sp.(i) (unnamed) (*U. affinis* sensu
 Hamlin 1959, non *U. riparia* var.
affinis Col. ex Clarke; see Nootboom
 1978)*
*U. sp.(ii) (unnamed; aff. *U. nervosa*)*
 (more slender than *U. nervosa*; see
 Edgar 1970, p. 23)

RUSHES AND ALLIED PLANTS

- Centrolepis ciliata*
Empodisma minus
Juncus australis
J. gragiflorus
J. novae-zelandiae
J. planifolius
Luzula colensoi
L. decipiens

L.migrata
L.picta s.s.
L.pufa s.s.
L.subclavata
L.sp.(unnamed; included in L.picta, as var. limosa, by Edgar 1966)
L.cotensoi × *L.migrata*

Brachyglottis lagopus var. *lagopus*
Brachyscome radicata
Celmisia glandulosa ssp. *glandulosa*
 (incl. *C.g. var. latifolia*)
C.gracilenta ssp.
C.incana (incl. *C.discolor*)
C.spectabilis ssp. *spectabilis*
C.sp.(unnamed; aff. C.gracilenta)
 (plants strongly rhizomatous)
C.gracilenta × *C.incana*
C.g. × C.spectabilis
C.incana × *C.spectabilis*
Craspedia minor var. *minor* (incl. *C.major*)
C.viscosa
C.sp.(i)(unnamed) (plants summer-green;
 lvs small, green; both surfaces densely
 clad in multicellular, non-glandular hairs,
 with or without cottony tips, over short
 glandular hairs; margins white with
 cottony hairs; capitula small, 0.8-1.5cm;
 fls white)
C.sp.(ii)(unnamed) (plants summer-green;
 lvs green, or greenish white; both sur-
 faces densely clad in multicellular,
 non-glandular hairs, with cottony tips;
 margins white with cottony hairs; cap-
 itula large, 1.8-2.5cm diam.; fls white)
Gnephium audax
G.delicatum
G.gymnocephalum
G.involucratum
G.limosum s.s.
G.paludosum s.s.
G.tuhinicum
G.sphaericum
Helichrysum filicaule
Lagenifera montana
L.pinnatifida
L.pumila
L.strangulata

COMPOSITE HERBS

Anaphalis kerriensis s.s.
A.sp.(i)(Helichrysum bellidioides s.s.)
A.sp.(ii)(unnamed) ("Helichrysum alpinum" of Cockayne (1928)) (lvs and
 capitula larger than in *A.sp.i*)

- Leptinella squalida* agg.
Leucogenes leontopodium
Microseris scapigera var.
Pseudognaphalium sp. (unnamed; aff. *P. luteoalbum*) (lvs mostly linear-spathulate to linear, acute)
Raoulia albo-sericea
R. glabra
R. grandiflora var. *grandiflora*
R. tenuicaulis (incl. *R. t.* var. *dimorpha* and *R. t.* var. *pusilla*)
R. albo-sericea × *R. glabra*
R. glabra × *Helichrysum filicavum*
Senecio glaucophyllus ssp. *toa*
S. hispidulus
S. minimus
S. quadridentatus
S. rufiglandulosus ssp. *rufiglandulosus*
S. wairauensis
S. sp. (unnamed; included in *S. glaucophyllus*, as ssp. *discoidaeus*, by Ornduff 1960) ssp.
Taraxacum magellanicum
Vittadinia australis agg.
- Cardamine* sp.(i) (*C. debilis* agg.) ("Long Style" of Pritchard 1957)
C. sp.(ii) (*C. debilis* agg.) ("Glossy Leaf" of Pritchard 1957)
C. sp.(iii) (*C. debilis* agg.) ("Narrow Petal" of Pritchard 1957)
C. sp.(iv) ("Scree Race" of Pritchard 1957)
C. sp.(v) ("Tussock Race" of Pritchard 1957)
Centella uniflora
Colobanthus acicularis
C. apetalus (incl. *C. a.* var. *alpinus*)
C. strictos
Crassula sp. (cf. *C. sieberiana* and *C. tetrameta*)
Dactylanthus taylorii
Dichondra repens agg.
D. sp. (unnamed; aff. *D. brevifolia*)
Drosera arcturi
D. binata
D. peltata ssp. *auriculata*
D. spathulata
Epilobium alsinoides s.s.
E. astonii
E. atriplicifolium
E. billardierianum s.s.
E. blennescens
E. chionanthum
E. chlorifolium
E. cinereum
E. cockayneanum
E. glabellum s.s.
E. insulare
E. komarovianum
E. linnaeoides
E. macropus
E. microphyllum
E. microphyllum
E. netteroides

DICOT HERBS OTHER THAN COMPOSITES

- Acaena anserinifolia*
A. emarginata
A. inermis
A. microphylla var. *microphylla*
A. profunda incisa
Australinia pusilla
Aciphylla colensoi var. *colensoi*
A. sp. (unnamed; aff. *A. squarroso*)
Anisotome aromatica

- E. microphyllum*
E. microphyllum
E. netteroides

- 11
- E. nummulariifolium*
E. pallidiflorum

E. pernitens
E. pubens
E. pycnostachyum
E. rotundifolium
E. tenuipes ss.
E. sp. (unnamed; included in *E. brunneascens*, as ssp. *minutiflorum*, by Raven and Engelhorn, 1971)
E. melanocaulon × *E. pycnostachyum*
Euphrasia cuneata
E. zelandica agg.
Forstera bidwillii (incl. *F. b.* var. *densifolia*)
Galium perpusillum s.s.
G. propinquum
G. trilobum
G. sp. (unnamed; aff. *G. perpusillum*)
 (lvs narrowly elliptic, 1-3 mm long, rounded to subacute; stipules slightly narrower and shorter, acute; apical bristle 0 or 1, rarely 2 or 3; fls c. 2 mm diam., white)
- Gentiana bellidifolia* var. *bellidi-folia*
G. grisebachii
G. montana (incl. *G. m.* var. *stolonifera*, *G. patula*, and *G. bellidi-folia* var. *pulchella*)
Geranium potentilloides var. *potentillordes*
G. sessiliflorum var. *novae-zelandiae*
G. sp. (unnamed; aff. *G. microphyllum*)
- Gaura* sp. (*G. parviflorum* agg.)
 (stock short; terminal leaflet about as long as broad; fls up to 13 mm diam.)
Glossostigma elatinoides
Gonocarpus aggregatus
G. micranthus ssp. *micranthus*
Gratiola sexdentata
Gunnera dentata var. *dentata*
G. prorepens
Haloragis erecta ssp. *erecta*
Hydrocotyle dissecta
H. elongata
H. heteromeria
H. hydrophila
H. microphylla
H. moschata
H. novae-zelandiae ss.
H. sulcata
H. sp. (unnamed; included in *H. nz.*, as var. *montana*, by Kirk 1899) var.
Hyperticum japonicum agg.
H. sp. (unnamed) (plants small; lvs lanceolate, 2-3 × 0.7-1 mm)
- Hypsela rivalis*
Jovellana repens
J. sinclairii
Kelleria dieffenbachii
K. laxa
Lilaeopsis ruthiana
Linum monogynum var. *monogynum*
Mazus radicans
Mentha cunninghamii
Montia fontana s.s.

- Myosotis forsteri*
M. sp.(i) (M. australis agg.) (fls yellow)
*M. sp.(ii) (unnamed; included in M. Pratia angulata agg. pygmaea, as var. *drucei*, by Moore 1961)*
Myrsiphyllum pedunculatum ssp. novae-zelandicum
M. propinquum
M. triphyllum
*Neopaxia sp. (*Montia calycina*)*
Nettlera ciliata
*N. depressa (incl. *N. cunninghamii*)*
N. setulosa
*N. sp. (unnamed; aff. *N. dichondrifolia* (hairs present on both surfaces of lf, straight, not hooked as in *N. dichondrifolia*))*
*Oreomyrtus colensoi ss. (incl. *Oc. Rotippa palustris* var. *hispida* and *O.c. multifida*)*
*O. sp.(i) (unnamed; aff. *O. ramosa*)*
*O. sp.(ii) (unnamed; included in *O. colensoi*, as var. *delicatula*, by Allan 1961)*
*Oxalis caespitosa ssp. *caespitosa**
O. colensoi
*O. lactea ssp. *drucei**
*O. macrophylla ssp. *robusta**
Oxalis exilis
O. magellanica
*Patahebe diffusa ssp. *diffusa**
P. olsenii
P. spathulata
*Phyllachne colensoi var. *colensoi**
Plantago novae-zelandiae
P. taoulii
P. triandra ss.
P. uniflora
Potentilla anserinoides
Pratia angulata agg.
P. petrosilla
Psychrophila novae-zelandiae
Ranunculus amphitrichus
*R. cheesemanii var. (*R. carsei*)*
R. glabrifolius
R. insignis
R. limosella
R. macropus
R. nivicola
R. reflexus
R. royi
*R. sp. (unnamed; included in *R. lappaceus*, as var. *macrophyllus*, by Kirk 1899)*
Rumex flexuosus
*Scandia sp. (*S. rosifolia* agg.)(leaflets ovate-lanceolate)*
Schizilema colensoi
S. nitens
S. trifoliolatum
S. sp. (unnamed) (lvs cut 1/3 - 2/3 to base; scattered hairs on upper surface; involucral bracts obtuse; fr = or < its pedicel)
Scleranthus biflorus
S. uniflorus
Stackhousia minima
*Stellaria decipiens (incl. *S. parviflora*)*
S. gracilenta
Urtica incisa

Viola cunninghamii

V. filicaulis

V. lyallii

Wahlenbergia albomarginata var.

pygmaea

W. colensoi

W. rupestris

No. of species: 615.

Doubtful and erroneous records not included in main list

<u>Record</u>	<u>Probable identity of plant on which record based</u>
(Conn. 1974 = Cunningham 1974)	
<i>Aciphylla squarrosa</i> (Conn. 1974)	<i>Aciphylla</i> sp. (unnamed)
<i>Astelia nervosa</i> (Conn. 1974)	<i>Astelia fragrans</i>
<i>Carex diandra</i> (Conn. 1974)	<i>Carex dipsacea</i>
<i>C. ternaria</i> (Conn. 1974)	<i>C. coriacea</i>
" <i>Calmisia graminifolia</i> × <i>C. spectabilis</i> " (Given 1984)	<i>Calmisia gracilenta</i> × <i>C. spectabilis</i>
<i>Chionochloa flavescens</i> (Elder 1959, as <i>Danthonia flavescens</i>)	<i>Chionochloa pallens</i> ssp. <i>pallens</i>
<i>C. rigida</i> (Elder 1959, as <i>Danthonia rigida</i>)	<i>C. rubra</i> ssp. <i>rubra</i>
<i>Clematis cunninghamii</i> (Conn. 1974, <i>Clematis foetida</i> as <i>C. parviflora</i>)	
<i>Coprosma parviflora</i> (Conn. 1974)	<i>Coprosma</i> sp. (i) (unnamed)
<i>Cotatia angustissima</i> (Conn. 1974)	<i>Cotatia plumosa</i>
<i>C. satmentosa</i> (Conn. 1974)	<i>C. arborea</i> × <i>C. pteridoides</i>
<i>Dracophyllum longifolium</i> var. (Conn. 1974, as <i>D. filifolium</i>)	
<i>Gaultheria depressa</i> (Conn. 1974)	<i>Gaultheria</i> sp. (unnamed)
<i>G. rupestris</i> ? (Conn. 1974)	
<i>Geranium microphyllum</i> (Conn. 1974)	<i>Geranium</i> sp. (unnamed)
<i>Hebe diosmifolia</i> (Cheeseman 1925, as <i>Veronica diosmifolia</i>)	<i>Hebe venusta</i>

Kirkianella novae-zelandiae

(Conn. 1974)

Koeleria novozelandica (Conn.) *Deyeuxia arenoides*
(1974, as *K. kurtzii*)*Lagenifera petiolata* (Conn. 1974) *Lagenifera strangulata**Melicytus crassifolius* (Conn. 1974), *Melicytus alpinus*
as *Hymenanthera crassifolia*)*Microtis parviflora* (Conn. 1974)*Myosotis pygmaea* (Conn. 1974)*Nectaria dichondrifolia* (Conn. 1974)*Olearia pachyphylla* (Carse 1929)*Oreomyrrhis rigida* (Conn. 1974)*Parahelbe cataractae* (Conn. 1974)*P. hookeriana**Pimalea aridula* (Conn. 1974)*P. gnidia* (Conn. 1974)*P. lyallii* (Elder 1959)*P. pseudolyallii* (Conn. 1974, p. 39)*Plantago triantha* (Conn. 1974)

"Poa laevis" (Elder 1959, p. 23)

Pterostylis banksii (Conn. 1974)*P. trullifolia* (Conn. 1974)*Raoulia australis* (Conn. 1974)*Rytidosperma nigricans* (Conn. 1974, as *Notodanthonia nigricans*)*Schizolema hydrocotyloides* (Conn. 1974) *Schizolema* sp. (unnamed)*Senecio scaberula* (Conn. 1974), *Senecio hispidulus*
as *Erechtites scaberula*)*Wahlenbergia gracilis* (Conn. 1974) *Wahlenbergia colensoi* / *W. ropestris*REFERENCES

Allan, H.H. 1961: Flora of New Zealand. Govt. Printer, Wellington. Vol. 1.

Carse, M. 1928: Botanical notes, new species & varieties. *Trans. N.Z. Inst.* 59: 315-6.

Cheseman, T.F. 1925: Manual of the N.Z. flora Ed. 2. Govt. Printer, Wellington.

Cockayne, L. 1928: The vegetation of New Zealand. 2nd ed. Engelmann, Leipzig.

Crookes, M. 1963: New Zealand ferns. Ed. 6. Whitcombe and Tombs.

- Cunningham, A. 1974: Headwaters of the Tuteaikuri catchment: study of an erosion problem in Hawke's Bay. N.Z. Forest Service Tech. Paper. No. 62. N.Z. Forest Service, Wellington. 59 pp.
- Edgar, E. 1966: Huzula in New Zealand. N.Z. Jl Bot. 4: 159-84
- 1970: In Flora of New Zealand. Govt. Printer, Wellington. Vol. 2.
- Elder, N.L. 1959: Vegetation of the Kaweka Ranges. Trans Roy Soc. N.Z. (Bot.) 87: 9-26.
- Franklin, D.A. 1962: The Ericaceae in New Zealand (Gaultheria and Pentstemon). Trans R. Soc. N.Z. Bot. 1: 155-73.
- Given, D.R. 1984: A taxonomic revision of Olearia subgenus Pelliculatae section Petiolatae (Compositae-Asterae). N.Z. Jl. Bot. 22: 139-58.
- Hamlin, B.G. 1959: A revision of the genus Uncinia (Cyperaceae-Caricoidae) in New Zealand. Dom. Mus. Bull. 19.
- Kirk, T. 1899. "The student's flora of New Zealand." Govt. Printer, Wellington.
- Moor, L.B. 1961: In Flora of New Zealand. Govt. Printer, Wellington. Vol. 1.
- Nooteboom, H.P. 1978: A taxonomic revision of the Malesian and Australian species of Uncinia (Cyperaceae). Blumea 24: 511-20.
- Oliver, W.R.B. 1935: The genus Coprosma. Bishop Mus. Bull. 132.
- Ornduff, R. 1960: An interpretation of the Senecio lautus complex in New Zealand. Trans. R. Soc. N.Z. 80: 63-77.
- Pritchard, G.G. 1957: Experimental taxonomic studies on species of Cardamine Linn. in New Zealand. Trans. R. Soc. N.Z. 85: 75-89.
- Raven, P.H.; Engelhorn, T. 1971: New taxa and new combinations in Australasian Epilobium (Onagraceae). N.Z. Jl Bot. 9: 345-50.
- Zotov, V.D. 1965: Grasses of the subantarctic islands of the New Zealand region. Rec. Dom. Mus. Wellington 5: 101-46.
-

