



Shakespear 2015/16 Seed Collection



Table of Contents

Introduction	1
<i>Arthropodium cirratum</i>	2
<i>Beilschmiedia tarairi</i>	3
<i>Carex virgata</i>	4
<i>Carpodetus serratus</i>	5
<i>Coprosma macrocarpa</i> subsp. <i>macrocarpa</i>	6
<i>Coprosma robusta</i>	7
<i>Cordyline australis</i>	8
<i>Dysoxylum spectabile</i>	9
<i>Entelea arborescens</i>	10
<i>Gahnia lacera</i>	11
<i>Kunzea robusta</i>	12
<i>Leptospermum scoparium</i> var. <i>scoparium</i>	13
<i>Melicytus ramiflorus</i>	14
<i>Metrosideros excelsa</i>	15
<i>Myrsine australis</i>	16
<i>Phormium tenax</i>	17
<i>Pittosporum crassifolium</i>	18
<i>Sophora microphylla</i>	19
<i>Vitex lucens</i>	20
Glossary	21

Made on the New Zealand Plant Conservation Network website – www.nzpcn.org.nz

Copyright

All images used in this book remain copyright of the named photographer. Any reproduction, retransmission, republication, or other use of all or part of this book is expressly prohibited, unless prior written permission has been granted by the New Zealand Plant Conservation Network (info@nzpcn.org.nz). All other rights reserved.

Arthropodium cirratum

Common Name(s):

Renga lily, Rengarenga, Rock lily

Current Threat Status (2012):

Not Threatened

Threats:

Not Threatened

References and further reading:

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 2009 Vol. 11 No. 4 pp. 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1515



Caption: In cultivation

Photographer: Jesse Bythell



Caption: Coromandel, November

Photographer: John Smith-Dodsworth

Beilschmiedia tarairi

Common Name(s):

Taraire

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. Confined to the North Island where it most common north of Auckland and Thames. However it also occurs in scattered pockets in the west south of Port Waikato to the Kawhia Harbour, inland at Pukemokemoke (near Tauhei), and in the east it occurs very locally from the eastern end of Papatea Bay to East Cape.

Habitat:

Common canopy forming tree in lowland and lower montane forests north of Auckland. Often associated with kauri (*Agathis australis*), and pohutukawa (*Metrosideros excelsa*), and on basalt rocks and soils puriri (*Vitex lucens*).

Features*:

Evergreen tree up to 22 m tall, with very broad canopy crown. Trunk to 1 m diam. Bark smooth, dark brown. Branches stout, spreading. Branchlets, young leaves, petioles and young inflorescences densely clad in reddish brown tomentum. Foliage closely alternate, erectopate, simple, leathery. Petioles (8-)10(-12) mm. Leaves (36-)50-72 (-85) x (26-)34-48(-56) mm, wide-elliptic to wide-obovate, dark green and glabrous above, bullate, glaucous below, with stout veins covered in reddish brown tomentum, margins entire, apex rounded, retuse and mucronate. Inflorescence and erect, axillary panicle up to 100 mm long. Flowers sexually perfect, 3-5 mm diam., greenish, often partially clothed in dense reddish-brown tomentum, perianth cleft into 6, stamens 12. Fruit an erect, ellipsoid to ovoid drupe (28-)30(-35) x (14-)16(-18) mm, 1-seeded, pericarp fleshy, dark purple when ripe, covered in waxy glaucous bloom.

Flowering:

(September-) November (-December)

Fruiting:

March - November

Threats:

Not Threatened

*Attribution:

Fact sheet prepared for NZPCN by P.J. de Lange 12 February 2004. Description adapted from Allan (1961) and Wright (1984).

References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Vol. I, Wellington, Government Printer.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 2009 Vol. 11 No. 4 pp. 285-309

Wilcox, M.D. 2001. Establishment of forest monitoring plots in Kirks Bush Papakura with special reference to Taraire (*Beilschmiedia tarairi*). *Auckland Botanical Society Journal*, 56: 76-79.

Wright, A. E. 1984: *Beilschmiedia* Nees (Lauraceae) in New Zealand. *New Zealand Journal of Botany* 22: 109-125.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1540



Caption: Wenderholm

Photographer: John Barkla



Caption: Taraire leaves

Photographer: DoC

Carex virgata

Common Name(s):

swamp sedge, pukio, toitoi, toetoe

Current Threat Status (2012):

Not Threatened

Distribution:

Indigenous. New Zealand: North, South, Stewart and Chatham Islands.

Habitat:

Widespread from sea level to about 1000 m a.s.l. in open, swampy conditions and also in damp sites within lowland forest. In parts of the country this sedge is often the dominant carice of lowland alluvial forest.

Features*:

Rhizomatous, densely clumped to tussock-forming sedge. Rhizome 5 mm. diameter. Culms 150–900 mm. x c.1.5 mm, trigonous, grooved, harshly scabrid; basal sheaths shining, grey-brown to dark brown, sometimes black. Lvs much > culms, 0.5–1.2 m tall, 1.5–4.5 mm wide, channelled, light green, harsh and rigid, keel and margins strongly scabrid. Inflorescence a narrow 100–260 mm long panicle with stiff erect branchlets, the lower-most quite distant. Spikes, androgynous, 4–6 mm. long, sessile, grey- or yellow-brown, male flowers terminal, lower spikes on each branchlet subtended by a pale membranous bract with a long scabrid awn often > spike. Glume ± = or slightly < utricles, membranous, ovate, acute, dull brown, with a prominent pale midrib, this often scabrid in lowermost glumes. Utricles 2.0–2.5 x c.1.0 mm, plano-convex, ovoid, light grey with distinct brown nerves; tapering to a brown beak c.0.5 mm long with a bifid orifice and conspicuously denticulate margins; abruptly contracted to a narrow stipe c.0.2 mm. long. Stigmas 2. Nut slightly > 1 mm. long, biconvex, ovoid, dark brown.

Flowering:

October - December

Fruiting:

December - May

Threats:

Not Threatened

*Attribution:

Fact Sheet prepared by P.J. de Lange (10 August 2006). Description adapted from Moore and Edgar (1970)

References and further reading:

Moore, L.B.; Edgar, E. 1970: Flora of New Zealand. Vol. II. Government Printer, Wellington.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 2009 Vol. 11 No. 4 pp. 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1426



Caption: Flower of *Carex virgata*
Photographer: Wayne Bennett



Caption: Flower of *Carex virgata*
Photographer: Wayne Bennett

Carpodetus serratus

Common Name(s):

putaputaweta, marbleleaf

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. Widespread. North, South and Stewart Islands.

Habitat:

Coastal to montane (10-1000 m a.s.l.). Moist broadleaf forest, locally common in beech forest. A frequent component of secondary forest. Streamsides and forest margins.

Features*:

Monoecious small tree up to 10 m tall. Trunk slender, bark rough, corky, mottled grey-white, often knobbed due to insect boring. Juvenile plants with distinctive zig-zag branching which is retained to a lesser degree in branchlets of adult. Leaves broad-elliptic to broad-ovate or suborbicular; dark green, marbled; membranous becoming thinly coriaceous; margin serrately toothed; tip acute to obtuse. Juvenile leaves 10-30 mm x 10-20 mm. Adult leaves 40-60 mm x 20-30mm. Petioles c. 10 mm; petioles, peduncles and pedicels pubescent; lenticels prominent. Flowers in panicles at branchlet tips; panicles to 50 x 50 mm; flowers 5-6 mm diam.; calyx lobes c. 1 mm long, triangular-attenuate; petals white, ovate, acute, 3-4 mm long. Stamens 5-6, alternating with petals; filaments short. Stigma capitate, tip dark; ovules many. Fruit an indehiscent subfleshy-fleshy capsule, 4-6 mm diam., black when mature; cupped in remains of calyx. Seeds many per capsule, in 3-5 locules, small, 1-2 mm long; testa reticulate.

Flowering:

November-
March

Fruiting:

January-February (though dried fruit
present at any time)

Threats:

Not Threatened.

*Attribution:

Description adapted from Allan (1961), puriri moth information modified from Martin (2010).

References and further reading:

Allan, H.H. 1961. Flora of NZ I. Government Printer, Wellington.

Martin, N. A. (2010). Puriri moth - *Aenetus virescens* fact sheet, retrieved from the website Interesting Insects and other Invertebrates. http://nzacfactsheets.landcareresearch.co.nz/factsheet/OrganismProfile/Puriri_moth_-_Aenetus_virescens.html

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 2009 Vol. 11 No. 4 pp. 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1605



Caption: Rotoiti Mainland Island, Nelson Lakes National Park
Photographer: John Sawyer



Caption: Rotoiti Mainland Island, Nelson Lakes National Park
Photographer: John Sawyer

Coprosma macrocarpa subsp.
macrocarpa

Common Name(s):

large-seeded Coprosma

Current Threat Status (2012):

At Risk - Naturally Uncommon

Distribution:

Endemic. Confined to the Three Kings Islands. A single specimen found on Aorangi Island (Poor Knights) may be a recent introduction from the adjacent mainland, as this plant is now commonly cultivated in northern New Zealand. Naturalised in Auckland and around Wellington cities

Threats:

A local endemic, common on but confined to the Three Kings Islands. A single record from the Poor Knights Islands is probably a chance naturalisation from the nearby mainland where it is now commonly cultivated

References and further reading:

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1720



Caption: Coprosma macrocarpa subsp. macrocarpa fruits

Photographer: John Smith-Dodsworth, Ex Cult. November



Caption: A plant of Coprosma macrocarpa subsp. macrocarpa fruits

Photographer: John Smith-Dodsworth, Ex Cult. November

Coprosma robusta

Common Name(s):

karamu, glossy karamu

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. North and South Islands. Naturalised on the Chatham Islands within a small area between Waitangi and Owenga.

Habitat:

Common throughout coastal, lowland and lower montane habitats within shrublands and open sites within forest.

Features:

Shrub or small tree up to 6 m tall. Branches numerous, stout, erect to somewhat spreading. Petioles stout, 10-20 mm long. Stipules fused towards base, obtuse, glabrous with one of two prominent, black, glandular denticles. Leaves 70-120 x 30-40-50 mm, leathery, dark green above, paler green beneath, glabrous, elliptic, elliptic-oblong to broad-ovate, acute or obtuse, apex mucronate. Venation reticulated, conspicuous. Male flowers in axillary many-flowered glomerules, corolla conspicuous, lobes triangular, acute, stamens 4-5, prominent. Females in compound clusters on peduncles 10-15 mm. Calyx and corolla much reduced, stigmas prominent. Drupe dark orange (rarely yellow), 8-8 x 4-5 mm, oblong to narrow-ovoid.

Flowering:

(July-) August-September (-November)

Fruiting:

(March-) April-May (-July)

Threats:

Not Threatened

References and further reading:

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1733



Caption: Fruit of *Coprosma robusta*

Photographer: Wayne Bennett



Caption: *Coprosma robusta* (Karamu)

Photographer: Wayne Bennett

Cordyline australis

Common Name(s):

cabbage tree, ti, ti kouka, palm lily

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. Common in the North, South and Stewart Islands. Probably naturalised on the Chatham Islands.

Habitat:

Widespread and common from coastal to montane forest. Most commonly encountered on alluvial terraces within riparian forest.

Features:

Tree up to 20 m tall, trunk stout, 1.5-2 m diam, many-branched above (prior to flowering, trunk slender and solitary, branching happens after the first flowering). Bark corky, persistent, fissured, pale to dark grey. Leaves numerous (0.2-)0.3-1(-1.5) x (0.2)-0.3(-0.6) m, dark to light green, narrowly lanceolate to lanceolate, erect to erecto-patent, scarcely inclined to droop, midrib indistinct. Petiole indistinct, short. Inflorescence a panicle. Peduncle stout, fleshy 40 mm or more in diam., panicle of numerous flowers, (0.6-)1(-1.8) x .3-0.6(-0.8) m, branching to third or fourth order, these well spaced, basal bracts green and leaf-like, ultimate racemes 100-200 mm long, 20 mm diam., bearing well-spaced to somewhat crowded, almost sessile to sessile flowers and axes. Flowers sweetly perfumed, perianth 5-6 mm diam., white, tepals free almost to base, reflexed. Stamens about same length as tepals. Stigma short, trifid.

Flowering:

(September-) October-
December (-January)

Fruiting:

(December-)
January-March

Threats:

Populations have been decimated from some parts of the country due to a mysterious illness linked to a Myoplast Like Organism (MLO) which is believed to cause the syndrome known as Sudden Decline. Plants stricken with this illness suddenly, and rapidly, wilt, with the leaves failing off still green. If the bark is peeled off the base of the tree near the soil line blackened or rotten spots are typically present. Once stricken with Sudden Decline there is no cure and the trees can die within days. Recently there has been some evidence to suggest the severity of Sudden Decline is lessening.

References and further reading:

Beever, R. et al. 1996. Sudden decline of cabbage tree. *NZ Journal of Ecology*, 20(1): 53-68

Duguid, F. 1976. *Cordyline australis* at Lake Kopureherehe. *Wellington Botanical Society Bulletin*, 39: 46-47

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1744



Caption: Awhitu Regional Park, Auckland region

Photographer: John Sawyer



Caption: *Cordyline australis*

Photographer: Wayne Bennett

Dysoxylum spectabile

Common Name(s):

kohekohe, New Zealand mahogany

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. North and South Islands. In the South Island not extending much beyond the Marlborough Sounds, reaching a southern limit near the Hurunui River (Napenape).

Habitat:

Common and sometimes dominant or co-dominant tree of coastal to lowland forest.

Features:

Tree up to 15 m tall usually with abroad, spreading canopy. Trunk up to 1 m diam., branches stout, erect then spreading. Bark pale brown, under bark green. Leaves compound, imparipinnate, alternate on pulvinate petioles up to 40 mm long, leaflet pairs 4-6, (50-)-150(-200) x (20-)30(-80) mm, opposite to subopposite, bright green, yellow-green to dark green, ovate to obovate-oblong, leathery, margins somewhat undulate. Plants gynodioecious, with fixed female and inconstant males on different trees. Inflorescence a cymose, drooping, panicle arising from trunk and branches (cauliflorous). Flowers c. 30 mm diam., fleshy. Pedicels short. Calyx divided to base, lobes broad-oblong, abruptly pointed, ciliate, petals linear, 10 mm, spreading, waxy white or greenish. Capsules, woody, broad-obovoid to subglobose, 3-4-celled, c. 25 mm long, green. Seeds 2 per cell, orange or scarlet.

Flowering:

March - June

Fruiting:

April - August

Threats:

Not Threatened. However, where possum and rat numbers are high this species is not actively regenerating. Possums defoliate trees, and will heavily browse inflorescences such that few succeed in flowering and setting fruit. Rats are major seed predators. Only where control of these animals is undertaken, or on possum and rodent-free offshore islands can one see kohekohe flowering, fruiting and regenerating freely. If numbers of these introduced animals remain unchecked, it is clear that kohekohe will decline and vanish from large parts of its natural range.

References and further reading:

Duguid, F. 1985. Kohekohe *Dysoxylum spectabile* as an accidental epiphyte. Wellington Botanical Society Bulletin, 42: 11

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 11: 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1825



Caption: Colonial Knob Scenic Reserve, Porirua.

Photographer: Jeremy Rolfe



Caption: Colonial Knob Scenic Reserve, Porirua.

Photographer: Jeremy Rolfe

Entelea arborescens

Common Name(s):

Whau

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. Three Kings, North (including Little and Great Barrier Islands) and South Islands. In the North Island, whau is locally common from Te Pahi to about Kawhia and Mahia Peninsula south of there it is known from a few sites in the eastern Wairarapa, at Paekakariki and Wellington. In the South Island it is confined to the Golden Bay area of North-West Nelson. Whau naturalises easily and has established south of these stations from both rural and urban plantings.

Habitat:

Coastal to lowland forest or shrubland. Usually in open sites such as around recent slips, tree falls, cliff faces, boulder slopes, sand dunes or on the margins of streams, rivers, lagoons and lakes. Mostly near the coast however it may occur well inland in some places e.g., the Waikato River near Hamilton, Rotorua. Some inland and southern North island occurrences are thought to be derived from deliberate plantings by Maori.

Features*:

Shrub or small spreading tree up to c. 8 m tall; trunk up to 0.25 m dbh; wood-weight very light; bark firm, grey, tearing in long fibrous strips when cut. branches numerous, upright than spreading; branchlets, leaves, petioles, inflorescences densely clad in soft whitish branched hairs; leaf-scars oval or lunate. Leaves alternate, softly membranous (wilting readily if picked), green, bright green to yellow green, ± glossy, venation distinct when fresh or dry; petioles 80-300 mm long, stout; stipules linear-acuminate, ± persistent. Lamina 50-100-150(-300) × 50-100-150-(260) mm, obliquely very broad-ovate, abruptly acuminate, cordate at base, margins doubly crenate-serrate, sometimes obscurely lobed, 3-5-7-subpalmately lobed. Inflorescence a subumbellate many-flowered cyme. Flowers (3-)4-5-merous. Peduncle 100-300 mm long, stout, pedicels 10-40 mm long. Sepals free, 8-10-12 mm long, narrowly lanceolate to triangular, acuminate; petals (3)-4-5, 10-30 × 10-30 mm, orbicular to suborbicular, white, crumpled. Stamens numerous, mostly free sometimes connate at base, filaments 10-18(-20) mm long, white, anthers versatile, yellow. Ovary 5-10 mm long, broadly to narrowly globose or ovoid, hispid, 5-7-locular, ovules numerous, style simple, stigma ± globular to broadly capitate, fringed or toothed. Fruit a bristly capsule 20-35 mm diameter, subglobose to globose, black to charcoal when ripe, invested by numerous, rigid, spinose, black to charcoal coloured hairs 15-25 mm long. Seeds numerous, 1.9-2.9 mm long, obovate, elliptic to broadly elliptic, glabrous, surface granular, orange-yellow, pale brown, or orange-brown. Description of seeds by Webb & Simpson (2001).

Flowering:

August - November

Fruiting:

December - June

Threats:

Not Threatened. However, recent field work gathering samples for a Marsden study into the possible past use of whau by Maori indicates that whau is much less common in the North island than it once was. browsing pressure from cattle, goats and horses, clearance of coastal scrub of housing and the spread of invasive woody shrubs and trees into many northern coastal areas may be threatening some populations.

*Attribution:

Fact Sheet Prepared for NZPCN by: P.J. de Lange 10 February 2011. Description of seeds by Webb & Simpson (2001).

References and further reading:

Webb, C.J.; Simpson, M.J.A. 2001: Seeds of New Zealand Gymnosperms and Dicotyledons. Christchurch, Manuka Press.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 11: 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1377



Caption: Fruits, Auckland
Photographer: John Barkla



Caption: Awhitu Regional Park, Auckland
Photographer: John Sawyer

Gahnia lacera

Common Name(s):

cutty grass

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. North Island from Te Pahi south to about Awakino in the west and East Cape in the East.

Habitat:

Coastal to lowland (rarely extending up to 500 m a.s.l. in mountain ranges close to the sea). Colonising a variety of substrates which may be seasonally waterlogged though otherwise dry. Usually found in scrub or open forest.

Features*:

Stout, bambusiform, perennial sedge forming dense, yellowish-green tufts. Rhizome shortly creeping, 5-8 mm diameter, very hard, lignaceous, long persistent when dead. Culms 0.6-2.0 m, 2-4 but up to 6 mm diameter at the base. Leaves numerous, almost all cauline and = or > culms; lamina up to 380 mm long and 9 mm wide, yellow-green, green or dark green, flat or involute, glabrous, margins scabrid; sheaths all closely appressed to and enclosing base of culm, rugose above, maturing dark brown to almost black, smooth and glossy towards the node. Panicle 20-600 x 30-60 mm, rigid, usually erect, many-flowered, light brown with branchlets 20-160 mm long, usually in dense though more or less distant clusters along the axis of the panicle. Spikelets 1-flowered, 8 mm long, alternate on the branchlets, sessile or shortly stalked. Glumes 4-5; the outer 2-3 light red-brown; inner glumes pale cream, with a red lacerate apex. Stamens 4. Style-branches 3. Nut 3.5-4.5 mm long, slightly < 2 mm diameter, oblong-ellipsoid or oblong-obovoid, lustrous black, with a minute apiculate apex, endocarp obscurely transverse striate within.

Flowering:

July -
November

Fruiting:

Fruits may be present throughout the
year

Threats:

Not Threatened

*Attribution:

Fact Sheet prepared by P.J. de Lange (30 October 2005). Description adapted from Moore and Edgar (1970)

References and further reading:

Gardner, R.O. 1996. *Gahnia pauciflora* and *G. procera* and a note on *G. lacera*. Auckland Botanical Society Journal, 51: 7-10.

Moore, L.B.; Edgar, E. 1970: Flora of New Zealand. Vol. II. Government Printer, Wellington.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 11: 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1903



Caption: *Gahnia lacera*

Photographer: Wayne Bennett



Caption: *Gahnia lacera*

Photographer: Wayne Bennett

Kunzea robusta

Common Name(s):

manuka, kanuka, kopuka, rawirinui, maru, manuka rauriki

Current Threat Status (2013):

Not Threatened

Distribution:

Endemic. New Zealand: North and South Islands.

Habitat:

Coastal to lowland shrubland, regenerating forest and forest margins, also present in montane forest, ultramafic shrubland and very occasionally present in subalpine shrubland (up to 900 m a.s.l.).

Features*:

Trees 8–30 m tall. Trunk 1–6, 0.10–1.0 m d.b.h. Bark stringy, or coarsely tessellated, coriaceous, firmly attached above, detaching basally, often hanging semidetached; peeling upwards along trunk in narrow to broad, tabular strips up to 4 m long. Branches initially erect, soon arching outwards and spreading; branchlets numerous, slender; sericeous, indumentum copious, hairs either long or short antrorse-appressed; if long, then weakly flexuose 0.15–0.38 mm long; if short, not flexuose, 0.09–0.15 mm long. In eastern Coromandel Peninsula and coastal East Cape to Mahia Peninsula, branchlet indumentum in mixtures of divergent 0.03–0.08 mm long hairs, and sparse, 0.1–0.2 mm long, antrorse-appressed hairs. In the Rangitikei region, branchlet hairs of seedling and juveniles divergent, short 0.04–0.10 mm long. Leaves sessile to shortly petiolate, light green or dark green above, paler beneath; oblanceolate, broadly oblanceolate, broadly lanceolate, lanceolate to linear-lanceolate, rarely elliptic to obovate; apex subacute to acute, rarely obtuse, rostrate or shortly apiculate, base attenuate to narrowly attenuate; lamina margin initially finely covered with a thin, interrupted band of spreading to antrorse-appressed hairs not or rarely meeting at apex; hairs shedding with age. Lamina of juvenile plants from coastal areas and northern North Island 14.6–28.4 × 1.6–2.5 mm; from inland areas, 3.2–6.3 × 0.7–1.5 mm; adult lamina of plants from coastal areas and northern North Island 4.9–20.1 × 0.9–3.0 mm; from inland areas, 5.8–12.3 × 1.2–2.2. Inflorescence mostly a compact corymbiform to shortly elongate 1–30-flowered botryum up to 60 mm long; extending near end of flowering season as an 4–12-flowered, elongate botryum up to 80 mm long; Pherophylls deciduous or persistent; squamiform grading into foliose; squamiform pherophylls 0.4–1.2 × 0.3–0.6 mm, broadly to narrowly deltoid or lanceolate, apex acute, subacute to obtuse, margins finely ciliate; foliose pherophylls 6.0–17.9 × 1.1–1.8 mm, elliptic, oblanceolate, broadly lanceolate to lanceolate, apex obtuse, base attenuate; margin densely covered by antrorse-appressed hairs. Pedicels 1.2–5.2 mm long at anthesis. Flower buds pyriform to obconic, apex flat or weakly domed prior to bud burst; calyx valves not meeting. Flowers 4.3–12.0 mm diameter. Hypanthium 2.1–4.1 × 3.0–5.2 mm, broadly obconic to turbinate, sometimes cupular, rim bearing five persistent calyx lobes. Hypanthium surface when fresh faintly ribbed and sparingly dotted with pink or colourless oil glands, these drying dull yellow-brown or brown; either finely pubescent with the ribs and veins conspicuously covered in longer silky, antrorse-appressed hairs, or glabrous; hypanthium similar when dry though with the ribs more strongly defined and clearly leading up to calyx lobes. Calyx lobes 5, coriaceous, 0.52–1.1 × 0.60–1.4 mm, broadly ovate, ovate-truncate to broadly obtuse, glabrate. Receptacle green or pink at anthesis, darkening to crimson after fertilisation. Petals 5–6, 1.5–3.8 × 1.3–3.6 mm, white, rarely pink, orbicular, suborbicular to ovate, apex rounded to obtuse, oil glands colourless. Stamens 15–58 in 2 weakly defined whorls, filaments white. Anthers 0.38–0.63 × 0.18–0.32 mm, ellipsoid to ovoid-ellipsoid or deltoid. Pollen white. Anther connective gland prominent, light pink, salmon pink, yellow to orange when fresh, drying dark orange, orange-brown or dark brown, spheroidal, finely rugulose or papillate. Ovary 5–6 locular. Style 2.0–3.5 mm long at anthesis, white or pinkish-white; stigma broadly capitate, flat, greenish-white or pale pink, flushing red after anthesis. Fruits 2.2–4.6 × 3.2–5.3 mm, maturing greyish white, obconic, broadly obconic to ± turbinate, rarely cupular; hairy, (rarely glabrous). Seeds 0.9–1.1 × 0.35–0.48 mm, oblong, oblong-obovate, oblong-elliptic; testa semi-glossy, orange-brown to dark brown, surface coarsely reticulate.

Flowering:

August–June

Fruiting:

Jul–May

Threats:

Not Threatened.

*Attribution:

Fact Sheet prepared for NZPCN by P.J. de Lange 10 September 2014. Description modified from de Lange (2014).

References and further reading:

de Lange, P.J. 2014: *A revision of the New Zealand Kunzea ericoides (Myrtaceae) complex. Phytokeys 40: 185p* doi: 10.3897/phytokeys.40.7973.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=7644



Caption: Otari Wilton's Bush, Wellington.

Photographer: Jeremy Rolfe

Leptospermum scoparium var. *scoparium*

Common Name(s):

manuka, tea tree, kahikatoa

Current Threat Status (2012):

Not Threatened

Distribution:

Indigenous to New Zealand and Australia. Most Australian forms of *L. scoparium* do not match the range seen in New Zealand. However, plants from Tasmania are very similar to, if not identical with some South Island forms, differing mainly by their wider leaf base, and longer, more pungent leaf apex. Manuka was also collected once from Rarotonga by Thomas Cheeseman in the 1800s. It has not been found there since, and is assumed to have been a failed introduction. Further study using DNA sequencing is underway to resolve the status of *L. scoparium* forms both here and in Australia.

Habitat:

Abundant from coastal situations to low alpine habitats.

Features*:

Decumbent shrub, subshrub, shrub, or small tree up to 5 m in height and in decumbent forms 2-4 m across. Bark light grey to charcoal grey, peeling in long papery flakes, these curling with age. Wood red. Branches numerous erect, spreading or decumbent, arising from base, sometimes sprouting adventitious roots and/or layering on contact with soil. Young branches, young leaves and flower buds densely to sparingly clad in long silky, white hairs. Leaves leathery, pale to dark green, glabrescent to glabrous, linear-filiform, narrowly lanceolate, lanceolate, oblanceolate, to elliptic or obovate (5-)10-15(-20) x 1-2-5(-8) mm, invariably apex drawn out into a long stiff, pungent point, midrib usually distinct sometimes obscure, leaf margin finely crenate, veins simple, scarcely branched. Flowers solitary in leaf axils, (8-)10-20(-25) mm diam. Receptacle dark red, crimson or pink. Petals white, sometimes flushed pink or dark red. Stamens numerous.

Flowering:

Throughout the year

Fruiting:

The capsules are long persistent so invariably mature plants always possess at least some capsules.

Threats:

Not threatened, though some stands are at risk from clearance for farmland or through felling for firewood.

*Attribution:

Fact Sheet prepared for NZPCN by P.J. de Lange 1 February 2004. Description by P.J. de Lange.


References and further reading:

Gardner, R. 2002. Notes towards an excursion Flora .Manuka *Leptospermum scoparium* myrtaceae. Auckland Botanical Society Journal, 57: 147-149

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 11: 285-309

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=2302



Photographer: © John Braggins



Caption: Flowers of *Leptospermum scoparium* var. *scoparium*

Photographer: Wayne Bennett

Melicytus ramiflorus

Common Name(s):

mahoe, hinahina, whitey wood

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic subspecies. Three other subspecies occur, one endemic to Norfolk (probably a different species), one to Fiji and one to Samoa. In addition forms from Raoul Island (Kermadec Islands Group) and the Three Kings and eastern Northland may warrant formal recognition. Research into this variation is in progress.

Habitat:

Abundant small tree of coastal, lowland, and lower montane forests throughout the country.

Features:

Shrub or small tree up to 15 m tall. Trunk 1 or more, 0.6-0.8 m diam, typically much branched from near base. Wood soft, white. Bark greyish-white, underbark bright green. Branchlets numerous, twiggy, rather brittle. Petioles 20 mm or more long. Leaves, firmly fleshy, 50-150 x 30-50 mm, light or dark green, lanceolate-oblong to elliptic oblong, apex acute to acuminate (rarely obtuse), leaf margins coarsely serrated (very rarely subentire, or irregularly coarsely toothed). Inflorescence 2-10 flowered fascicles arising from branchlets or leaf axils. Flowers 3-4 mm diam., female or inconstant male (flowers types on separate plants) borne on slender pedicels 5-10 mm long. Bracts subtending flowers, calyx lobes minute, petals greenish-yellow, yellow (rarely cream), lanceolate, apex obtuse. Anthers sessile, stigma 4-6-lobed. Fruit a violet, dark blue or purple berry, 4-5 mm diam., obovoid to globose. Seeds 3-6 per berry.

Flowering:

November - February

Fruiting:

November - March

Threats:

Not Threatened

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=973



Caption: Carter Scenic Reserve, Wairarapa

Photographer: John Sawyer



Caption: Carter Scenic Reserve, Wairarapa

Photographer: John Sawyer

Metrosideros excelsa

Common Name(s):

Pohutukawa, New Zealand Christmas tree

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. New Zealand: Three Kings Islands and North Island from North Cape to about Pukearuhe, (northern Taranaki) in the west and near Mahia Peninsula (in the east). However, exact southern limit is difficult to ascertain as it has been widely planted and there is evidence that old time Maori cultivated the tree in some southerly areas. Found inland around the Rotorua Lakes and at Lake Taupo - though these occurrences could stem from Maori plantings (though the association of other normally coastal species around these lakes argues against this). Now widely planted throughout the rest of New Zealand (especially around Nelson, the Marlborough Sounds, the Kaikoura Coast and on the west coast to about Hokitika).

Habitat:

Coastal forest and on occasion inland around lake margins. Also in the far north occasionally an associate of kauri forest. In some northerly locations it forms forest type in its own right - this forest is dominated by pohutukawa, other associates often include tawapou (*Pouteria costata*), kohekohe (*Dysoxylum spectabile*), puriri (*Vitex lucens*), karaka (*Corynocarpus laevigatus*), and on rodent-free offshore islands the frequent presence of coastal maire (*Nestegis apetala*), and milk tree (*Streblus banksii*) suggests these species too may once have been important in mainland examples of pohutukawa forest.

Features*:

Tree up to 20 m tall with canopy spread of 10-50m. Specimens typically multi-trunked from base, trunks up to 2 m diameter, branches spreading, and often arching, sometimes looping over ground, and/or bearing "brooms" of aerial adventitious roots. Branchlets numerous, twiggy and long-persistent. Bark firm, persistent and difficult to detach, often deeply furrowed, grey to grey-brown, somewhat corky. Young branchlets tomentose, being covered in fine, deciduous, greyish-white hairs. Leaves of all but water shoots leathery, 25-120 × 25-60 mm, elliptic, oblong, rarely lanceolate, apex acute or obtuse, dark olive-green, undersides thickly clad in white tomentum, adaxial surface at first distinctly tomentose but hairs shedding with leaf maturation. Flowers borne on stout, tomentose pedicels crimson, orange, pink, yellow (or very rarely white). Hypanthium obconic, calyx lobes triangular (deltoid).

Flowering:

(August-) November-December (-March)

Fruiting:

(January-) March-April (-May)

Threats:

Like all New Zealand tree *Metrosideros*, pohutukawa is most at risk from possum (*Trichosurus vulpecula*) browse. These can seriously damage and even kill trees. Often where their browsing occurs within sites of unrestricted stock and vehicle access, pohutukawa forest is in danger of becoming locally extinct. It does remain common over large parts of its range, a situation being greatly improved by the efforts of people encouraged by the national coordination of Project Crimson - a non profit organisation set up to protect, enhance and/or establish pohutukawa forest, as well as promote the species use, and its conservation.

*Attribution:

Fact sheet prepared for NZPCN by: P.J. de Lange (4 January 2004). Description adapted from Allan (1961).

References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Vol. I. Wellington, Government Printer.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=975



Caption: Wellington

Photographer: John Sawyer



Caption: *Metrosideros excelsa*

Photographer: Wayne Bennett

Myrsine australis

Common Name(s):

Red mapou, red matipo, mapau, red maple

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. Three Kings, North, South and Stewart Islands.

Habitat:

Common tree of regenerating and mature forest in coastal to montane situations. Often common on northern offshore islands.

Features*:

Shrub or small tree up 6 m tall. Trunk stout, 0.2-0.6 m diam. Bark dark black or purple-black, red on younger branches. Branchlets numerous erect to spreading, very leafy. Petioles stout, fleshy, 5 mm long, often red or green mottled red. Leaves 30-60 x 15-25 mm, dark green to yellow-green variously mottled or blotched with red, or purple spots, leathery, glabrous except for finely pubescent mid vein, obovate-oblong to broad-elliptic, apex obtuse, margins entire, strongly undulate, rarely flat. Inflorescence a fascicle, usually numerous and crowded, produced along branchlets and in leaf axils. Fixed female and inconstant male flowers on different plants, 1.5-2.5 mm diam., white, cream or pale green. Pedicels short, stout, dark red or purple-black. Calyx-lobes 4, sometimes heavily reduced, long persistent. Petals 4, lanceolate, obtuse, free, revolute. Fruit a 1-seeded drupe, 2-3 mm diam., purple-black to black when mature.

Flowering:

August - January

Fruiting:

September - May

Threats:

Not Threatened

*Attribution:

Fact Sheet Prepared for NZPCN by: P.J. de Lange 28 October 2009.
Description based on Allan (1961)

References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Vol. I. Wellington, Government Printer.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1007



Caption: Male flowers. Rimutaka Forest Park.

Photographer: Jeremy Rolfe



Caption: Male flowers. Rimutaka Forest Park.

Photographer: Jeremy Rolfe

Phormium tenax

Common Name(s):

flax, harakeke, korari (maori name for inflorescence).

Current Threat Status (2012):

Not Threatened

Distribution:

Indigenous to New Zealand and Norfolk Island. A broad circumscription has been adopted here - many botanists feel that plants from the Chatham Islands could be distinguished at species rank from the mainland New Zealand species, other distinctive variants occur on the Three Kings and outer Hauraki Gulf Islands, and along the Kaikoura coast. Norfolk Island plants though uniform differ in subtle ways from the New Zealand forms of *P. tenax*. Further study into this variation is underway.

Habitat:

Common from lowland and coastal areas to montane forest, usually but not exclusively, in wetlands and in open ground along riversides.

Features:

Stout liliaceous herb, 1-5(-6) m tall. Leaves numerous, arising from fan-like bases. Individual leaves rather stiff at first, but becoming decurved, somewhat pendulous or "floppy" in upper half to a third, 1-3 x 50-120 mm, usually blue-grey (glaucous) or dark green, lamina margin, entire, somewhat thickened and pigmented black, dark red, pink, yellow or cream. Inflorescence 5(-6) m tall, somewhat woody and fleshy when fresh, long persistent, drying charcoal grey or black, with the fibrous interior becoming progressively more exposed. Peduncle 20-30 mm diam., erect, dark grey-green or red-green, glabrous. Flowers 25-50 mm long, tubular, predominantly dull red but may also be pink or yellow; tips of inner tepals slightly recurved. Ovary erect. Capsules 50-100 mm long, dark green, red-green or black, trigonous in cross-section, erect, abruptly contract at tip, not twisted, initially fleshy becoming woody with age, long persistent. Seeds 9-10 x 4-5 mm, black, elliptic, flat and plate-like, margins frilled or twisted.

Flowering:

(September-) October-November (-January)

Fruiting:

(November-) December (-March)

Threats:

Not threatened although see the discussion below about flax dieback. This die back phenomenon is characterised by abnormal yellowing of the leaves and may result in collapse of flax plants or whole populations.

References and further reading:

Boyce, et al. 1951. Preliminary note on yellowleaf disease. NZJ of Science and Technology, 32(3): 76-77

Scheele, S. 1997. Insect pests and diseases of harakeke, Manaaki Whenua Press


For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=2219



Caption: *Phormium tenax*

Photographer: Wayne Bennett



Caption: Flowers of *Phormium tenax*

Photographer: Wayne Bennett

Pittosporum crassifolium

Common Name(s):

Karo

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. New Zealand, Great Barrier and North Island. In the North indigenous from Te Pahi south to about White Cliffs, and East Cape. Widely naturalised further south to Wellington. Naturalised in the South, Stewart and Chatham Islands. Also naturalised on Norfolk Island, and in Hawaii.

Habitat:

Coastal and offshore islands. Favouring steep slopes, cliff faces, boulder beaches, rock stacks and the margins of petrel burrowed land. Sometimes forms major canopy dominant on offshore islands, and on occasion can be a significant component of dune forest. Often an urban weed because its fruits/seeds are avidly taken by indigenous and exotic birds and dispersed widely.

Features*:

Gynodioecious shrubs to small trees 1-10 m tall. Trunk stout, grey-black, often distinctly lenticillate. Branches and branchlets erect, dark grey-black or brown, immature branchlets densely invested in grey-white or white tomentum, this maturing black. Leaves alternate, usually densely crowded toward branch and branchlet apices. Petioles 4-14 x 1-3 mm, grey-white to grey-black tomentose. Leaves 30-100 x 10-30 mm, obovate to oblanceolate, apices obtuse to acute, base attenuate, margins entire, both surfaces densely white, grey-white or brown tomentose when young, soon glabrate above but remain densely covered in dirty white or grey-white, appressed tomentum beneath, very coriaceous, margins thickened and often strongly revolute, surfaces often blistered with insect galls. Flowers in terminal 1-10-flowered fascicles; pedicels 6-50 mm, accrescent in fruit, tomentose, subtended by a whorl of leaves and numerous, 3-15 mm long, caducous, brown-tomentose, ciliate bud scales. Sepals 7-11 x 1.5-3 mm, oblong to linear-lanceolate, acute, greyish-white, dirty white or brown tomentose on outer surfaces, inner surface only toward the middle, margins ciliate. Petals 10-16 x 3-5 mm, oblanceolate to lanceolate, subacute, free to base, recurved at apices, dark red, purple, yellow, pink or white; stamens 5-9 mm long, anthers 1-3 x 0.5-1.5 mm, sagittiform to elliptic-oblong. Ovary 3-6 x 2-5 mm, white or grey-white tomentose; style 3-2.5 mm long, stigma capitate or 3-lobed truncate. Capsules woody, 10-30 x 10-30 mm, (2-)3(-4)-valved, woody, trigonous, sometimes 2-4-lobed

Flowering:

August - October

Fruiting:

September - August (Old fruits persist on trees)

Threats:

Not Threatened. However, the fruits are eaten by rats, and on rodent infested offshore islands this species rarely regenerates.

*Attribution:

Fact sheet prepared for NZPCN by P.J. de Lange 30 August 2006. Description adapted from Cooper (1956).

References and further reading:

Cooper, R.C. 1956: The Australian and New Zealand species of *Pittosporum*. *Annals of the Missouri Botanical Garden* 43: 87-188

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1133



Caption: Masterton

Photographer: John Barkla



Caption: Meola Reef, Westmere, Auckland

Photographer: John Sawyer

Sophora microphylla

Common Name(s):

Kowhai, weeping kowhai, small-leaved kowhai

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. Throughout the main islands of New Zealand but scarce in parts of Northland.

Habitat:

In the North Island, especially the northern half this is a species of mainly riparian forest. South of about Hamilton it can be found in a diverse range of habitats from coastal cliff faces and associated wetlands to inland grey scrub communities. Scarce to absent over large parts of the eastern North Island from about East Cape south to the northern Wairarapa.

Features*:

Tree up to 25 m tall, usually a single trunk. Branches weeping, and spreading. Juveniles divaricating and/or strongly flexuose, and interlacing. Leaves on seedlings sparsely to moderately leafy, 3-5.8 x 2.3-4.9 mm, broadly obovate to orbicular, glabrous to sparsely pubescent, distant, not crowded or overlapping. Adult leaves up to 150 mm long, imparipinnate, moderately to sparsely hairy, hairs, straight, appressed. Leaflets 30-50, not crowded or overlapping, distant, 4.5-12.5 x 2.3-5.7 mm, elliptic, broadly elliptic, obovate to ovate, sometimes orbicular, distal and proximal leaflets of similar size. Inflorescences racemose with up to 7 flowers. Calyx 5-11 x 7-10 mm, cupulate. Flowers yellow, keel petal blade 18-50 x 7-13 mm, wing petal blade 18-50 x 6-11 mm, standard petal blade 20-35 x 14-25 mm; petals with distinct claws 4-8 mm long. Fruit 50-200 mm long, 4-winged, brown, with up to 12 seeds. Seeds 5.5-8.5 x 4-5.5 mm, oblong, elliptic to orbicular, yellow to light yellow-brown.

Flowering:

(May-) August-October

Fruiting:

October - May

Threats:

The main threat that faces all wild New Zealand kowhai species is the risk posed through planting for revegetation and horticultural purposes of hybrid material, foreign species, such as the Chilean Pelu (*S. cassioides*) and also of kowhai species outside their natural range. In many places *S. microphylla* occurs as isolated stands within otherwise cleared alluvial forest, and in this situations the loss of trees over time is inevitable. The species is genuinely uncommon in Northland, and in that area inadequately represented within reserves and other conservation land.

*Attribution:

Fact Sheet prepared for NZPCN by P.J. de Lange (31 July 2004).
Description adapted from Heenan et al. (2001).

References and further reading:

Anonymous. 1944. Kowhai. *Wellington Botanical Society Bulletin* 9: 4-5

Duguid, F. 1971. Germination of kowhai at Hokio beach. *Wellington Botanical Society Bulletin* 37: 65-66.

Heenan, P.B.; de Lange, P. J.; Wilton, A. D. 2001: *Sophora* (Fabaceae) in New Zealand: taxonomy, distribution, and biogeography. *New Zealand Journal of Botany* 39: 17-53


For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1302



Caption: Bark, Dunedin Botanic Gardens

Photographer: John Barkla



Caption: *Sophora microphylla* (Kowhai)

Photographer: Wayne Bennett

Vitex lucens

Common Name(s):

puriri

Current Threat Status (2012):

Not Threatened

Distribution:

Endemic. New Zealand: Three Kings Islands and North Island from Te Pahi to Taranaki, Mahia Peninsula and the northern Hawkes Bay. Puriri is, as a rule, scarce south of about Opotiki and Kawhia.

Habitat:

In the northern part of its range Puriri is a common co-dominant with Taraire (*Beilschmiedia tarairi*) and karaka (*Corynocarpus laevigatus*) especially on rich fertile soils derived from basaltic and basaltic-andesitic igneous rocks. South of the northern Bay of Plenty and Raglan Harbours it is rarely found inland and is more commonly found in coastal forest where it co-habits with pohutukawa (*Metrosideros excelsa*) and karaka. Puriri is also an important forest tree on many of the smaller islands of the Hauraki Gulf, where it may at times be the canopy dominant.

Features*:

Tree up to c. 20 m. tall with a broad spreading canopy; trunk up to c.1.5 m. diameter; bark grey-brown, firm, flaking in small irregular-shaped shards. Branches stout, spreading; branchlets 4-angled, green. Leaves opposite, glabrous, coriaceous, compound, on petioles up to 110 mm long; Leaflets 3-4-5, somewhat undulose, adaxially dark green, glossy, abaxially lighter green, mat; basal one or pair of leaflets usually much smaller than the terminal 3, digitate; lamina of 3 main leaflets 50-140 × 30-60 mm; elliptic-oblong to obovate, abruptly acute to subacuminate, margin entire. Domatia (pit-type) present at axils of costa and main veins. Inflorescence in axillary, dichotomous, (4)-10-15-flowered panicles. Calyx cupular, minutely 5-toothed; corolla dull red, pink or white, pubescent, 2-lipped, c.25-35 mm long. Upper lip entire or bifid, lower deflexed, 3-lobed. Style slender, bifid, c.25 mm long. Drupe 20-26 mm diameter subglobose, bright red, pink or white.

Flowering:

May - October

Fruiting:

January - October

Threats:

Not Threatened. However, in some parts of Northland puriri "die-back" has been observed (the exact causes of which are much debated). Puriri is at times heavily browsed by possums, to such an extent that trees can die.

*Attribution:

Factsheet prepared for NZPCN by P.J. de Lange 9 February 2011. Description adapted from Allan (1961).

References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Vol. I, Wellington, Government Printer.

For more information, visit:

http://nzpcn.org.nz/flora_details.asp?ID=1359



Caption: In cultivation.

Photographer: John Braggins



Caption: In cultivation.

Photographer: John Braggins

Definitions of botanical terms

A glossary has been provided below with definitions for many of the botanical terms used in the species descriptions.

Glossary

Term	Definition
Abaxial	Facing away from the stem of a plant (especially denoting the lower surface of a leaf).
Acerose	Narrow with a sharp stiff point.
Achene	A simple, dry, one-seeded (one-celled) fruit
Acicular	Needle-shaped.
Acidic	Having a low pH, opposite of basic or alkaline.
Acroscopic	Pointing towards, or on the side of, the apex
Acuminate	Gradually tapered to a point. Sharply pointed.
Acute	Pointed or sharp, tapering to a point with straight sides.
Adnate	Fusion of unlike parts, e.g. stamens fused to petals.
Adventive	A plant that grows in the wild in New Zealand but which was introduced to the country by humans.
Agglutinated	Stuck together.
Allelopath	An organism that releases compounds that are toxic to other species.
Allelopathy	The release by an organism of compounds that are toxic to other species.
Alternate	Attached singly at each node but changing from one side of a stem to the other.
Alveolate	Honeycombed with ridged partitions.
Amplexicaul	clasping or surrounding the stem
Anamorph	Asexual fruiting stage, usually of an ascomycete fungus.
Anastomosing	Rejoining after branching, as in some leaf veins.
Annual	A plant that completes its complete life cycle within the space of a year
Annual evergreen	Plants that lose their over-wintering leaves rapidly in the first half of the growing season. Annual evergreens never present a leafless appearance, but are closer in a functional sense to a deciduous plant than they are to multi-annual evergreens.
Annulus	Line of thickened cells that governs the release of spores from a sporangium
Anterior	Towards the front.
Anther	The pollen-bearing portion of the stamen.
Antheridium	Male reproductive organ formed on the prothallus of a fern
Anthesis	When the flower is fully developed and functioning. The time of pollination or bloom.
Apex	Tip; the point furthest from the point of attachment.
Apices	Plural of apex. Tip, the point furthest from the point of attachment
Apiculate	Bearing a short slender and flexible point.
Apiculus	A small, slender point.
Apomixis	A form of reproduction whereby seed is formed without the usual mode of sexual fusion
Appressed	Pressed against another organ or surface.
Aquatic	Growing, or living in, or frequenting water. Applied to plants and animals and their habitats. Opposite of terrestrial (land living).
Archegonium	Female reproductive organ of a fern formed on the prothallus
Arcuate	Curved into an arch.
Aril	An often fleshy appendage on the outside of a seed.
Artificial thinning	Selectively removing vegetation to create gaps to facilitate natural invasion of native plants, or to plant later successional plants.
Ascending	Growing obliquely upward.
Asexual	Vegetative reproduction, lacking sexual involvement by sperm or egg cells
Attenuate	Narrowing gradually
Auricle	A small, ear-shaped appendage.
Auriculate	Bearing a small, ear-shaped appendage.
Autogamous	Self-fertilising flowers.
Autotrophic	Of or relating to organisms (as green plants) that can make complex organic nutritive compounds from simple inorganic sources by photosynthesis
awn	A stiff or bristle like projection often from the tip or back of an organ
Axil	The upper angle between the leaf and the stem.
Axis	The longitudinal supporting structure around which organs are borne, e.g., a stem bearing leaves.
Barbellate	Barbed, having or covered with protective barbs or quills or spines or thorns or setae
Basal	At the base.
Basisopic	Pointing towards the base
Beak	A prominent extension of an organ
Bifid	Deeply split into two lobes.
Bifurcate	Divided into two.

Term	Definition
Biosecurity	Preventing, eradicating, controlling and managing risks posed by pests and diseases.
Biotic	Pertaining to the living parts of the environment
Bipinnate	With each primary pinna divided to the midrib into a secondary pinna
Biserrate	Doubly serrate.
Blade	The flattened part of a leaf.
Blunt	Not pointed at the ends
Bog	A quagmire covered with specialised plants including sphagnum moss, grasses, sedges, rushes, sundews, umbrella ferns and other plants; has wet, spongy ground, a marsh-plant community on wet, very acid peat. Fed only by rainfall.
Bottleneck	A genetic term; refers to the fact that in smaller populations there could be lower genetic variability
Brachyblasts	Short shoots
Bract	A reduced leaf or leaf-like structure at the base of a flower.
Bracteate	Bearing bracts: leaves or leaf-like structure reduced at the base of a flower.
Bracteolate	With small bracts.
Bracteole	A small bract.
Bracteoles	Bracts directly below the flower
Brevideciduous	Brief (1 month or less) loss of most leaves from the canopy just before flowering or during flushing of a new cohort of leaves.
Bryophyte	Plant group including mosses, liverworts and hornworts
Bryophytes	Plant group including mosses, liverworts and hornworts
Bulbil	A bud produced vegetatively on the stem or frond that is capable of breaking off and growing into a new plant
Bullate	With rounded projections covering the surface as if blistered
Caespitose	Growing in dense tufts
Calli	Circular, warty, stalked thickenings commonly found on the lip (labellum) of the orchid (plural of callus).
Callose	Hardened or thickened.
Callus	Stalked thickening on the lip (labellum) of an orchid.
Calyx	The group of sepals, or outer floral leaves, of a flower
Campanulate	Bell-shaped.
Canaliculate	With longitudinal channels or grooves.
Canopy	The uppermost cover formed by the branches and leaves of trees or the spread of bushes, shrubs and ground covers.
Canopy closure	Stage where canopies of shrub and tree species meet.
Canopy manipulation	Selectively removing vegetation to create gaps to facilitate natural invasion of native plants, or to plant later successional plants.
Capillary	Hair-like
Capitula	Plural of capitulum: A dense head-like inflorescence of many flowers as occurs in most Asteraceae (daisies)
Capitulum	A dense head-like inflorescence of many flowers as occurs in most Asteraceae (daisies)
Capsule	A dry fruit formed from two or more fused carpels that splits open when ripe.
Carbon sinks	Carbon locked away, or sequestered e.g. by trees
Carpel	One unit of the female part of a flower that consists of a basal seed-bearing ovary joined to a receptive stigma by a stalk-like style.
Cauda	Tail-like appendage. (pl. caudae; adj. caudate)
Caudex	The axis of a woody plant, esp. a palm or tree fern, comprising the stem and root.
Cauline	Belonging to the stem, as in cauline leaves emerging from the stem.
Cerise	Bright or deep red.
Chartaceous	Having a papery texture.
Chlorophyll	The green pigment of plants.
Chlorotic	Lacking chlorophyll, therefore yellowish, suffering from chlorosis.
Cilia	Short small hair-like structures on a cell or microorganism
Ciliate	With small hairs (cilia).
Ciliolate	Diminutive of ciliate, i.e., having very small hairs
Cladode	Flattened stem with the function of a leaf
Cladodes	Usually flattened, photosynthetically active branches, these may be leaf-like (e.g., Phyllocladus) or branch-like (e.g., Carmichaelia)
Clavate	Club-shaped, gradually widening towards apex.
Cleft	Having indentations that extend about halfway to the center, as in certain leaves.
Cleistogamous	Flowers that self-fertilise without opening.
Coherent	Sticking together of like parts.
Column	Stamen and stigmas fused to form a single organ.

Term	Definition
Columnar	Shaped like a column
Composite	many small flowers tightly packed together e.g., daisy flowers.
Compound	Composed of several similar parts (cf simple)
Concave	Curved inward.
Concolorous	Of the same colour.
Conical	Cone-shaped.
Connate	Fusion of like parts.
Conspecific	Individuals of the same species.
Cordate	Heart-shaped with the notch at the base.
Coriaceous	Leather-like; thick, tough, and somewhat rigid.
Corolla	The whorl of petals of a flower.
Corymb	Modified raceme where stalks of lower flowers are elongated to same level as the upper flowers.
Cosmopolitan	A species or other taxonomic group that is distributed widely throughout the world.
Costa	The midrib
Crenate	With rounded teeth (bluntly toothed) along the margin.
Crisped	Margin tightly wavy or crinkled, curled or wavy.
Cristate	With a crest.
Crown	The growing point of an upright rhizome or trunk. This usually produces a tuft or ring of fronds.
Crura	The two small projections at the mouth of a utricle in Carex
Cucullate	Hood-shaped.
Culm	The erect stem of a grass.
Cuneate	Wedge-shaped.
Cupular	Cup-shaped.
Cuttings	Stems and/or leaves taken from plants for propagation
Cyathium	A cup-like structure that surrounds the inflorescence in Euphorbia
Cyme	Inflorescence at the terminus of a branch and where new flowering branches emerge laterally below the flower.
Cytorace	Populations (or infraspecific taxa) that differ in chromosome number or chromosome morphology, e.g., <i>Nematoceras trilobum</i> agg. has two cytoraces, a diploid and a tetraploid (in which the chromosomes are doubled).
Cytotype	Populations (or infraspecific taxa) that differ in chromosome number or chromosome morphology, e.g., <i>Nematoceras trilobum</i> agg. has two cytotypes, a diploid and a tetraploid (in which the chromosomes are doubled).
Deciduous	Marked leaflessness in winter, and greater than 90% leaves lost by beginning of spring flush.
Decrescent	Diminishing.
Decumbent	With a prostrate or curved base and an erect or ascending tip.
Decurrent	Attached by a broadened base.
Decurved	Curved downward.
Deflexed	Bent abruptly downward.
Dehiscence	The time of opening at maturity to release the contents, e.g., a capsule releasing the seeds.
Dehiscent	Splitting open at maturity to release contents (of a fruit).
Deltoid	Shaped broadly like an equilateral triangle.
Dentate	Toothed along the margin with the teeth pointing outward, not forward.
Denticles	minute teeth
Denticulate	having a very finely toothed margin
Dichotomous	Divided into two equal branches.
Digitiform	Finger-like.
Dioecious	Having male and female flowers on separate plants of the same species.
Diploid	With two complete sets of chromosomes in each cell.
Disarticulating	Separating at a joint.
Discoid	Disc-shaped.
Disjunct	A species or other taxonomic group that occupies areas that are widely separated and scattered and therefore have a discontinuous distribution.
Distal	Toward the apex, away from the point of attachment (cf. proximal).
Distichous	In two rows on opposite sides of the axis.
Divaricating	Branching at a very wide angle with stiff intertwined stems.
Domatia	small structures on the lower surface of a leaf in some woody dicotyledons, located in the axils of the primary veins and usually consisting of depressions partly enclosed by leaf tissue or hairs.

Term	Definition
Dorsal	Of the back or outer surface relative to the axis. (cf. ventral)
Drupe	A stone fruit, the seed enclosed in a bony covering (endocarp) which is surrounded by a + fleshy layer (mesocarp)
Early successional species	Plants which are able to colonise an open area after disturbance but which are often temporary and are replaced by taller plants in time and shaded out.
Echinate	having sharply pointed spines or bristles.
Ecological district	A characteristic landscape and biological community defined in the PNA (Protected Natural Area) programme.
Ecological restoration	Attempt to reinstate original (pre-disturbance) state of a habitat, plant community or ecosystem.
Ecosourced	Plants sourced from seed collected from similar naturally growing plants in the area of the planting site.
Ecosourcing	Using native plants grown from locally grown seeds. Eco-sourced plants help to preserve the ecological distinctiveness of an area, and ecosourced plants fare better and are adapted to survive in the local conditions.
Eglandular	Without glands.
Elaiosome	Fleshy, oil-rich structure attached to seed that attracts ants which act as dispersers.
Ellipsoid	Elliptic in long section and circular in cross-section.
Elliptic	Broadest at the middle
Emarginate	With a notch at the apex.
Emarginated	Having a shallow notch at the tip, as in some petals and leaves.
Emergent	In an aquatic sense - wetland herbs that are rooted in the substrate below water level, but carry leaves and stems above the water level e.g. rushes and raupo. Found on the shallow margins of lakes, ponds and waterways. In a forest sense - tree that is appearing above the surrounding canopy.
Emergent marginals	An aquatic plant having most of its structure above water. Other aquatic plants are submerged or floating.
Endemic	Unique or confined to a place or region, found naturally nowhere else.
Endophyte	An endosymbiont (usually a bacterium or fungus) that lives within a plant for at least part of its life without causing any apparent disease.
Endophytes	Endosymbionts (usually bacteria or fungi) that live within plants for at least part of their lives without causing any apparent disease.
Endosperm	The nutritive tissue of a seed, consisting of carbohydrates, proteins, and lipids.
Enrichment planting	Returning to a revegetation site and creating gaps, or filling existing gaps, with different plants of plants, usually later successional plants which may not have survived being planted in the first phases of the project.
Ensiform	Sword shaped
Entire	Smooth. Without teeth, notches or divisions.
Entomophilous	Pollinated by insects.
Epicalyx	Calyx-like structure outside, but close to, the true calyx.
Epigeal	Growing on or close to the ground or emerging from the ground after germination (often used for cotyledons).
Epiphyte	A plant that grows upon another plant but is not parasitic and does not draw nourishment from it.
Epiphytic	Growing upon another plant but not parasitic and not drawing nourishment it
Erose	Irregularly toothed, as if gnawed.
Estuarine	Pertaining to the meeting of freshwater and seawater wetlands.
Ethnobotany	The study of people's classification, management and use of plants.
Eusporangia	Sporangia that arise from groups of epidermal cells
Evanescent	Lasting a very short time or running a short distance.
Ex situ	Away from the place of natural occurrence.
Ex-situ	Maintenance of plants as live specimens or propagules in cultivation as insurance against the loss of wild populations and as source for material for translocation.
Excurrent	Having the axis prolonged to form an undivided main stem or trunk (as in conifers).
Extravaginal	Outside an enclosing sheath
Falcate	Hooked or curved like a sickle.
Fastigate	Branches erect and close to central axis.
Fen	A type of wet land that accumulates peat deposits. Fens are less acidic than bogs, deriving most of their water from groundwater rich in calcium and magnesium.
Ferruginous	Rust-like (a colour term)
Fertile frond	Fronds that bear sporangia.
Filamentous	Resembling a filament.
Filiform	Thread like, resembling a filament.
Filiramulate	Branching at a very wide angle with stiff intertwined stems.
Fimbriae	Plural of fimbria: Fringe. A fimbria is composed of many fimbriellae (individual hair-like structures).
fimbriate	With fringes.
Flabellate	Fan shaped.
Flaccid	Limp, not rigid, flabby.
Flange	A projecting rim.

Term	Definition
Flexuose	With curves or bends.
Floccose	Having tufts of soft woolly hairs
Floret	A small flower, usually one of a cluster - the head of a daisy for example.
Foliaceous	Leaf-like.
Foliate	Having leaflets.
Founder effect	When a small number of plants (and therefore their genes) from a larger population are selected some genetic information is lost.
Fronnd	A leaf, the complete leaf of a fern including the stipe and lamina
Fulvous	Orange–yellow.
Funneliform	Funnel-shaped.
Fusiform	Broadest near the middle and tapering toward both ends.
Galea	Helmet- or hood-shaped.
Galeate	Shaped like a helmet or hood.
Gametophyte	A plant that produces sperm and egg cells and in which sexual reproduction takes place - in ferns this is known as the prothallus
Gene pool	The mixture of all genes and gene variations of a group or population.
Genetic diversity	The variety of genes in a plants or populations.
Genetic variation	Differences displayed by individuals within a plant which may be favoured or eliminated by selection.
geniculate	abruptly bent
Genus	A taxonomic rank of closely related forms that is further subdivided in to species (plural = genera). In a scientific name (e.g., <i>Sicyos australis</i>), the first word is the genus, the second the species.
Gibbous	Swollen or enlarged on one side, as in a gibbous moon.
Glabrescent	Lacking hair or a similar growth or tending to become hairless
Glabrous	Without or devoid of hairs, smooth.
Gland	A structure that secretes a sticky or oily substance.
Glandular	A structure that secretes a sticky or oily substance.
Glaucous	Covered with a fine, waxy, removable powder that imparts a white or bluish cast to the surface.
Gley	A soil prone to seasonal inundation.
Globose	Globe-shaped.
Glume	One of two bracts at the base of a grass spikelet.
Groundwater	Groundwater is the water beneath the surface that can be collected with wells, tunnels, or drainage galleries, or that flows naturally to the earth's surface via seeps or springs. Groundwater is the water that is pumped by wells and flows out through springs.
Gymnosperm	Plants in the class Gymnospermae that have seeds which are not enclosed in an ovary.
Gynodioecious	A species population containing plants that produce bisexual (perfect) flowers, and plants that produce only female (pistillate) flowers.
Gynoeceium	The female reproductive organs of a flower; the pistil or pistils considered as a group. Means literally "womans house" i.e., the overall structure that contains the female sex organs
Hastate	Spear like. Shaped like an arrowhead, but with basal lobes pointing outward rather than downward.
Haustorium	The absorbing organ of a parasite or hemiparasite
Hemi–parasite	Obtains water and nutrients from the roots of other plants but also manufactures food through photosynthesis.
Hemi–parasitic	Obtaining water and nutrients from the roots of other plants then manufacturing food through photosynthesis.
Herbarium	The place where collections of dried/pressed plants are kept.
Hermaphrodite	Having both male and female sexual characteristics and organs.
Heteroblastic	Exhibiting differences in leaf shapes or forms in juvenile and adult phases of the plant.
Heteroblasty	The state of being heteroblastic (i.e., exhibiting differences in leaf shapes or forms in juvenile and adult phases of the plant).
Hirsute	Hairy.
Hyaline	Membranous, thin and translucent.
Hybrid	An individual that is the offspring of a cross between two different varieties or species.
Hybridise	Breeding with a member of a different plant or type.
Hydrophyte	A plant species adapted to growing in or on water or in wet situations. Aquatic or semi-aquatic.
Hymenium	The fertile, spore–bearing layer of a fruitbody.
Hypanthium	A ring–like, cup–shaped, or tubular structure of a flower on which the sepals, petals, and stamens are borne.
Imbricate	Overlapping.
imbricating	Overlapping.
Imparipinnate	Odd–pinnate, a leaf shape; pinnate with a single leaflet at the apex.
In-situ	On site conservation relating to the maintenance of plants in the wild.
Inbreeding	Genetic similarity in offspring of closely related individuals.

Term	Definition
Incoherent	Not sticking together.
Incursion	Entrance of a pest into an area where it is not present
Indumentum	A covering of fine hairs (or sometimes scales)
Indusia	Plural of indusium, a membrane covering a sorus of a fern
Indusium	A thin tissue that covers the sorus in many ferns. Plural: indusia.
Inflorescence	The arrangement of flowers on the stem. A flower head.
Infundibuliform	Funnel-like.
Interkeel	The space between the keel and the leaf blade
Internode	The part of an axis between two nodes; the section of the stem between leaves.
Internodes	Part of a stem between two nodes.
Intramarginal	Within or near the margin.
Involucral bracts	The scales surrounding the flower head or capitula.
Involucre	A group of bracts surrounding a flower head.
Involute	With margins rolled inward toward the upper side.
Irritable	Responding to touch.
Jugate	Paired.
Juvenile	A plant of non-reproducing size.
Keel	A prominent or obvious longitudinal ridge (as in a boat).
Labellar	Pertaining to the labellum: a lip; in orchid flowers referring to the middle petal which usually differs in size, shape or ornamentation from the two lateral petals.
Labellum	A lip; in orchid flowers referring to the highly modified middle petal which usually differs in size, shape or ornamentation from the two lateral petals.
Lacinia	A jagged lobe.
Laciniae	Jagged lobes.
Laciniate	Cut into narrow, irregular lobes or segments.
Lacustrine	Of or having to do with a lake, of, relating to, or formed in lakes, growing or living in lakes.
Lamina	The expanded flattened portion or blade of a leaf, fern frond or petal.
Lanceolate	Lance-shaped; of a leaf several times longer than wide with greatest width about one third from the base, tapering gradually to apex and more rapidly to base
Lateral	On or at the side.
Lax	With parts open and spreading, not compact.
Laxly	With parts open and spreading, not compact
Leaflet	One section of a compound leaf.
Lemma	The lower of two bracts enclosing the flower in grasses.
Lenticillate	Bark that is covered in fine lenticles (breathing pores)
Ligulate	Strap-like, tongue-shaped
Ligule	The membrane between the leaf and the stem of a grass; the "petal" of a ray floret in a composite inflorescence
Linear	Long and narrow with more or less parallel sides.
Littoral	Occurring at the border of land and sea (or lake). On or pertaining to the shore. The shallow sunlit waters near the shore to the depth at which rooted plants stop growing.
Lobe	A recognisable, but not separated, rounded division or segment of a leaf or pinna. Used to describe ferns and leaves in <i>Cotula</i> and <i>Leptinella</i> .
Lobed	Part of a leaf (or other organ), often rounded, formed by incisions to about halfway to the midrib.
Lobule	A small lobe or sub-division of a lobe
Lustrous	Glossy, shiny.
Lycophytes	Seedless vascular plants that belong to the phylum Lycophyta (characterised by microphylls -primitive leaves found in ancient plants).
Lyrate	Pinnatifid or pinnatisect terminal lobe much larger than lower lobes.
Maculate	Blotched or spotted.
Mangrove	Coastal wetland dominated by Manawa or mangrove <i>Avicennia marina</i> var. <i>resifera</i> . Northern New Zealand only, salt marsh replaces it further south.
Margin	The edge or border of a leaf
Marine	Pertaining to the sea and saltwater systems.
Marsh	A tract of wet land principally inhabited by partially-submerged herbaceous vegetation. Has fewer woody plants than swamplier habitats.
Mealy	Dry, powdery, crumbly.
Median	In the middle.
Membranous	Very thin, like a membrane.
Mid-lobe	The middle part into which a leaf is divided.
Midrib	The central or principal vein of a leaf or pinna of a fern.
Mire	Synonymous with any peat-accumulating wetland. Term covers bogs and peaty swamps, fens, carr, moor, muskeg and peatland. Term excludes marsh which is non-peat forming.

Term	Definition
Molecular techniques	Where proteins and genes are used to investigate plant relationships
Monitoring	Recording of quantitative data over time to document changes in condition or state of species or ecosystems.
Monoecious	Having male and female flowers on the same plant of the same species.
Montane	Land between 300 and 800 metres above sea level.
Mucronate	Tipped with a short, sharp, point.
Mucronulate	Having a very small mucro; diminutive of mucronate.
Multi-annual evergreen	Overlapping annual cohorts of leaves always present.
Multifid	Cleft into many lobes or segments
Multiseptate	With many septa.
muricate	Rough with short, hard points like the shell of Murex, a genus of tropical sea snails with elaborately pointed shells.
Mycorrhiza	A symbiotic relationship between a fungus and a plant.
Mycorrhizal associations	Symbiotic association between fungi and plant roots which assists plant health by allowing increased ability for uptake of nutrients and promote plant growth.
Napiform	A long swollen but tapering root – like a parsnip, or carrot.
Native	Naturally occurring in New Zealand (i.e., not introduced accidentally or deliberately by humans).
naturalised	Referring to plants that have escaped from cultivation (including gardens or forest plantations) and can now reproduce in the wild (without human assistance)
Nectary	Organ that produces nectar.
Nerve	Prominent vein or rib.
Nerves	Strands of conducting and usually strengthening tissue in a leaves or similar structures
Net veins	Veins that repeatedly divide and re-unite.
Net venation	Feather-like or hand-like venation on a leaf.
Nival	Growing at high altitudes. From Latin: nivalis, snowy etc. from nix, nivis, snow.
Node	The point at which leaves, branches or roots arise on a stem.
Ob-	Prefix meaning inverted, in reverse direction.
Obcordate	Heart shaped with the notch at the apex.
Oblanceolate	Tapering and widest towards the apex or inversely lanceolate.
Oblique	Slanting; of a leaf, larger on one side of the midrib than the other, in other words asymmetrical.
Oblong	Rectangular.
Obovate	Roughly elliptical or reverse egg shaped and widest near the apex (i.e., the terminal half broader than the basal half).
Obtuse	Blunt or rounded at the apex, with the sides meeting at an angle greater than 90°.
Operculate	With a small lid.
Opposite	A pair of organs attached at nodes in pairs on either side of a stem or axis.
Orbicular	Almost or approximately circular.
Outbreeding depression	A reduction in vigor of offspring from distant parents. It can occur when a locally adapted population is moved and mixed with plants adapted to different conditions.
Outer canopy deciduous	Marked reduction in leaf number in the outer canopy in exposed high light environments over winter.
Oval	Planar, shaped like a flattened circle, symmetrical about both the long and the short axis; about twice as long as broad, tapering equally both to the tip and the base. Synonymous with elliptical.
Ovary	Part of a flower containing the ovules and later the seeds.
Ovate	Egg-shaped and widest at base.
Ovoid	Oval; egg-shaped, with rounded base and apex.
Pakihi	A term which in its strict sense refers to open clears within forest dominated by low scrub and rushes. However, more usually used to refer natural and induced wetlands and their associated shrublands. A vernacular most frequently used in the West Coast for impoverished soils and their associated peats, left after forest has been cleared
Palea	The small upper bract enclosing the flower of a grass
palea	1. The upper of the two bracts that enclose each floret in a grass spikelet. 2. A small bract at the base of a disc floret in some plants of the composite family. 3. Scales on various parts of ferns (referred to as paleate or paleaceous). From the Latin word for 'chaff'.
paleae	Plural of palea, from the Latin word for 'chaff'. 1. The upper of the two bracts that enclose each floret in a grass spikelet. 2. A small bract at the base of a disc floret in some plants of the composite family. 3. Scales on various parts of ferns (referred to as paleate or paleaceous).
Palmately	Radiating from a point, as fingers radiating from the palm of a hand.
Palmatifid	Deeply divided into several lobes arising from more or less the same level.
Palmatisect	Intermediate between palmate and palmatifid, i.e. the segments are not fully separated at the base; often more or less digitate.
Palustrine	Pertaining to wet or marshy habitats. Term covers mires and marshes
Pandurate	Fiddle-shaped.
Panicle	Highly branched (multiple raceme).

Term	Definition
Papilla	A short rounded projection.
Papillae	A soft, fleshy projection, usually small and nipple-like.
Papillate	With short rounded projections.
Papillose	Warty, with short rounded projections or gland-dotted
Parallel venation	Veins are parallel along leaf.
Parasite	An organism that derives all its nourishment from its host.
Patent	Spreading or expanded, e.g., spreading petals.
Peat	A mass of partially carbonised plant tissue formed by partial decomposition in water of various plants and especially of mosses of the genus Sphagnum, widely found in many parts of the world, varying in consistency from a turf to a slime used as a fertiliser, as stable litter, as a fuel, and for making charcoal. Partially carbonized vegetable matter saturated with water; can be used as a fuel when dried. A type of soil deriving from dead organic material situated in a wet area, where the reduced amount of [[oxygen available in the wet conditions results in the organic material not decomposing as much as it usually would do so in the presence of more oxygen. Used in growing media. Represents an important carbon sink –drainage of peat releases large amounts of carbon (CO ₂) to the atmosphere.
Pedicel	The stalk of a single flower in an inflorescence or fruit (either in a cluster or existing singularly).
Peduncle	The stalk of a solitary flower or the main stalk of an inflorescence or flower cluster.
Pedunculate	Describing fruits, which are borne on a stalk (a peduncle).
Pellucid	Transparent.
Peltate	Shield-like, with the stalk attached well inside the margin
Pendent	Hanging down from its support
Pendulous	Hanging or drooping.
Penicillate	With a tuft of hairs at the end, like a brush.
Perennial	A plant lasting for three seasons or more
Perianth	A collective term for the calyx (sepals or tepals) and corolla (petals) of the flower, especially when these are indistinguishable
Petal	Part of flower inside the sepals; usually coloured.
Petiolate	Having a petiole.
Petiole	Leaf stalk.
phloem	The vascular tissue in land plants that is primarily responsible for the distribution of sugars and nutrients manufactured in a shoot.
Photopoint	A monitoring technique where repeat photos are taken of the same scene from the same point over a period of time in order to quantify changes.
Pilose	Bearing long, soft hairs.
Pinna	A segment of a divided lamina that is classified as primary, secondary or tertiary according to the degree of dissection of the lamina.
Pinnae	Divisions of a pinnate leaf
Pinnate	With leaflets arranged regularly in two rows on either side of a stalk as in a feather; the lamina on a fern is divided into separate pinnae
Pinnatifid	Pinnately lobed, cleft more than halfway to the midrib. Not cleft all the way to the rachis.
Pinnatisect	Pinnately divided almost to midrib but segments still confluent.
Pioneer	Plant species are hardy species that should be planted first to establish a good canopy cover that restricts weed growth and promotes natural regeneration. In natural ecosystems these are the first plants to arrive and grow on a site.
Pistil	The female reproductive organ of a flower, consisting of an ovary, style, and stigma.
Pistillate	A flower with one or more pistils, but no stamens.
Plano-convex	Flat on one side, convex on the other.
Plumose	Feathery.
Podzol	Infertile, acidic soil, strongly leached to form a whitish-grey subsoil underlain by a layer enriched in iron, aluminium and organic matter; usually under forest in a wet temperate climate.
Pole	A subcanopy size individual with a long thin trunk and foliage tuft of a potential canopy tree.
Pollinia	Compact masses of orchid pollen.
Population enhancement	Increasing a population for a specific biological purpose, e.g., when a species is already present in an area but extra individuals are added to address a sex imbalance.
Porrect	Extending forward.
Procumbent	Lying and flat along the ground but not rooting
Propagate	To reproduce a plant by sexual (i.e., from seed) or asexual (e.g., from cuttings) means.
Prostrate	A general term for lying flat along the ground. This includes procumbent (that is lying and flat along the ground but not rooting) and decumbent (with a prostrate or curved base and an erect or ascending tip).
Provenance	The place of origin (of a plant that is in cultivation).
Proximal	Toward the base or point of attachment (cf. distal).
Pseudobulb	Thickened surface stem; usually looking like a bulb.
Pseudoterminal	Falsely terminal – as in a bud which appears to occupy a terminal position but does not

Term	Definition
Puberulent	Minutely clad in short, soft hairs
Pubescence	Covering of soft, fine hairs
Pubescent	Covered in short, soft hairs.
Pungent	Ending in a stiff sharp point
Pustule	Small blister-like elevation.
Quadrate	Square, rectangular.
Raceme	An unbranched, elongated inflorescence with pedicellate flowers maturing from the bottom upward i.e., flowers attached to the main stem by short stalks.
Rachis	the axis of an inflorescence or of a compound leaf
Ray	An outer ring of strap-like florets in the head of Asteraceae (daisy) flowers.
Re-introduction	Translocating wild or cultivated individuals to sites where the taxon has been known to occur in the past, but from which it has disappeared.
Recurved	Curved backward.
Reflexed	Bent back on itself
Reniform	Kidney shaped.
Repend	With a slightly wavy margin.
Replum	The outer structure of a pod in which the valves have dehisced (persists after the opening of the fruit)
Restiad	Area dominated by rush-like plants (collectively known as restiads) of the family Restionaceae. Includes Chatham Island and North Island Sporodanthus and oiioi (<i>Apodasmia similis</i>)
Retorse	Pointing backward.
Retuse	A shallow notch at the rounded or blunt apex of a leaf.
Rhizoid	Any of various slender filaments that function as roots in mosses and ferns and fungi.
Rhizomatous	With underground creeping stems.
Rhizome	An underground stem (usually spreading horizontally or creeping) or short and erect.
Rhombic	Diamond-shaped.
Rhomboid	Diamond shaped, nearly rhombic.
Riparian	Relating to or living or located on the bank of a natural watercourse (as a river) or sometimes of a lake or a tidewater.
Riparian margin	Refers to the edges of streams, rivers, lakes or other waterways.
Riparian plants	Refers to plants found growing near the edges of streams, rivers or other waterways.
Riparian zone	A strip of land next to streams, rivers, and lakes where there is a transition from terrestrial (land vegetation) to aquatic (water) vegetation. Also known as "berm".
Riverine	Pertaining to rivers, streams and such like flowing water systems.
Rootstock	A short, erect, underground stem.
Rosette	A radiating cluster of leaves.
Rostellum	In orchids, a modified stigma that prevents self-fertilisation.
Rosulate	A dense radiating cluster of leaves.
Rugose	Wrinkled.
Rugulose	Having small wrinkles.
Runcinate	Sharply pinnatifid or cleft, the segments directed downward.
Runner	A trailing stem that roots at the nodes.
Rupestral	Growing on rocks.
Rushes	A group of distinctive wetland plants. They have solid stems (grasses have hollow stems), true rushes <i>Juncus</i> sp. have rounded leaves.
Sagittate	Shaped like the head of an arrow; narrow and pointed but gradually enlarged at base into two straight lobes directed downwards; may refer only to the base of a leaf with such lobes; cf. hastate.
Salt marsh	A coastal wetland, with specialized salt tolerant plants (halophytes).
Sapling	A juvenile tree that has reached the stage of 1 or 2 main stems but is still in the shrub layer.
Saprophyte	A plant lacking chlorophyll and living on dead organic matter.
Saprophytic	Lacking chlorophyll and living on dead organic matter.
Sarcotesta	The fleshy, often highly coloured outer layer of the seed coat in some species, e.g., titoki (<i>Alectryon excelsus</i>).
Scabrid	Roughened or rough with delicate and irregular projections.
Scale	Any thin, flat, membranous structure.
Scape	A leafless flower stem.
schizocarp	A fruit which splits when dry, from the Greek <i>skhizein</i> 'split' and <i>karpos</i> 'fruit'
schizocarps	Plural of schizocarp, a fruit which splits when dry, from the Greek <i>skhizein</i> 'split' and <i>karpos</i> 'fruit'
Scutiform	Shield-shaped.
Sedges	A group of grass-like or rush-like herbaceous plants belonging to the family Cyperaceae. Many species are found in wetlands some are forest floor plants. Leaves are usually angular. Hence the saying "rushes are round and sedges have edges".

Term	Definition
Seedling	A newly germinated plant.
Self sustaining	Able to sustain itself, or replace itself, independently of management i.e. regenerate naturally
Self thinning	Natural tree death in a crowded, even-aged forest or shrubland.
Semi-deciduous	Partial leaflessness in winter, and greater than 50% leaves lost by the beginning of spring flush.
Sepal	Outer part of flower; usually green.
Serrate	Sharply toothed with teeth pointing forwards towards apex.
Serrulate	Finely serrate, i.e., finely toothed with asymmetrical teeth pointing forward; like the cutting edge of a saw.
Sessile	Attached by the base without a stalk or stem.
Seta	The stalk of a fruiting moss capsule
Sheath	A portion of an organ that surrounds (at least partly) another organ (e.g., the tubular envelope enclosing the stem in grasses and sedges).
Silicles	The flattened usually circular capsule – compared with the narrow, elongated fruit (silique) – containing the seed/seeds. A term used almost exclusively for plants within the cabbage family (Brassicaceae)
Silique	A capsule, usually 2-celled, with 2 valves falling away from a frame (replum) bearing
Simple	Of one part; undivided (cf compound).
Sinuate	With a wavy margin.
Sinus	The space or recess between lobes; in hebes a gap between the margins of two leaves of an opposite pair that may be present in the bud before the pair of leaves separate.
Sorus	A cluster of two or more sporangia on the margin or underside of the lamina of a fern, sometimes protected by an indusium.
Spathulate	Spatula or spoon-shaped, a rounded blade tapering gradually to the base.
Spheroidal	Almost spherical but elliptic in cross section.
Spicate	Arranged in a spike.
Spike	Flowers attached to main stem without stalks.
Spikelet	Collection of individual grass florets borne at the end of the smallest branch of the inflorescence.
Sporangia	Plural of sporangium. Structures in which spores are produced.
Sporangium	Structure in which spores are produced.
Spore	A single-celled reproductive unit similar in function to that of the seed in a flowering plant.
sporophyte	The spore producing plant in ferns that is usually the visible part.
Stamen	The male reproductive organ of a flower where pollen is produced. Consists of an anther and its stalk.
Stamens	The male, pollen bearing organ of a flower.
Standing water	Where water lies above the soil surface for much of the year.
Stellate	Irregularly branched or star shaped.
Stigma	Female part of the flower that is receptive to pollen, usually found at or near the tip (apical end) of the style where deposited pollen enters the pistil.
Stipe	The stalk of a frond.
Stipitate	Borne on a stipe or stalk.
Stipulate	A leaf with stipules.
Stipule	A scale-like or leaf-like appendage at the base of a petiole, usually paired.
Stolon	A stem which creeps along the ground, or even underground.
Stoloniferous	Producing stolons
Stramineous	Chaffy, like straw or straw-colored.
Stria	A fine line or groove.
Striae	Fine lines or grooves.
Striate	Fine longitudinal lines or minute ridges
Style	The elongated part of the flower between the ovary and the stigma.
Sub-	A prefix meaning under, somewhat or almost.
Subglabrous	Very slightly, but persistently, hairy.
Suborbicular	Slightly rounded in outline
Substrate	The surface upon which an orchid grows.
Subtended	Immediately beneath, occupying a position immediately beneath a structure, i.e., flower subtended by bract
Subulate	Slender and tapering to a point.
Succession	Progressive replacement of one species or plant community type by another in an ecosystem.
Successional	Referring to species, plant communities or habitats that tend to be progressively replaced by another.
Succulent	Fleshy and juicy.
Summer-green	Used in New Zealand to indicate herbs or sub-shrubs that die down to a root stock or rhizomatous network.
Supplementary planting	Returning to a revegetation site and creating gaps, or filling existing gaps, with different plants of plants, usually later successional plants which may not have survived being planted in the first phases of the project.

Term	Definition
Surface water	Water present above the substrate or soil surface.
Surveillance	Regular survey for pests inside operational and managed areas e.g. nurseries, standout areas on parks.
Survey	Collection of observations on the spatial distribution or presence or absence of species using standardised procedures.
Sustainable Land Management	The use of farming practices which are sustainable both financially and environmentally including management of nutrient runoff, waste disposal or stock effluent, reducing impacts of nutrients on waterways, preventing erosion and soil loss, and protecting native forest and wetland habitats from stock damage.
Swamp	Low land that is seasonally flooded; has more woody plants than a marsh and better drainage than a bog. They are more fertile and less acidic than bogs because inflowing water brings silt, clay and organic matter. Typical swamp plants include raupo, purei and harakeke (flax). Zonation and succession often leads through manuka to kahikatea swamp forest as soil builds up and drainage improves.
Symbiote	An organism that has an association with organisms of another species whereby the metabolic dependence of the two associates is mutual.
Symbiotic	The relation between two different species of organisms that are interdependent; each gains benefits from the other (see also symbiosis).
Sympatric	Occupying the same geographical region.
Synangia	Structures made up of fused sporangia
Synonym	A botanical name that also applies to the same taxon.
Systematics	The study of taxonomy, phylogenetics, and taxagenetics.
Tabular	Shaped like a rectangular tablet.
Taxa	Taxonomic groups. Used to refer to a group at any level e.g., genus, species or subspecies.
Taxon	A taxonomic group. Used to refer to a group at any level e.g., genus, species or subspecies.
Taxonomy	The process or science of classifying, naming, and describing organisms
Tepal	An individual member of the perianth.
Terete	Cylindrical and tapering.
Terminal	At the tip or apex.
Ternatifid	Leaflets In threes,
Tetrad	A group of four.
Tomentum	A hairy covering of short closely matted hairs.
Translocation	The movement of living organisms from one area to another.
Trifid	Divided into three.
Trifoliate	Having three leaflets.
Trigonus	Three-angled
Tripinnate	With each secondary pinna divided to the midrib into tertiary pinnae
Triquetrous	Triangular in cross section and acutely angled.
Truncate	With the apex or base squared at the end as if cut off.
Tuberculate	Bearing small swellings.
Tubular	Tube-shaped.
turbinate	Top-shaped.
Turgid	Distended through internal pressure
Type locality	The place or source where a holotype or type specimen was found for a species.
Ultramafic	A type of dark, usually igneous, rock that is chemically dominated by magnesium and iron-rich minerals, the partially metamorphosed form of which is serpentinite.
Umbel	Umbrella like; the flower stalks arise from one point at the stem.
Undulate	Wavy edged.
Undulose	Wavy edged.
Unitubular	A tube partitioned once – literally one tube (compare – multitubular – many tubes)
Utricle	A thin loose cover enveloping some fruits (eg., Carex, Uncinia)
Valvate	Opening by valves.
Vascular plant	A plant that possesses specialised conducting tissue (xylem and phloem). This includes flowering plants, conifers and ferns but excludes mosses, algae, lichens and liverworts.
Velutinous	Thickly covered with delicate hairs; velvety.
Ventral	Of the front or inner (adaxial) surface relative to the axis. (cf. dorsal)
Vermiform	Worm-shaped.
Vernicose	Glossy, literally as if varnished, e.g., Hebe vernicosa has leaves that appear as if varnished
Verrucose	Having small rounded warts.
Verticillium	A fungus disease that will cause wilting and death.
Villous	Covered with long, soft, fine hairs.
Water table	The level at which water stays in a soil profile. The zone of saturation at the highest average depth during the wettest season.
Wetland	A site that regularly has areas of open water for part or all of the year, or has a water table within 10 cm of the surface for at least 3 months of the year. Wetland ecosystems support a range of plant and animal species adapted to an aquatic or semi-aquatic environment.

Term	Definition
-------------	-------------------

Whipcord	A shrub in which the leaves are reduced to scales that are close-set and pressed against the stem.
-----------------	--

Whorl	A ring of branches or leaves arising at the same level around the stem of a plant.
--------------	--