



# Small trees shrubs, and flax. Lake Hāwea Southern Foreshore Reserve. June 201



## Table of Contents

Introduction	1
<i>Coprosma crassifolia</i>	2
<i>Coprosma dumosa</i>	3
<i>Coprosma lucida</i>	4
<i>Coprosma propinqua</i> var. <i>propinqua</i>	5
<i>Coprosma robusta</i>	6
<i>Corokia cotoneaster</i>	7
<i>Discaria toumatou</i>	8
<i>Helichrysum lanceolatum</i>	9
<i>Kunzea serotina</i>	10
<i>Lophomyrtus obcordata</i>	11
<i>Olearia avicenniifolia</i>	12
<i>Olearia hectorii</i>	13
<i>Olearia lineata</i>	14
<i>Ozothamnus vauvilliersii</i>	15
<i>Phormium cookianum</i> subsp. <i>cookianum</i>	16
<i>Phormium tenax</i>	17
<i>Pittosporum eugenioides</i>	18
<i>Pittosporum tenuifolium</i>	19
<i>Pseudopanax crassifolius</i>	20
<i>Rubus schmidelioides</i> var. <i>schmidelioides</i>	21
Glossary	22

Made on the New Zealand Plant Conservation Network website – [www.nzpcn.org.nz](http://www.nzpcn.org.nz)

### Copyright

All images used in this book remain copyright of the named photographer. Any reproduction, retransmission, republication, or other use of all or part of this book is expressly prohibited, unless prior written permission has been granted by the New Zealand Plant Conservation Network ([info@nzpcn.org.nz](mailto:info@nzpcn.org.nz)). All other rights reserved.

## Introduction

This book was compiled from information stored on the website of the New Zealand Plant Conservation Network ([www.nzpcn.org.nz](http://www.nzpcn.org.nz)).

This website was established in 2003 as a repository for information about New Zealand's threatened vascular plants. Since then it has grown into a national database of information about all plants in the New Zealand botanic region including both native and naturalised vascular plants, threatened mosses, liverworts and fungi.

Funding to develop the website was provided by the New Zealand Government's Terrestrial and Freshwater Biodiversity Information System Programme (TFBIS).

The species information used on the website has come from a variety of sources. The indigenous vascular plant text was written largely by Dr Peter de Lange (former Network Vice President). Peter based the descriptions on a wide range of sources including the Flora of NZ Series (Allan 1961, Moore and Edgar 1970 and Webb et al 1987) as well as numerous other taxonomic treatments. For a full bibliography of information sources see the References at the end of this book.

Where no published treatment was available Peter used herbarium specimens and his own knowledge of the flora to prepare species pages. Various other contributors have provided text and additional information to many species pages including botanists such as Mike Thorsen, John Barkla, Cathy Jones, Simon Walls, Nick Singers and many others. The threatened fungi text was written by Eric Mackenzie and Peter Buchanan (Landcare Research).

More than 200 photographers have kindly provided images to illustrate the website and for use in this book especially John Smith-Dodsworth, Jeremy Rolfe, Peter de Lange, Wayne Bennett and Gillian Crowcroft.

## The New Zealand Botanic Region

The information on the Network website, from which this book was compiled, is for species that are indigenous to or naturalised within the New Zealand Botanic Region as defined by Allan (1961). The New Zealand botanic region encompasses the Kermadec, Manawatawhi/Three Kings, North, South, Stewart Island/Rakiura, Chatham, Antipodes, Bounties, Snares, Auckland Campbell island/Motu Ihupuku and Macquarie.

## About the Network

The Network has more than 800 members worldwide and is New Zealand's largest non-governmental organisation solely devoted to the protection and restoration of New Zealand's indigenous plant life.

The vision of the New Zealand Plant Conservation Network is that '*no indigenous species of plant will become extinct nor be placed at risk of extinction as a result of human action or indifference, and that the rich, diverse and unique plant life of New Zealand will be recognised, cherished and restored*'.

Since it was founded in 2003 the Network has undertaken a range of conservation initiatives in order to achieve its vision.

That work has included:

- Training people in plant conservation
- Publishing plant books, reports and posters
- Raising money for the David Given Threatened Plant Research Trust to pay for plant conservation research scholarships
- Advocacy to raise awareness of the importance of plant life in general and especially New Zealand's status as a Global Centre of Plant Diversity
- Lobbying central and regional government and business to protect indigenous plant life
- Educating people about plant life through the Network website
- Connecting people through the monthly newsletter, the Network conference and the annual general meeting

## What is a threatened plant?

The NZ Threatened Plant Committee was formed in 1991 and ever since then it has met at regular intervals to review the status of indigenous vascular plants. It is made up of a small group of botanists that between them have an extensive knowledge of the native plants of New Zealand. This group is chaired by Dr Peter de Lange of the New Zealand Department of Conservation.

This committee applies a set of criteria to each native plant to determine its conservation status. The resulting list of species classified as threatened is published in the NZ Journal of Botany (see for example de Lange et al. 2009). The main threat categories used are: Extinct, Critical, Endangered, Vulnerable, Declining. Other categories used are: Recovering, Relict, Naturally Uncommon, Coloniser, Vagrant and Data Deficient. For vascular plants the threat status used in this book is taken from the 2009 conservation assessment (see de Lange et al 2009).

More recently other committees have been established to review the status of non-vascular plants but their lists are yet to be published.

## *Coprosma crassifolia*

### **Current Threat Status (2012):**

Not Threatened

### **Threats:**

Not Threatened

### **References and further reading:**

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

### **For more information, visit:**

[http://nzpcn.org.nz/flora\\_details.asp?ID=1708](http://nzpcn.org.nz/flora_details.asp?ID=1708)



**Caption:** Windshorn plant.

Aramoana

**Photographer:** John Barkla



**Caption:** Bark. Hayward Point,  
Dunedin

**Photographer:** John Barkla

## *Coprosma dumosa*

### **Current Threat Status (2012):**

Not Threatened

### **Distribution:**

Endemic. North Island from Kaimai Range south; South Island (rare in Fiordland), Stewart Island.

### **Flowering:**

October - November (-  
January)

### **Fruiting:**

February-April (-  
October)

### **Threats:**


Not Threatened

### **References and further reading:**


Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

### **For more information, visit:**

[http://nzpcn.org.nz/flora\\_details.asp?ID=2322](http://nzpcn.org.nz/flora_details.asp?ID=2322)



**Caption:** *Coprosma dumosa*  
**Photographer:** Graeme Jane



**Caption:** *Coprosma dumosa*  
**Photographer:** Graham Jane

## *Coprosma lucida*

### **Common Name(s):**

karamu, shining karamu

### **Current Threat Status (2012):**

Not Threatened

### **Threats:**

Not Threatened

### **References and further reading:**

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

### **For more information, visit:**

[http://nzpcn.org.nz/flora\\_details.asp?ID=1719](http://nzpcn.org.nz/flora_details.asp?ID=1719)



**Caption:** Coprosma lucida

**Photographer:** Wayne Bennett



**Caption:** Leaf of Coprosma lucida

**Photographer:** Wayne Bennett

## *Coprosma propinqua* var. *propinqua*

**Common Name(s):**

mingimingi

**Current Threat Status (2012):**

Not Threatened

**References and further reading:**

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

**For more information, visit:**

[http://nzpcn.org.nz/flora\\_details.asp?ID=1728](http://nzpcn.org.nz/flora_details.asp?ID=1728)



**Caption:** Waikanae Estuary.  
**Photographer:** Jeremy Rolfe



**Caption:** *Coprosma propinqua*  
var. *propinqua*  
**Photographer:** Wayne Bennett

# *Coprosma robusta*

## Common Name(s):

karamu, glossy karamu

## Current Threat Status (2012):

Not Threatened

## Distribution:

Endemic. North and South Islands. Naturalised on the Chatham Islands within a small area between Waitangi and Owenga.

## Habitat:

Common throughout coastal, lowland and lower montane habitats within shrublands and open sites within forest.

## Features:

Shrub or small tree up to 6 m tall. Branches numerous, stout, erect to somewhat spreading. Petioles stout, 10-20 mm long. Stipules fused towards base, obtuse, glabrous with one of two prominent, black, glandular denticles. Leaves 70-120 x 30-40-50 mm, leathery, dark green above, paler green beneath, glabrous, elliptic, elliptic-oblong to broad-ovate, acute or obtuse, apex mucronate. Venation reticulated, conspicuous. Male flowers in axillary many-flowered glomerules, corolla conspicuous, lobes triangular, acute, stamens 4-5, prominent. Females in compound clusters on peduncles 10-15 mm. Calyx and corolla much reduced, stigmas prominent. Drupe dark orange (rarely yellow), 8-8 x 4-5 mm, oblong to narrow-ovoid.

## Flowering:

(July-) August-September (-November)

## Fruiting:

(March-) April-May (-July)

## Threats:

Not Threatened

## References and further reading:

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

## For more information, visit:

[http://nzpcn.org.nz/flora\\_details.asp?ID=1733](http://nzpcn.org.nz/flora_details.asp?ID=1733)



**Caption:** Fruit of *Coprosma robusta*

**Photographer:** Wayne Bennett



**Caption:** *Coprosma robusta* (Karamu)

**Photographer:** Wayne Bennett



## *Corokia cotoneaster*

### **Common Name(s):**

Korokio, wire-netting bush

### **Current Threat Status (2012):**

Not Threatened

### **Threats:**

Not Threatened

### **References and further reading:**

Moorfield, J. C. (2005). Te aka : Maori-English, English-Maori dictionary and index. Pearson Longman: Auckland, N.Z.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 11: 285-309

### **For more information, visit:**

[http://nzpcn.org.nz/flora\\_details.asp?ID=1756](http://nzpcn.org.nz/flora_details.asp?ID=1756)



**Caption:** Upper Oreti River, Southland

**Photographer:** Jesse Bythell



**Caption:** Fuit, Greenstone Valley

**Photographer:** John Barkla

## *Discaria toumatou*

### **Common Name(s):**

matagouri, wild Irishman

### **Current Threat Status (2012):**

Not Threatened

### **Distribution:**

Endemic. North and South Islands. In the North Island known from near Waiuku south to the southern Wairarapa and Wellington coastline. Very uncommon in the North Island. In the South Island mainly east of the main divide, appearing to avoid areas of high rainfall

### **Threats:**

Not Threatened for most of its range. However, very uncommon and under threat throughout the North Island, where it is now known from very few sites and viable populations.

### **References and further reading:**

Chrystall, L. 1976. Further record of matagouri in the North Island. *Wellington Botanical Society Bulletin*, 39: 47

Duguid, F. 1976. Matagouri at Herbertville. *Wellington Botanical Society Bulletin*, 39: 45

Elder, N.L. 1966. Matagouri in the North Island. *Wellington Botanical Society Bulletin*, 33: 5

Elder, N.L. 1967. Matagouri in the North Island - Part 2. *Wellington Botanical Society Bulletin*, 34: 19-20

Moorfield, J. C. (2005). *Te aka : Maori-English, English-Maori dictionary and index*. Pearson Longman: Auckland, N.Z.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

### **For more information, visit:**

[http://nzpcn.org.nz/flora\\_details.asp?ID=1795](http://nzpcn.org.nz/flora_details.asp?ID=1795)



**Caption:** Matukituki, Mt Aspiring  
**Photographer:** John Sawyer



**Caption:** Matukituki, Mt Aspiring  
**Photographer:** John Sawyer

## *Helichrysum lanceolatum*

### **Current Threat Status (2012):**

Not Threatened

### **References and further reading:**

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285-309

### **For more information, visit:**

[http://nzpcn.org.nz/flora\\_details.asp?ID=809](http://nzpcn.org.nz/flora_details.asp?ID=809)



**Caption:** Otago Peninsula  
**Photographer:** John Barkla



**Caption:** Cooks Cove, East Coast  
**Photographer:** John Sawyer

## *Kunzea serotina*

### Common Name(s):

makahikatoa

### Current Threat Status (2013):

Not Threatened

### Distribution:

Endemic. New Zealand. North and South Islands from Central Volcanic Plateau south through central North Island and east to the southern Wairarapa, thence easterly from Marlborough to Central Otago

### Habitat:

*Kunzea serotina*, in the North Island part of its range is mostly a montane to subalpine species, extending into lowland sites in forest flats and other places where temperature inversions occur. In the South Island it is more wide ranging but still most confined to mountain areas and intermontane basins.

### Features\*:

Erect, columnar to tightly pyramidal, fastigiata, densely branched shrubs or trees 3–20 × 2–4 m developing with time into less densely branched, open pyramidal crowns. Trunk usually single, very rarely 2–3 arising from base, 0.10–0.86 m d.b.h., with basal portion of trunk covered in numerous, curled, chartaceous bark remnants. Bark chartaceous, greyish-white to pinkish-white, heavily cracked, often crumpled, detaching as inrolled, curled, sinuous, irregular pieces, pieces often congregating in branch forks and forming deep drifts at trunk base. Branches numerous arising at or near trunk base; short and stout, obliquely ascending, fastigiata; branchlets numerous, very leafy; indumentum copious, sericeous; persistent, divergent 0.05–0.08 μm, apices curved or slightly curled. Leaves heterophyllous; seedling, subadult leaves and that of reversion shoots, spreading to patent; lamina 0.8–7.8 × 0.6–1.2 mm, red-green, pale green suffused with red, rarely bright green, linear-lanceolate to lanceolate; flat or involute, apex acute to obtuse, finely cuspidate; adult leaves, usually densely aggregated along brachyblasts, lamina 2.0–6.3 × 0.8–1.8 mm, dark glossy green or bronze-green, linear-oblong, oblanceolate to obovate; strongly recurved from about ½ of total length, apex initially acute to subacute, maturing obtuse to rounded, often cuspidate; base attenuate; glandular punctate, glabrous, very rarely with fine antrorse hairs near base; lamina margin sparsely hairy, usually in one interrupted row failing well short of leaf apex. Inflorescence a compact 1–12-flowered corymbiform botryum up to 25 mm long, borne on alternate, distinctly spiralled, densely leafy, brachyblasts up to 15 mm long. Pherophylls deciduous, mostly foliose, 0.9–2.5 mm long, green to bronze-green, spatulate, spatulate-orbicular, rarely pandurate or lanceolate, margins and apex finely ciliate. Flower buds clavate to pyriform, apex flat to weakly domed prior to bud burst, calyx valves not or scarcely meeting. Fresh flowers 2.8–8.8 mm diam. Hypanthium 1.6–3.4 × 1.5–3.8 mm, dark green or red-green, drying brown-green to red-brown; urceolate to campanulate terminating in a distinctly thicker rim bearing five persistent calyx lobes; copiously dotted with red oil glands, finely puberulent to ± glabrescent; hairs if present short, divergent. Calyx lobes 5, upright, 0.8–1.2 × 0.7–1.2 mm, persistent, ovate to broadly ovate, central portion of lobe pale green or yellow-green, with margins usually cream to pale pink, surface glandular punctate, oil glands usually pink in exposed situations otherwise ± colourless, glabrous except ciliate margins. Receptacle pink at anthesis, darkening to dark crimson magenta after fertilisation. Petals 5–6, 1.4–2.0 × 1.2–2.0 mm, white, sometimes basally flushed pink, narrowly orbicular to broadly ovate or cuneate, apex obtuse to rounded, margins ± frayed to finely and irregularly toothed, oil glands yellow when fresh, when dried very pale yellow to colourless. Stamens 20–38 in 1–2 weakly defined whorls, arising from receptacular rim, filaments white occasionally tinged rose-pink toward base. Anthers dorsifixed, 0.04–0.06 × 0.02–0.04 mm, testiculate to ellipsoid. Pollen white. Anther connective gland prominent, orange often flushed rose when fresh, drying dark orange-brown or purple, spheroidal, distinctly papillate. Ovary 3–5 locular, each with 10–23 ovules in two rows on each placental lobe. Style 0.6–1.2 mm long, white; stigma capitate, scarcely wider than style, usually flat to very weakly domed, greenish-white, cream or pale pink, surface finely papillate. Fruits rarely persistent 1.2–3.0 × 1.2–3.4 mm, light brown to grey, finely hairy, urceolate to shortly-campanulate, rarely cupular, splits concealed by dried, suberect to erect, free portion of hypanthium and incurved calyx lobes. Seeds 0.60–1.00 × 0.48–0.60 mm, narrowly oblong, oblong, oblong-obovate, orange-brown to dark brown, surface coarsely reticulate.

### Flowering:

November - May

### Fruiting:

January - December

### Threats:

Not Threatened

### \*Attribution:

Fact Sheet prepared for NZPCN by P.J. de Lange 25 August 2014. Description modified from de Lange (2014).

### References and further reading:

de Lange, P.J. 2014: A revision of the New Zealand *Kunzea ericoides* (Myrtaceae) complex. *Phytokeys* 40: 185p doi: 10.3897/phytokeys.40.7973.

### For more information, visit:

[http://nzpcn.org.nz/flora\\_details.asp?ID=7640](http://nzpcn.org.nz/flora_details.asp?ID=7640)



**Caption:** In cultivation ex Te Porere. Jul 2007.

**Photographer:** Jeremy Rolfe

# *Lophomyrtus obcordata*

## Common Name(s):

Rohutu, New Zealand myrtle

## Current Threat Status (2012):

Not Threatened

## Distribution:

Endemic. North and South Islands. Patchy and often absent over large parts of the country. More common in the eastern North and South island though locally prominent in some parts of western Northland and Auckland.

## Habitat:

Coastal to montane in forest - though mostly found in coastal and lowland forested habitats. *Lophomyrtus obcordata* is often rather local over large parts of its range though it seems to be reasonably common on the Pouto and Kaipara Peninsula where it grows on stable sand dunes in a forest dominated by an as yet unnamed species of *Kunzea ericoides* (known as rawirinui to northern Maori and referred to by de Lange & Murray 2004, de Lange et al., 2005 and de Lange et al. 2010 as *Kunzea* aff. *ericoides* (a)). *Lophomyrtus obcordata* is also occasionally dominant in alluvial forest remnants of the eastern South Island. In these places it is often parasitised by the dwarf mistletoe *Korthalsella lindsayi*.

## Features\*:

Shrub up to c.6 m tall. Trunk slender, up to 0.2 m diameter. Bark greyish pink, chartaceous, flaking in large, irregular shards, underbark pale cream. Branches numerous, erect, compactly branched to spreading. Branchlets initially 4-angled maturing subterete, rather brittle, minutely pubescent. Leaves opposite (sometimes in fascicles), coriaceous, puberulent when young (hairs patent), becoming glabrous with age, glandular punctate, oil glands colourless, leaf lamina and petioles shortly decurrent with branchlet; petiole 0.7-1.0 mm long, brittle; leaf lamina 5-12 × 5-10 mm, obcordate, cuneately narrowed to base, adaxially dark green to grey-green (sometimes tinged red), paler beneath (sometimes tinged pink). Flowers 4-merous, 6-8 mm diameter, borne in axillary, solitary monads on slender, 10-14(-20) mm long, pubescent pedicels. Hypanthium subturbinate, not extending beyond ovary summit, glandular punctate, oil glands colourless (rarely pink tinged), calyx lobes 4, 1.0-1.8 mm long, persistent, spreading, pubescent, oblong, acute. Petals 6-8 × 5-8 mm, suborbicular, white, margins entire to slightly irregular, oil glands colourless. Stamens 60-80(-100 or more), free, in 4 weakly defined whorls, filaments 6-8 mm long, anthers cream, dorsifixed, latrorse. Ovary inferior, 2-3-locular, ovules numerous, in a single row on each linear placenta. Style 6-8 mm long, slender, white, stigma capitate, scarcely dilated. Fruit a broadly ovate, bright to dark red (rarely black or violet) 6-7 mm long berry. Seeds numerous, reniform, 2.0-4.3 mm diameter, testa pale brown, glossy ± smooth, very hard.

## Flowering:

November - March

## Fruiting:

January - May

## Threats:

Not Threatened

## \*Attribution:

Fact sheet including description prepared for NZPCN by P.J. de Lange 9 February 2011. Seed description which is modified from Webb & Simpson (2001).

## References and further reading:

de Lange, P.J.; Murray, B.G. 2004: Chromosome numbers of New Zealand *Kunzea* (Myrtaceae). *Australian Journal of Botany* 52: 609-617.

de Lange, P.J.; Datson, P.M.; Murray, B.G.; Toelken, H.R. 2005: Hybridism in the *Kunzea ericoides* complex (Myrtaceae): an analysis of artificial crosses. *Australian Systematic Botany* 18: 117-131.

de Lange, P.J.; Smissen, R.D.; Wagstaff, S.J.; Keeling, D.J.; Murray, B.G.; Toelken, H.R. 2010: A molecular phylogeny and infrageneric classification for *Kunzea* (Myrtaceae) inferred from rDNA ITS and ETS sequences. *Australian Systematic Botany* 23: 309-319.

Webb, C.J.; Simpson, M.J.A. 2001: *Seeds of New Zealand Gymnosperms and Dicotyledons*. Christchurch, Manuka Press.

## For more information, visit:

[http://nzpcn.org.nz/flora\\_details.asp?ID=943](http://nzpcn.org.nz/flora_details.asp?ID=943)



**Caption:** Bark. Upper Hutt, eb 2011.

**Photographer:** Jeremy Rolfe



**Caption:** Stevensons Island, Lake Wanaka.

**Photographer:** John Barkla

## *Olearia avicenniifolia*

**Common Name(s):**

Mountain akeake

**Current Threat Status (2012):**

Not Threatened

**Threats:**

Not Threatened

**For more information, visit:**

[http://nzpcn.org.nz/flora\\_details.asp?ID=1044](http://nzpcn.org.nz/flora_details.asp?ID=1044)



**Caption:** Minaret Burn, March

**Photographer:** John Barkla



**Caption:** Olearia avicenniifolia,  
South Otago

**Photographer:** John Barkla

# *Olearia hectorii*

## Common Name(s):

Deciduous tree daisy, Hectors tree daisy

## Current Threat Status (2012):

Threatened - Nationally Endangered

## Distribution:

Endemic. Eastern South Island.

## Habitat:

Lowland to subalpine often at the base of steep hills on colluvium, or on alluvium in situations affected by flooding, debris avalanching, water-logging, drought and/or frost.

## Features\*:

Deciduous shrub or small tree up to 10 m tall. Trunk up to 1 m diam., bark thick, somewhat corky, grey, persistent, deeply marked with longitudinal furrows. Branches one to many, often spreading. Branchlets slender, grooved, glabrescent; bark red, red-brown to bronze-red. Adult leaves 2-4 on short shoots or widely spaced along fast growing branchlets; petioles 5 mm, slender; leaf lamina 20-50 x 5-20 mm, grey-green to green above, silvery-grey beneath, narrow-oblong, oblong-ovate to broadly-ovate, undersides clad in silvery tomentum, upper surface glabrescent; lamina margins flat and entire. Capitula in fascicles of 2-6, 5 x 5 mm; pedicels slender, silky hairy, 15 mm long. Florets 20-25, pale yellow, ray-florets 10-15, narrow, rather short, disc florets 10-15. Phyllaries in 2 series, weakly imbricate, oblong, obtuse, exposed surface pilose hairy. Achenes 1-2 mm, narrow-obovate. Pappus-hairs 3-5 mm long.

## Flowering:

October - December

## Fruiting:

December - February

## Threats:

This species is seriously threatened by recruitment failure. The seed of this species requires open sites to germinate in, and in most places such sites are scarce due to the presence of introduced grasses and herbs. Very few *O. hectorii* populations occur on protected land, and many are now dominated by old senescent trees. This species is also susceptible to browsing animals, and because of the dynamic habitats it occupies floods and slips once so critical for this species regeneration is now a serious threat. Isolated plants produce little viable seed.

## \*Attribution:

Fact Sheet prepared for the NZPCN by P.J. de Lange 14 April 2006. Description by P.B Heenan (adapted from Heads (1998) and subsequently published in de Lange et al. (2010).

## References and further reading:

de Lange, P.J.; Heenan, P.B.; Norton, D.A.; Rolfe, J.R.; Sawyer, J.W.D. 2010: Threatened Plants of New Zealand. Canterbury University Press, Christchurch.

Heads, M. 1998. Biodiversity in the New Zealand divaricating tree daisies: *Olearia* sect. nov. (Compositae). Botanical Journal of the Linnean Society 127(3): 239-285.

Hooker, J.D. 1864. *Handbook of the New Zealand Flora: a systematic description of the native plants of New Zealand and the Chatham, Kermadec's, Lord Auckland's, Campbell's and Macquarie's Islands.* Part I ed. London, Reeve. 392 p.

## For more information, visit:

[http://nzpcn.org.nz/flora\\_details.asp?ID=115](http://nzpcn.org.nz/flora_details.asp?ID=115)



**Caption:** Tree in Matukituki Valley

**Photographer:** John Barkla



**Caption:** Swales Bush Scenic Reserve

**Photographer:** Gillian Crowcroft

# *Olearia lineata*

## Common Name(s):

None known

## Current Threat Status (2012):

At Risk - Declining

## Distribution:

Endemic. South Island, easterly from north Canterbury south to Southland and Stewart Island.

## Habitat:

Lowland to montane (10-300 m a.s.l.) grey scrub, tussock grassland and forest margins. Often on river terraces in or near seepages and ephemeral wetlands, on occasion even growing in shallow water. Also found on the margins of steep river gorges, and in and amongst rock outcrops, boulder field and at the toe of alluvial fans.

## Features:

Small tree up to 8 m tall with narrow to broad canopy crowns. Trunk stout, erect, solitary, sometimes several arising from the ground, up to 0.6 m d.b.h. Bark grey or charcoal-grey, firm, deeply furrowed, shedding in tough, corky shards. Branches sparse to numerous, at first ascending then widely spreading; branchlets grey to charcoal grey, more or less square and angled in cross-section, deeply and longitudinally grooved, slender, at first erect then spreading, ultimately pendulous. Brachyblasts 10-30 mm long distantly spaced. Leaves 2-10-fascicled; 20-60 x 0.4-0.8 mm, linear to very narrow-linear, upper surface dark green more or less covered with finely appressed greyish-white indument, glabrate to glabrous with age, undersides clad in soft, white to greyish-white appressed tomentum, margin often strongly revolute. Capitula discoid, 1-8-fascicled, 2-4(-6) mm diameter, pedicellate, pedicels up to 40 mm long; florets 6-10, off-white to white (rarely creamy yellow), involucre bracts 2-4-seriate, narrowly lanceolate to oblanceolate, undersides finely grey-white villous. Cypselas 1-2 mm long, compressed, finely pubescent, puberulent to glabrescent, pappus hairs 2-3 mm long, off white to buff.

## Flowering:

November - January

## Fruiting:

January - April

## Threats:

Widespread and at times locally abundant (especially in some parts of Central Otago) *O. lineata* is otherwise often known from only widely scattered sites with few individuals. Although widespread the majority of the known populations are not officially protected and recruitment is often lacking. *Olearia lineata* together with the majority of Eastern South Island endemic *Olearia* Sect. *Divaricaster* Heads is the subject of a major Department of Conservation initiated Recovery Plan. As part of that work this species has been subject to intensive survey.

## For more information, visit:

[http://nzpcn.org.nz/flora\\_details.asp?ID=294](http://nzpcn.org.nz/flora_details.asp?ID=294)



**Caption:** Hunter Valley

**Photographer:** John Barkla



**Caption:** Bark, Hunter Valley

**Photographer:** John Barkla



## *Ozothamnus vauvilliersii*

**Common Name(s):**

mountain tauhinu

**Current Threat Status (2012):**

Not Threatened

**Threats:**

Not Threatened

**For more information, visit:**

[http://nzpcn.org.nz/flora\\_details.asp?ID=1606](http://nzpcn.org.nz/flora_details.asp?ID=1606)



**Caption:** Ozothamnus vauvilliersii

**Photographer:** John Barkla



**Caption:** Ozothamnus

**Photographer:** John Barkla

## *Phormium cookianum* subsp. *cookianum*

### Common Name(s):

Mountain flax, wharariki

### Current Threat Status (2012):

Not Threatened

### Distribution:

Endemic. Scarce in North Island where only known from high alpine situations in the Tararua Ranges, and possible elsewhere within the central axial ranges. Common in the South Island, in subalpine/alpine situations.

### Habitat:

Strictly confined to subalpine, alpine situations.

### Features:

Stout liliaceous herb, 0.6-1(-2) m tall. Leaves numerous, arising from fan-like bases. Individual leaves "paddle-shaped", erect, stiff, rarely decurved or pendulous 0.6-1(-1.5) x 20-80 mm, glaucous. Lamina margin, entire, somewhat thickened and distinctly pigmented by a dark, rather broad often encircling band 3-5 mm wide. Inflorescence (0.8-)1(-2) m tall, somewhat woody and fleshy when fresh, long persistent, drying charcoal grey or black, with the fibrous interior becoming progressively more exposed. Peduncle 20-30 mm diam., inclined, dark red-green, glabrous. Flowers 25-40 mm long, tubular, dull pink or yellow; tips of inner tepals markedly recurved. Ovary erect. Capsules 100-120 mm long, dark green, trigonous in cross-section, pendulous, tapering toward tip, twisted, initially fleshy becoming papery with age, long persistent. Seeds 8-10 x 4-5 mm, black, elliptic, flat and plate-like, margins frilled or twisted.

### Flowering:

(September-) October-  
November (-January)

### Fruiting:

(November-) December  
(-March)

### Threats:

Not Threatened

### For more information, visit:

[http://nzpcn.org.nz/flora\\_details.asp?ID=1110](http://nzpcn.org.nz/flora_details.asp?ID=1110)



**Caption:** Swampy Summit,  
Dunedin

**Photographer:** John Barkla



**Caption:** Hollyford Valley,  
Fiordland

**Photographer:** John Sawyer

## *Phormium tenax*

### Common Name(s):

flax, harakeke, korari (maori name for inflorescence).

### Current Threat Status (2012):

Not Threatened

### Distribution:

Indigenous to New Zealand and Norfolk Island. A broad circumscription has been adopted here - many botanists feel that plants from the Chatham Islands could be distinguished at species rank from the mainland New Zealand species, other distinctive variants occur on the Three Kings and outer Hauraki Gulf Islands, and along the Kaikoura coast. Norfolk Island plants though uniform differ in subtle ways from the New Zealand forms of *P. tenax*. Further study into this variation is underway.

### Habitat:

Common from lowland and coastal areas to montane forest, usually but not exclusively, in wetlands and in open ground along riversides.

### Features:

Stout liliaceous herb, 1-5(-6) m tall. Leaves numerous, arising from fan-like bases. Individual leaves rather stiff at first, but becoming decurved, somewhat pendulous or "floppy" in upper half to a third, 1-3 x 50-120 mm, usually blue-grey (glaucous) or dark green, lamina margin, entire, somewhat thickened and pigmented black, dark red, pink, yellow or cream. Inflorescence 5(-6) m tall, somewhat woody and fleshy when fresh, long persistent, drying charcoal grey or black, with the fibrous interior becoming progressively more exposed. Peduncle 20-30 mm diam., erect, dark grey-green or red-green, glabrous. Flowers 25-50 mm long, tubular, predominantly dull red but may also be pink or yellow; tips of inner tepals slightly recurved. Ovary erect. Capsules 50-100 mm long, dark green, red-green or black, trigonous in cross-section, erect, abruptly contract at tip, not twisted, initially fleshy becoming woody with age, long persistent. Seeds 9-10 x 4-5 mm, black, elliptic, flat and plate-like, margins frilled or twisted.

### Flowering:

(September-) October-November (-January)

### Fruiting:

(November-) December (-March)

### Threats:

Not threatened although see the discussion below about flax dieback. This die back phenomenon is characterised by abnormal yellowing of the leaves and may result in collapse of flax plants or whole populations.

### References and further reading:

Boyce, et al. 1951. Preliminary note on yellowleaf disease. NZJ of Science and Technology, 32(3): 76-77

Scheele, S. 1997. Insect pests and diseases of harakeke, Manaaki Whenua Press


### For more information, visit:

[http://nzpcn.org.nz/flora\\_details.asp?ID=2219](http://nzpcn.org.nz/flora_details.asp?ID=2219)



**Caption:** *Phormium tenax*

**Photographer:** Wayne Bennett



**Caption:** Flowers of *Phormium tenax*

**Photographer:** Wayne Bennett

# *Pittosporum eugenioides*

**Common Name(s):**

Tarata, lemonwood

**Current Threat Status (2012):**

Not Threatened

**Distribution:**

Endemic. Common in the North and South Islands.

**Habitat:**

Common tree of regenerating and mature forest in coastal to montane situations.

**Features\*:**

Gynodioecious tree up to 12 m tall but usually much less. Trunk 0.6-1 m diam, stout, clad in persistent pale-grey bark, branches numerous, erect then spreading. Leaf buds sticky, resinous. Leaves borne on slender petioles 10-20 mm long, alternate, 50-100(-150) x 25-40 mm, yellow-green, green, more or less blotched and mottled with paler green or yellow-green (sometimes white), somewhat leathery, glossy, smelling strongly when crushed of ivy or resin, elliptic to elliptic-oblong, apex acute to subacute; leaf margin undulate (very rarely not so), midrib pale green. Inflorescences terminal, numerous, subcorymbose compound umbels. Flowers pale yellow to yellow, very fragrant. Peduncles 10-20 mm, pedicels 5 mm, both sparsely hairy. Sepals 2 mm, ovate to narrow-ovate, pale caducous. Petals 5, 5-7 mm long, narrow-oblong. Capsules 2-valved (rarely 3), 5-6 mm, ovoid to elliptic, caducous, seeds immersed in dark yellow viscid pulp, whole structure covered in long persistent papery endocarp.

**Flowering:**

October - December

**Fruiting:**

October - January

**Threats:**

Not Threatened

**\*Attribution:**

Fact sheet prepared for NZPCN by P.J. de Lange 30 August 2006. Description adapted from Cooper (1956).

**References and further reading:**

Cooper, R.C. 1956: The Australian and New Zealand species of *Pittosporum*. *Annals of the Missouri Botanical Garden* 43: 87-188

Gardner, R. 1999. Notes towards an excursion Flora. *Pittosporum eugenioides* as a wild plant. *Auckland Botanical Society Journal*, 54, 1


**For more information, visit:**

[http://nzpcn.org.nz/flora\\_details.asp?ID=1135](http://nzpcn.org.nz/flora_details.asp?ID=1135)



**Caption:** Masterton

**Photographer:** John Barkla



**Caption:** Maidstone Park, Upper Hutt.

**Photographer:** Jeremy Rolfe

# *Pittosporum tenuifolium*

## Common Name(s):

Kohukohu, kohuhu, black matipo

## Current Threat Status (2012):

Not Threatened

## Distribution:

Endemic and widespread throughout country.

## Habitat:

A small tree of coastal to montane shrubland and forested habitats. Preferring successional habitats.

## Features\*:

Shrub or small gynodioecious tree up to 10 m tall (usually much less). Trunk 0.3-0.4(-0.6) m diam., stout, clad in dark grey-black or brown persistent bark. Branches numerous, erect then spreading. Branchlets and young leaves pubescent, hairs pale yellow or cream. Petioles short, somewhat fleshy. Leaves alternate, (10-)30(-70) x (5-)10(-20) mm, leathery, pale-green to dark green above, lighter below, oblong, oblong-ovate or elliptic-obovate, apex obtuse to acute, rarely acuminate, margins entire, often undulose. Flowers solitary or in axillary cymes, rather fragrant, especially at night. Pedicels stout, pale green, fleshy, bracts entire, lanceolate, caducous. Sepals narrowly ovate-oblong, subacute to obtuse, silky hairy. Petals 12 mm long, lanceolate, dark red, black (rarely yellow or white). Capsules 2-valved (rarely 3), subglobose, valves woody, black when mature, long persistent. Seeds immersed in sticky, red or yellow viscid pulp.

## Flowering:

October - November (-December)

## Fruiting:

January - March

## Threats:

Not Threatened

## \*Attribution:

Fact sheet prepared for NZPCN by P.J. de Lange 10 January 2004. Description adapted from Allan (1961).

## References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Vol. I. Wellington, Government Printer

## For more information, visit:

[http://nzpcn.org.nz/flora\\_details.asp?ID=1139](http://nzpcn.org.nz/flora_details.asp?ID=1139)



**Caption:** *Pittosporum tenuifolium* in flower Dunedin

**Photographer:** John Barkla



**Caption:** Quail Island

**Photographer:** John Barkla

# *Pseudopanax crassifolius*

## Common Name(s):

Horoeka, lancewood

## Current Threat Status (2012):

Not Threatened

## Distribution:

Endemic. North, South and Stewart Islands. Widespread and common

## Habitat:

Lowland to montane forest. Sealevel to c. 750 m a.s.l.

## Features\*:

Bushy topped tree to 15 m tall, branchlets fleshy, trunk unbranched in lower part, to 50 cm diam., distinctly ridged when young, bark dark becoming paler with age, wood tough. Leaves alternate; leaflets 1-3 in seedling, palmate, sessile or subsessile on very short petiolule, submembranous coarsely toothed, absent from juvenile and adult. Juvenile leaves dark green, narrow-linear, deflexed, to 1 m long, coriaceous, midrib pale cream-yellow, raised, margins distantly sharply toothed, distal margin of tooth perpendicular to midvein, not swollen. Adult leaves shorter, 10-20 x 2-3 cm, dark green, very occ. trifoliate (probably due to hybridisation with other species), narrow elliptic-cuneate to lanceolate or linear-obovate, acute or obtuse, margins entire to sunuate or coarsely serrate, subsessile or on petioles to 10 mm long, petiole base expanded around stem. Inflorescence a terminal umbel, irregularly compound; primary rays (branchlets) 5-10, c. 6 cm long; umbellules sometimes racemosely arranged. Ovary 5-loculed, each containing 1 ovule; style branches 5, connate, tips sometimes free. Fruit fleshy, subglobose, 4-5 mm diam., style branches retained on an apical disc, dark purple when ripe. Seeds 4-5 per fruit, easily separated, broadly ovate, grooved, 2.2-3.5(-5.5) mm long.

## Flowering:

January-April

## Fruiting:

January-April

## Threats:

Not Threatened

## \*Attribution:

Description adapted from Allan (1961) and Webb and Simpson (2001).

## References and further reading:

Allan, H.H. 1961. Flora of NZ, Vol. I. Government Printer, Wellington

Webb, C.J. & Simpson, M.J.A. 2001. Seeds of NZ gymnosperms and dicotyledons. Manuka Press, Christchurch.

## For more information, visit:

[http://nzpcn.org.nz/flora\\_details.asp?ID=1196](http://nzpcn.org.nz/flora_details.asp?ID=1196)



**Caption:** *Pseudopanax crassifolius*

**Photographer:** Wayne Bennett



**Caption:** Seeds of *Pseudopanax crassifolius*

**Photographer:** Wayne Bennett

## *Rubus schmidelioides* var. *schmidelioides*

### Common Name(s):

Tataramoa, bush lawyer, white-leaved lawyer

### Current Threat Status (2012):

Not Threatened

### Distribution:

Endemic. New Zealand: North Island (from about Hikurangi - Dargaville South), South and Stewart Islands

### Habitat:

Coastal to montane in scrub and forest.

### Features\*:

Much-branched dioecious vine with stems up to 10 m or more long when growing through supporting vegetation or forming sprawling bushes with interlacing branchlets when exposed. Young branchlets terete, up to 40 mm diameter; pubescent, becoming glabrous with age, armed or unarmed; when armed then armed with small, reddish prickles. Stipules minute, caducous, linear to linear-filiform. Leaves of adults ternate to palmate, petioles 20-50 mm long; lamina of adult leaflets 30-80 × 10-50 mm, coriaceous, ovate, oval, lanceolate, elliptic to broadly elliptic, acute, subacute to obtuse, rounded to oblique at base, margins coarsely and often unevenly serrate, adaxially dark green to bronze-green, more or less glossy, glabrous or almost glabrous, rugose to bullate, veins distinct (these often minutely pubescent), abaxially glaucous to white, usually finely, densely tomentose, tomentum white to grey-white, veins prominent below, pubescent, midrib armed or not; petiolules 3-5(-20 mm long). Inflorescence a much-branched panicle up to 100 mm long (rarely reduced to few-flowered racemes), branchlets and pedicels unarmed, more or less finely pubescent. Flowers white to cream, usually densely aggregate, on pedicels 5-10 mm long. Sepals 2-6 mm long, ovate-oblong to broad-ovate, pubescent; petals 5, 4-7 mm long, ovate to broadly ovate. Male flowers with numerous stamens, ovary absent or rudimentary. Female flowers with rudimentary stamen, carpels numerous. Fruits 5-9 mm diameter, drupelets 8-12 (or more), yellow or orange. Endocarp 'seed' 1.9-2.5 mm long, dorsally ridged 2x.

### Flowering:

September - November

### Fruiting:

December - April

### Threats:

Not Threatened

### \*Attribution:

Fact Sheet prepared for NZPCN by P.J. de Lange 26 October 2016. Description based on Allan (1961), Webb et al. (1988) and Webb & Simpson (2001)

### References and further reading:

Allan, H.H. 1961: Flora of New Zealand. Vol. I, Wellington, Government Printer.

Webb CJ, Sykes WR, Garnock-Jones PJ 1988: Flora of New Zealand. Vol. IV. Botany Division, DSIR, Christchurch.

Webb, C.J.; Simpson, M.J.A. 2001: Seeds of New Zealand Gymnosperms and Dicotyledons. Christchurch, Manuka Press.

### For more information, visit:

[http://nzpcn.org.nz/flora\\_details.asp?ID=1265](http://nzpcn.org.nz/flora_details.asp?ID=1265)



**Caption:** Whirinaki, October  
**Photographer:** John Smith-Dodsworth



**Caption:** Whirinaki, October  
**Photographer:** John Smith-Dodsworth

## Definitions of botanical terms

A glossary has been provided below with definitions for many of the botanical terms used in the species descriptions.

### Glossary

Term	Definition
<b>Abaxial</b>	Facing away from the stem of a plant (especially denoting the lower surface of a leaf).
<b>Acerose</b>	Narrow with a sharp stiff point.
<b>Achene</b>	A simple, dry, one-seeded (one-celled) fruit
<b>Acicular</b>	Needle-shaped.
<b>Acidic</b>	Having a low pH, opposite of basic or alkaline.
<b>Acroscopic</b>	Pointing towards, or on the side of, the apex
<b>Acuminate</b>	Gradually tapered to a point. Sharply pointed.
<b>Acute</b>	Pointed or sharp, tapering to a point with straight sides.
<b>Adnate</b>	Fusion of unlike parts, e.g. stamens fused to petals.
<b>Adventive</b>	A plant that grows in the wild in New Zealand but which was introduced to the country by humans.
<b>Agglutinated</b>	Stuck together.
<b>Allelopath</b>	An organism that releases compounds that are toxic to other species.
<b>Allelopathy</b>	The release by an organism of compounds that are toxic to other species.
<b>Alternate</b>	Attached singly at each node but changing from one side of a stem to the other.
<b>Alveolate</b>	Honeycombed with ridged partitions.
<b>Amplexicaul</b>	clasping or surrounding the stem
<b>Anamorph</b>	Asexual fruiting stage, usually of an ascomycete fungus.
<b>Anastomosing</b>	Rejoining after branching, as in some leaf veins.
<b>Annual</b>	A plant that completes its complete life cycle within the space of a year
<b>Annual evergreen</b>	Plants that lose their over-wintering leaves rapidly in the first half of the growing season. Annual evergreens never present a leafless appearance, but are closer in a functional sense to a deciduous plant than they are to multi-annual evergreens.
<b>Annulus</b>	Line of thickened cells that governs the release of spores from a sporangium
<b>Anterior</b>	Towards the front.
<b>Anther</b>	The pollen-bearing portion of the stamen.
<b>Antheridium</b>	Male reproductive organ formed on the prothallus of a fern
<b>Anthesis</b>	When the flower is fully developed and functioning. The time of pollination or bloom.
<b>Apex</b>	Tip; the point furthest from the point of attachment.
<b>Apices</b>	Plural of apex. Tip, the point furthest from the point of attachment
<b>Apiculate</b>	Bearing a short slender and flexible point.
<b>Apiculus</b>	A small, slender point.
<b>Apomixis</b>	A form of reproduction whereby seed is formed without the usual mode of sexual fusion
<b>Appressed</b>	Pressed against another organ or surface.
<b>Aquatic</b>	Growing, or living in, or frequenting water. Applied to plants and animals and their habitats. Opposite of terrestrial (land living).
<b>Archegonium</b>	Female reproductive organ of a fern formed on the prothallus
<b>Arcuate</b>	Curved into an arch.
<b>Aril</b>	An often fleshy appendage on the outside of a seed.
<b>Artificial thinning</b>	Selectively removing vegetation to create gaps to facilitate natural invasion of native plants, or to plant later successional plants.
<b>Ascending</b>	Growing obliquely upward.
<b>Asexual</b>	Vegetative reproduction, lacking sexual involvement by sperm or egg cells
<b>Attenuate</b>	Narrowing gradually
<b>Auricle</b>	A small, ear-shaped appendage.
<b>Auriculate</b>	Bearing a small, ear-shaped appendage.
<b>Autogamous</b>	Self-fertilising flowers.
<b>Autotrophic</b>	Of or relating to organisms (as green plants) that can make complex organic nutritive compounds from simple inorganic sources by photosynthesis
<b>awn</b>	A stiff or bristle like projection often from the tip or back of an organ
<b>Axil</b>	The upper angle between the leaf and the stem.
<b>Axis</b>	The longitudinal supporting structure around which organs are borne, e.g., a stem bearing leaves.
<b>Barbellate</b>	Barbed, having or covered with protective barbs or quills or spines or thorns or setae
<b>Basal</b>	At the base.
<b>Basisopic</b>	Pointing towards the base
<b>Beak</b>	A prominent extension of an organ
<b>Bifid</b>	Deeply split into two lobes.
<b>Bifurcate</b>	Divided into two.



<b>Term</b>	<b>Definition</b>
<b>Biosecurity</b>	Preventing, eradicating, controlling and managing risks posed by pests and diseases.
<b>Biotic</b>	Pertaining to the living parts of the environment
<b>Bipinnate</b>	With each primary pinna divided to the midrib into a secondary pinna
<b>Biserrate</b>	Doubly serrate.
<b>Blade</b>	The flattened part of a leaf.
<b>Blunt</b>	Not pointed at the ends
<b>Bog</b>	A quagmire covered with specialised plants including sphagnum moss, grasses, sedges, rushes, sundews, umbrella ferns and other plants; has wet, spongy ground, a marsh-plant community on wet, very acid peat. Fed only by rainfall.
<b>Bottleneck</b>	A genetic term; refers to the fact that in smaller populations there could be lower genetic variability
<b>Brachyblasts</b>	Short shoots
<b>Bract</b>	A reduced leaf or leaf-like structure at the base of a flower.
<b>Bracteate</b>	Bearing bracts: leaves or leaf-like structure reduced at the base of a flower.
<b>Bracteolate</b>	With small bracts.
<b>Bracteole</b>	A small bract.
<b>Bracteoles</b>	Bracts directly below the flower
<b>Brevideciduous</b>	Brief (1 month or less) loss of most leaves from the canopy just before flowering or during flushing of a new cohort of leaves.
<b>Bryophyte</b>	Plant group including mosses, liverworts and hornworts
<b>Bryophytes</b>	Plant group including mosses, liverworts and hornworts
<b>Bulbil</b>	A bud produced vegetatively on the stem or frond that is capable of breaking off and growing into a new plant
<b>Bullate</b>	With rounded projections covering the surface as if blistered
<b>Caespitose</b>	Growing in dense tufts
<b>Calli</b>	Circular, warty, stalked thickenings commonly found on the lip (labellum) of the orchid (plural of callus).
<b>Callose</b>	Hardened or thickened.
<b>Callus</b>	Stalked thickening on the lip (labellum) of an orchid.
<b>Calyx</b>	The group of sepals, or outer floral leaves, of a flower
<b>Campanulate</b>	Bell-shaped.
<b>Canaliculate</b>	With longitudinal channels or grooves.
<b>Canopy</b>	The uppermost cover formed by the branches and leaves of trees or the spread of bushes, shrubs and ground covers.
<b>Canopy closure</b>	Stage where canopies of shrub and tree species meet.
<b>Canopy manipulation</b>	Selectively removing vegetation to create gaps to facilitate natural invasion of native plants, or to plant later successional plants.
<b>Capillary</b>	Hair-like
<b>Capitula</b>	Plural of capitulum: A dense head-like inflorescence of many flowers as occurs in most Asteraceae (daisies)
<b>Capitulum</b>	A dense head-like inflorescence of many flowers as occurs in most Asteraceae (daisies)
<b>Capsule</b>	A dry fruit formed from two or more fused carpels that splits open when ripe.
<b>Carbon sinks</b>	Carbon locked away, or sequestered e.g. by trees
<b>Carpel</b>	One unit of the female part of a flower that consists of a basal seed-bearing ovary joined to a receptive stigma by a stalk-like style.
<b>Cauda</b>	Tail-like appendage. (pl. caudae; adj. caudate)
<b>Caudex</b>	The axis of a woody plant, esp. a palm or tree fern, comprising the stem and root.
<b>Cauline</b>	Belonging to the stem, as in cauline leaves emerging from the stem.
<b>Cerise</b>	Bright or deep red.
<b>Chartaceous</b>	Having a papery texture.
<b>Chlorophyll</b>	The green pigment of plants.
<b>Chlorotic</b>	Lacking chlorophyll, therefore yellowish, suffering from chlorosis.
<b>Cilia</b>	Short small hair-like structures on a cell or microorganism
<b>Ciliate</b>	With small hairs (cilia).
<b>Ciliolate</b>	Diminutive of ciliate, i.e., having very small hairs
<b>Cladode</b>	Flattened stem with the function of a leaf
<b>Cladodes</b>	Usually flattened, photosynthetically active branches, these may be leaf-like (e.g., Phyllocladus) or branch-like (e.g., Carmichaelia)
<b>Clavate</b>	Club-shaped, gradually widening towards apex.
<b>Cleft</b>	Having indentations that extend about halfway to the center, as in certain leaves.
<b>Cleistogamous</b>	Flowers that self-fertilise without opening.
<b>Coherent</b>	Sticking together of like parts.
<b>Column</b>	Stamen and stigmas fused to form a single organ.

<b>Term</b>	<b>Definition</b>
<b>Columnar</b>	Shaped like a column
<b>Composite</b>	many small flowers tightly packed together e.g., daisy flowers.
<b>Compound</b>	Composed of several similar parts (cf simple)
<b>Concave</b>	Curved inward.
<b>Concolorous</b>	Of the same colour.
<b>Conical</b>	Cone-shaped.
<b>Connate</b>	Fusion of like parts.
<b>Conspecific</b>	Individuals of the same species.
<b>Cordate</b>	Heart-shaped with the notch at the base.
<b>Coriaceous</b>	Leather-like; thick, tough, and somewhat rigid.
<b>Corolla</b>	The whorl of petals of a flower.
<b>Corymb</b>	Modified raceme where stalks of lower flowers are elongated to same level as the upper flowers.
<b>Cosmopolitan</b>	A species or other taxonomic group that is distributed widely throughout the world.
<b>Costa</b>	The midrib
<b>Crenate</b>	With rounded teeth (bluntly toothed) along the margin.
<b>Crisped</b>	Margin tightly wavy or crinkled, curled or wavy.
<b>Cristate</b>	With a crest.
<b>Crown</b>	The growing point of an upright rhizome or trunk. This usually produces a tuft or ring of fronds.
<b>Crura</b>	The two small projections at the mouth of a utricle in Carex
<b>Cucullate</b>	Hood-shaped.
<b>Culm</b>	The erect stem of a grass.
<b>Cuneate</b>	Wedge-shaped.
<b>Cupular</b>	Cup-shaped.
<b>Cuttings</b>	Stems and/or leaves taken from plants for propagation
<b>Cyathium</b>	A cup-like structure that surrounds the inflorescence in Euphorbia
<b>Cyme</b>	Inflorescence at the terminus of a branch and where new flowering branches emerge laterally below the flower.
<b>Cytorace</b>	Populations (or infraspecific taxa) that differ in chromosome number or chromosome morphology, e.g., <i>Nematoceras trilobum</i> agg. has two cytoraces, a diploid and a tetraploid (in which the chromosomes are doubled).
<b>Cytotype</b>	Populations (or infraspecific taxa) that differ in chromosome number or chromosome morphology, e.g., <i>Nematoceras trilobum</i> agg. has two cytotypes, a diploid and a tetraploid (in which the chromosomes are doubled).
<b>Deciduous</b>	Marked leaflessness in winter, and greater than 90% leaves lost by beginning of spring flush.
<b>Decrescent</b>	Diminishing.
<b>Decumbent</b>	With a prostrate or curved base and an erect or ascending tip.
<b>Decurrent</b>	Attached by a broadened base.
<b>Decurved</b>	Curved downward.
<b>Deflexed</b>	Bent abruptly downward.
<b>Dehiscence</b>	The time of opening at maturity to release the contents, e.g., a capsule releasing the seeds.
<b>Dehiscent</b>	Splitting open at maturity to release contents (of a fruit).
<b>Deltoid</b>	Shaped broadly like an equilateral triangle.
<b>Dentate</b>	Toothed along the margin with the teeth pointing outward, not forward.
<b>Denticles</b>	minute teeth
<b>Denticulate</b>	having a very finely toothed margin
<b>Dichotomous</b>	Divided into two equal branches.
<b>Digitiform</b>	Finger-like.
<b>Dioecious</b>	Having male and female flowers on separate plants of the same species.
<b>Diploid</b>	With two complete sets of chromosomes in each cell.
<b>Disarticulating</b>	Separating at a joint.
<b>Discoid</b>	Disc-shaped.
<b>Disjunct</b>	A species or other taxonomic group that occupies areas that are widely separated and scattered and therefore have a discontinuous distribution.
<b>Distal</b>	Toward the apex, away from the point of attachment (cf. proximal).
<b>Distichous</b>	In two rows on opposite sides of the axis.
<b>Divaricating</b>	Branching at a very wide angle with stiff intertwined stems.
<b>Domatia</b>	small structures on the lower surface of a leaf in some woody dicotyledons, located in the axils of the primary veins and usually consisting of depressions partly enclosed by leaf tissue or hairs.

<b>Term</b>	<b>Definition</b>
<b>Dorsal</b>	Of the back or outer surface relative to the axis. (cf. ventral)
<b>Drupe</b>	A stone fruit, the seed enclosed in a bony covering (endocarp) which is surrounded by a + fleshy layer (mesocarp)
<b>Early successional species</b>	Plants which are able to colonise an open area after disturbance but which are often temporary and are replaced by taller plants in time and shaded out.
<b>Echinate</b>	having sharply pointed spines or bristles.
<b>Ecological district</b>	A characteristic landscape and biological community defined in the PNA (Protected Natural Area) programme.
<b>Ecological restoration</b>	Attempt to reinstate original (pre-disturbance) state of a habitat, plant community or ecosystem.
<b>Ecosourced</b>	Plants sourced from seed collected from similar naturally growing plants in the area of the planting site.
<b>Ecosourcing</b>	Using native plants grown from locally grown seeds. Eco-sourced plants help to preserve the ecological distinctiveness of an area, and ecosourced plants fare better and are adapted to survive in the local conditions.
<b>Eglandular</b>	Without glands.
<b>Elaiosome</b>	Fleshy, oil-rich structure attached to seed that attracts ants which act as dispersers.
<b>Ellipsoid</b>	Elliptic in long section and circular in cross-section.
<b>Elliptic</b>	Broadest at the middle
<b>Emarginate</b>	With a notch at the apex.
<b>Emarginated</b>	Having a shallow notch at the tip, as in some petals and leaves.
<b>Emergent</b>	In an aquatic sense - wetland herbs that are rooted in the substrate below water level, but carry leaves and stems above the water level e.g. rushes and raupo. Found on the shallow margins of lakes, ponds and waterways. In a forest sense - tree that is appearing above the surrounding canopy.
<b>Emergent marginals</b>	An aquatic plant having most of its structure above water. Other aquatic plants are submerged or floating.
<b>Endemic</b>	Unique or confined to a place or region, found naturally nowhere else.
<b>Endophyte</b>	An endosymbiont (usually a bacterium or fungus) that lives within a plant for at least part of its life without causing any apparent disease.
<b>Endophytes</b>	Endosymbionts (usually bacteria or fungi) that live within plants for at least part of their lives without causing any apparent disease.
<b>Endosperm</b>	The nutritive tissue of a seed, consisting of carbohydrates, proteins, and lipids.
<b>Enrichment planting</b>	Returning to a revegetation site and creating gaps, or filling existing gaps, with different plants of plants, usually later successional plants which may not have survived being planted in the first phases of the project.
<b>Ensiform</b>	Sword shaped
<b>Entire</b>	Smooth. Without teeth, notches or divisions.
<b>Entomophilous</b>	Pollinated by insects.
<b>Epicalyx</b>	Calyx-like structure outside, but close to, the true calyx.
<b>Epigeal</b>	Growing on or close to the ground or emerging from the ground after germination (often used for cotyledons).
<b>Epiphyte</b>	A plant that grows upon another plant but is not parasitic and does not draw nourishment from it.
<b>Epiphytic</b>	Growing upon another plant but not parasitic and not drawing nourishment it
<b>Erose</b>	Irregularly toothed, as if gnawed.
<b>Estuarine</b>	Pertaining to the meeting of freshwater and seawater wetlands.
<b>Ethnobotany</b>	The study of people's classification, management and use of plants.
<b>Eusporangia</b>	Sporangia that arise from groups of epidermal cells
<b>Evanescent</b>	Lasting a very short time or running a short distance.
<b>Ex situ</b>	Away from the place of natural occurrence.
<b>Ex-situ</b>	Maintenance of plants as live specimens or propagules in cultivation as insurance against the loss of wild populations and as source for material for translocation.
<b>Excurrent</b>	Having the axis prolonged to form an undivided main stem or trunk (as in conifers).
<b>Extravaginal</b>	Outside an enclosing sheath
<b>Falcate</b>	Hooked or curved like a sickle.
<b>Fastigate</b>	Branches erect and close to central axis.
<b>Fen</b>	A type of wet land that accumulates peat deposits. Fens are less acidic than bogs, deriving most of their water from groundwater rich in calcium and magnesium.
<b>Ferruginous</b>	Rust-like (a colour term)
<b>Fertile frond</b>	Fronds that bear sporangia.
<b>Filamentous</b>	Resembling a filament.
<b>Filiform</b>	Thread like, resembling a filament.
<b>Filiramulate</b>	Branching at a very wide angle with stiff intertwined stems.
<b>Fimbriae</b>	Plural of fimbria: Fringe. A fimbria is composed of many fimbriae (individual hair-like structures).
<b>fimbriate</b>	With fringes.
<b>Flabellate</b>	Fan shaped.
<b>Flaccid</b>	Limp, not rigid, flabby.
<b>Flange</b>	A projecting rim.

Term	Definition
<b>Flexuose</b>	With curves or bends.
<b>Floccose</b>	Having tufts of soft woolly hairs
<b>Floret</b>	A small flower, usually one of a cluster - the head of a daisy for example.
<b>Foliaceous</b>	Leaf-like.
<b>Foliate</b>	Having leaflets.
<b>Founder effect</b>	When a small number of plants (and therefore their genes) from a larger population are selected some genetic information is lost.
<b>Fronnd</b>	A leaf, the complete leaf of a fern including the stipe and lamina
<b>Fulvous</b>	Orange–yellow.
<b>Funneliform</b>	Funnel-shaped.
<b>Fusiform</b>	Broadest near the middle and tapering toward both ends.
<b>Galea</b>	Helmet- or hood-shaped.
<b>Galeate</b>	Shaped like a helmet or hood.
<b>Gametophyte</b>	A plant that produces sperm and egg cells and in which sexual reproduction takes place - in ferns this is known as the prothallus
<b>Gene pool</b>	The mixture of all genes and gene variations of a group or population.
<b>Genetic diversity</b>	The variety of genes in a plants or populations.
<b>Genetic variation</b>	Differences displayed by individuals within a plant which may be favoured or eliminated by selection.
<b>geniculate</b>	abruptly bent
<b>Genus</b>	A taxonomic rank of closely related forms that is further subdivided in to species (plural = genera). In a scientific name (e.g., <i>Sicyos australis</i> ), the first word is the genus, the second the species.
<b>Gibbous</b>	Swollen or enlarged on one side, as in a gibbous moon.
<b>Glabrescent</b>	Lacking hair or a similar growth or tending to become hairless
<b>Glabrous</b>	Without or devoid of hairs, smooth.
<b>Gland</b>	A structure that secretes a sticky or oily substance.
<b>Glandular</b>	A structure that secretes a sticky or oily substance.
<b>Glaucous</b>	Covered with a fine, waxy, removable powder that imparts a white or bluish cast to the surface.
<b>Gley</b>	A soil prone to seasonal inundation.
<b>Globose</b>	Globe-shaped.
<b>Glume</b>	One of two bracts at the base of a grass spikelet.
<b>Groundwater</b>	Groundwater is the water beneath the surface that can be collected with wells, tunnels, or drainage galleries, or that flows naturally to the earth's surface via seeps or springs. Groundwater is the water that is pumped by wells and flows out through springs.
<b>Gymnosperm</b>	Plants in the class Gymnospermae that have seeds which are not enclosed in an ovary.
<b>Gynodioecious</b>	A species population containing plants that produce bisexual (perfect) flowers, and plants that produce only female (pistillate) flowers.
<b>Gynoeceium</b>	The female reproductive organs of a flower; the pistil or pistils considered as a group. Means literally "womans house" i.e., the overall structure that contains the female sex organs
<b>Hastate</b>	Spear like. Shaped like an arrowhead, but with basal lobes pointing outward rather than downward.
<b>Haustorium</b>	The absorbing organ of a parasite or hemiparasite
<b>Hemi–parasite</b>	Obtains water and nutrients from the roots of other plants but also manufactures food through photosynthesis.
<b>Hemi–parasitic</b>	Obtaining water and nutrients from the roots of other plants then manufacturing food through photosynthesis.
<b>Herbarium</b>	The place where collections of dried/pressed plants are kept.
<b>Hermaphrodite</b>	Having both male and female sexual characteristics and organs.
<b>Heteroblastic</b>	Exhibiting differences in leaf shapes or forms in juvenile and adult phases of the plant.
<b>Heteroblasty</b>	The state of being heteroblastic (i.e., exhibiting differences in leaf shapes or forms in juvenile and adult phases of the plant).
<b>Hirsute</b>	Hairy.
<b>Hyaline</b>	Membranous, thin and translucent.
<b>Hybrid</b>	An individual that is the offspring of a cross between two different varieties or species.
<b>Hybridise</b>	Breeding with a member of a different plant or type.
<b>Hydrophyte</b>	A plant species adapted to growing in or on water or in wet situations. Aquatic or semi-aquatic.
<b>Hymenium</b>	The fertile, spore–bearing layer of a fruitbody.
<b>Hypanthium</b>	A ring–like, cup–shaped, or tubular structure of a flower on which the sepals, petals, and stamens are borne.
<b>Imbricate</b>	Overlapping.
<b>imbricating</b>	Overlapping.
<b>Imparipinnate</b>	Odd–pinnate, a leaf shape; pinnate with a single leaflet at the apex.
<b>In-situ</b>	On site conservation relating to the maintenance of plants in the wild.
<b>Inbreeding</b>	Genetic similarity in offspring of closely related individuals.

Term	Definition
<b>Incoherent</b>	Not sticking together.
<b>Incursion</b>	Entrance of a pest into an area where it is not present
<b>Indumentum</b>	A covering of fine hairs (or sometimes scales)
<b>Indusia</b>	Plural of indusium, a membrane covering a sorus of a fern
<b>Indusium</b>	A thin tissue that covers the sorus in many ferns. Plural: indusia.
<b>Inflorescence</b>	The arrangement of flowers on the stem. A flower head.
<b>Infundibuliform</b>	Funnel-like.
<b>Interkeel</b>	The space between the keel and the leaf blade
<b>Internode</b>	The part of an axis between two nodes; the section of the stem between leaves.
<b>Internodes</b>	Part of a stem between two nodes.
<b>Intramarginal</b>	Within or near the margin.
<b>Involucral bracts</b>	The scales surrounding the flower head or capitula.
<b>Involucre</b>	A group of bracts surrounding a flower head.
<b>Involute</b>	With margins rolled inward toward the upper side.
<b>Irritable</b>	Responding to touch.
<b>Jugate</b>	Paired.
<b>Juvenile</b>	A plant of non-reproducing size.
<b>Keel</b>	A prominent or obvious longitudinal ridge (as in a boat).
<b>Labellar</b>	Pertaining to the labellum: a lip; in orchid flowers referring to the middle petal which usually differs in size, shape or ornamentation from the two lateral petals.
<b>Labellum</b>	A lip; in orchid flowers referring to the highly modified middle petal which usually differs in size, shape or ornamentation from the two lateral petals.
<b>Lacinia</b>	A jagged lobe.
<b>Laciniae</b>	Jagged lobes.
<b>Laciniate</b>	Cut into narrow, irregular lobes or segments.
<b>Lacustrine</b>	Of or having to do with a lake, of, relating to, or formed in lakes, growing or living in lakes.
<b>Lamina</b>	The expanded flattened portion or blade of a leaf, fern frond or petal.
<b>Lanceolate</b>	Lance-shaped; of a leaf several times longer than wide with greatest width about one third from the base, tapering gradually to apex and more rapidly to base
<b>Lateral</b>	On or at the side.
<b>Lax</b>	With parts open and spreading, not compact.
<b>Laxly</b>	With parts open and spreading, not compact
<b>Leaflet</b>	One section of a compound leaf.
<b>Lemma</b>	The lower of two bracts enclosing the flower in grasses.
<b>Lenticillate</b>	Bark that is covered in fine lenticles (breathing pores)
<b>Ligulate</b>	Strap-like, tongue-shaped
<b>Ligule</b>	The membrane between the leaf and the stem of a grass; the "petal" of a ray floret in a composite inflorescence
<b>Linear</b>	Long and narrow with more or less parallel sides.
<b>Littoral</b>	Occurring at the border of land and sea (or lake). On or pertaining to the shore. The shallow sunlit waters near the shore to the depth at which rooted plants stop growing.
<b>Lobe</b>	A recognisable, but not separated, rounded division or segment of a leaf or pinna. Used to describe ferns and leaves in <i>Cotula</i> and <i>Leptinella</i> .
<b>Lobed</b>	Part of a leaf (or other organ), often rounded, formed by incisions to about halfway to the midrib.
<b>Lobule</b>	A small lobe or sub-division of a lobe
<b>Lustrous</b>	Glossy, shiny.
<b>Lycophytes</b>	Seedless vascular plants that belong to the phylum Lycophyta (characterised by microphylls -primitive leaves found in ancient plants).
<b>Lyrate</b>	Pinnatifid or pinnatisect terminal lobe much larger than lower lobes.
<b>Maculate</b>	Blotched or spotted.
<b>Mangrove</b>	Coastal wetland dominated by Manawa or mangrove <i>Avicennia marina</i> var. <i>resifera</i> . Northern New Zealand only, salt marsh replaces it further south.
<b>Margin</b>	The edge or border of a leaf
<b>Marine</b>	Pertaining to the sea and saltwater systems.
<b>Marsh</b>	A tract of wet land principally inhabited by partially-submerged herbaceous vegetation. Has fewer woody plants than swamper habitats.
<b>Mealy</b>	Dry, powdery, crumbly.
<b>Median</b>	In the middle.
<b>Membranous</b>	Very thin, like a membrane.
<b>Mid-lobe</b>	The middle part into which a leaf is divided.
<b>Midrib</b>	The central or principal vein of a leaf or pinna of a fern.
<b>Mire</b>	Synonymous with any peat-accumulating wetland. Term covers bogs and peaty swamps, fens, carr, moor, muskeg and peatland. Term excludes marsh which is non-peat forming.

Term	Definition
<b>Molecular techniques</b>	Where proteins and genes are used to investigate plant relationships
<b>Monitoring</b>	Recording of quantitative data over time to document changes in condition or state of species or ecosystems.
<b>Monoecious</b>	Having male and female flowers on the same plant of the same species.
<b>Montane</b>	Land between 300 and 800 metres above sea level.
<b>Mucronate</b>	Tipped with a short, sharp, point.
<b>Mucronulate</b>	Having a very small mucro; diminutive of mucronate.
<b>Multi-annual evergreen</b>	Overlapping annual cohorts of leaves always present.
<b>Multifid</b>	Cleft into many lobes or segments
<b>Multiseptate</b>	With many septa.
<b>muricate</b>	Rough with short, hard points like the shell of Murex, a genus of tropical sea snails with elaborately pointed shells.
<b>Mycorrhiza</b>	A symbiotic relationship between a fungus and a plant.
<b>Mycorrhizal associations</b>	Symbiotic association between fungi and plant roots which assists plant health by allowing increased ability for uptake of nutrients and promote plant growth.
<b>Napiform</b>	A long swollen but tapering root – like a parsnip, or carrot.
<b>Native</b>	Naturally occurring in New Zealand (i.e., not introduced accidentally or deliberately by humans).
<b>naturalised</b>	Referring to plants that have escaped from cultivation (including gardens or forest plantations) and can now reproduce in the wild (without human assistance)
<b>Nectary</b>	Organ that produces nectar.
<b>Nerve</b>	Prominent vein or rib.
<b>Nerves</b>	Strands of conducting and usually strengthening tissue in a leaves or similar structures
<b>Net veins</b>	Veins that repeatedly divide and re-unite.
<b>Net venation</b>	Feather-like or hand-like venation on a leaf.
<b>Nival</b>	Growing at high altitudes. From Latin: nivalis, snowy etc. from nix, nivis, snow.
<b>Node</b>	The point at which leaves, branches or roots arise on a stem.
<b>Ob-</b>	Prefix meaning inverted, in reverse direction.
<b>Obcordate</b>	Heart shaped with the notch at the apex.
<b>Oblanceolate</b>	Tapering and widest towards the apex or inversely lanceolate.
<b>Oblique</b>	Slanting; of a leaf, larger on one side of the midrib than the other, in other words asymmetrical.
<b>Oblong</b>	Rectangular.
<b>Obovate</b>	Roughly elliptical or reverse egg shaped and widest near the apex (i.e., the terminal half broader than the basal half).
<b>Obtuse</b>	Blunt or rounded at the apex, with the sides meeting at an angle greater than 90°.
<b>Operculate</b>	With a small lid.
<b>Opposite</b>	A pair of organs attached at nodes in pairs on either side of a stem or axis.
<b>Orbicular</b>	Almost or approximately circular.
<b>Outbreeding depression</b>	A reduction in vigor of offspring from distant parents. It can occur when a locally adapted population is moved and mixed with plants adapted to different conditions.
<b>Outer canopy deciduous</b>	Marked reduction in leaf number in the outer canopy in exposed high light environments over winter.
<b>Oval</b>	Planar, shaped like a flattened circle, symmetrical about both the long and the short axis; about twice as long as broad, tapering equally both to the tip and the base. Synonymous with elliptical.
<b>Ovary</b>	Part of a flower containing the ovules and later the seeds.
<b>Ovate</b>	Egg-shaped and widest at base.
<b>Ovoid</b>	Oval; egg-shaped, with rounded base and apex.
<b>Pakihi</b>	A term which in its strict sense refers to open clears within forest dominated by low scrub and rushes. However, more usually used to refer natural and induced wetlands and their associated shrublands. A vernacular most frequently used in the West Coast for impoverished soils and their associated peats, left after forest has been cleared
<b>Palea</b>	The small upper bract enclosing the flower of a grass
<b>palea</b>	1. The upper of the two bracts that enclose each floret in a grass spikelet. 2. A small bract at the base of a disc floret in some plants of the composite family. 3. Scales on various parts of ferns (referred to as paleate or paleaceous). From the Latin word for 'chaff'.
<b>paleae</b>	Plural of palea, from the Latin word for 'chaff'. 1. The upper of the two bracts that enclose each floret in a grass spikelet. 2. A small bract at the base of a disc floret in some plants of the composite family. 3. Scales on various parts of ferns (referred to as paleate or paleaceous).
<b>Palmately</b>	Radiating from a point, as fingers radiating from the palm of a hand.
<b>Palmatifid</b>	Deeply divided into several lobes arising from more or less the same level.
<b>Palmatisect</b>	Intermediate between palmate and palmatifid, i.e. the segments are not fully separated at the base; often more or less digitate.
<b>Palustrine</b>	Pertaining to wet or marshy habitats. Term covers mires and marshes
<b>Pandurate</b>	Fiddle-shaped.
<b>Panicle</b>	Highly branched (multiple raceme).

Term	Definition
<b>Papilla</b>	A short rounded projection.
<b>Papillae</b>	A soft, fleshy projection, usually small and nipple-like.
<b>Papillate</b>	With short rounded projections.
<b>Papillose</b>	Warty, with short rounded projections or gland-dotted
<b>Parallel venation</b>	Veins are parallel along leaf.
<b>Parasite</b>	An organism that derives all its nourishment from its host.
<b>Patent</b>	Spreading or expanded, e.g., spreading petals.
<b>Peat</b>	A mass of partially carbonised plant tissue formed by partial decomposition in water of various plants and especially of mosses of the genus Sphagnum, widely found in many parts of the world, varying in consistency from a turf to a slime used as a fertiliser, as stable litter, as a fuel, and for making charcoal. Partially carbonized vegetable matter saturated with water; can be used as a fuel when dried. A type of soil deriving from dead organic material situated in a wet area, where the reduced amount of [[oxygen available in the wet conditions results in the organic material not decomposing as much as it usually would do so in the presence of more oxygen. Used in growing media. Represents an important carbon sink –drainage of peat releases large amounts of carbon (CO <sub>2</sub> ) to the atmosphere.
<b>Pedicel</b>	The stalk of a single flower in an inflorescence or fruit (either in a cluster or existing singularly).
<b>Peduncle</b>	The stalk of a solitary flower or the main stalk of an inflorescence or flower cluster.
<b>Pedunculate</b>	Describing fruits, which are borne on a stalk (a peduncle).
<b>Pellucid</b>	Transparent.
<b>Peltate</b>	Shield-like, with the stalk attached well inside the margin
<b>Pendent</b>	Hanging down from its support
<b>Pendulous</b>	Hanging or drooping.
<b>Penicillate</b>	With a tuft of hairs at the end, like a brush.
<b>Perennial</b>	A plant lasting for three seasons or more
<b>Perianth</b>	A collective term for the calyx (sepals or tepals) and corolla (petals) of the flower, especially when these are indistinguishable
<b>Petal</b>	Part of flower inside the sepals; usually coloured.
<b>Petiolate</b>	Having a petiole.
<b>Petiole</b>	Leaf stalk.
<b>phloem</b>	The vascular tissue in land plants that is primarily responsible for the distribution of sugars and nutrients manufactured in a shoot.
<b>Photopoint</b>	A monitoring technique where repeat photos are taken of the same scene from the same point over a period of time in order to quantify changes.
<b>Pilose</b>	Bearing long, soft hairs.
<b>Pinna</b>	A segment of a divided lamina that is classified as primary, secondary or tertiary according to the degree of dissection of the lamina.
<b>Pinnae</b>	Divisions of a pinnate leaf
<b>Pinnate</b>	With leaflets arranged regularly in two rows on either side of a stalk as in a feather; the lamina on a fern is divided into separate pinnae
<b>Pinnatifid</b>	Pinnately lobed, cleft more than halfway to the midrib. Not cleft all the way to the rachis.
<b>Pinnatisect</b>	Pinnately divided almost to midrib but segments still confluent.
<b>Pioneer</b>	Plant species are hardy species that should be planted first to establish a good canopy cover that restricts weed growth and promotes natural regeneration. In natural ecosystems these are the first plants to arrive and grow on a site.
<b>Pistil</b>	The female reproductive organ of a flower, consisting of an ovary, style, and stigma.
<b>Pistillate</b>	A flower with one or more pistils, but no stamens.
<b>Plano-convex</b>	Flat on one side, convex on the other.
<b>Plumose</b>	Feathery.
<b>Podzol</b>	Infertile, acidic soil, strongly leached to form a whitish-grey subsoil underlain by a layer enriched in iron, aluminium and organic matter; usually under forest in a wet temperate climate.
<b>Pole</b>	A subcanopy size individual with a long thin trunk and foliage tuft of a potential canopy tree.
<b>Pollinia</b>	Compact masses of orchid pollen.
<b>Population enhancement</b>	Increasing a population for a specific biological purpose, e.g., when a species is already present in an area but extra individuals are added to address a sex imbalance.
<b>Porrect</b>	Extending forward.
<b>Procumbent</b>	Lying and flat along the ground but not rooting
<b>Propagate</b>	To reproduce a plant by sexual (i.e., from seed) or asexual (e.g., from cuttings) means.
<b>Prostrate</b>	A general term for lying flat along the ground. This includes procumbent (that is lying and flat along the ground but not rooting) and decumbent (with a prostrate or curved base and an erect or ascending tip).
<b>Provenance</b>	The place of origin (of a plant that is in cultivation).
<b>Proximal</b>	Toward the base or point of attachment (cf. distal).
<b>Pseudobulb</b>	Thickened surface stem; usually looking like a bulb.
<b>Pseudoterminal</b>	Falsely terminal – as in a bud which appears to occupy a terminal position but does not

Term	Definition
<b>Puberulent</b>	Minutely clad in short, soft hairs
<b>Pubescence</b>	Covering of soft, fine hairs
<b>Pubescent</b>	Covered in short, soft hairs.
<b>Pungent</b>	Ending in a stiff sharp point
<b>Pustule</b>	Small blister-like elevation.
<b>Quadrante</b>	Square, rectangular.
<b>Raceme</b>	An unbranched, elongated inflorescence with pedicellate flowers maturing from the bottom upward i.e., flowers attached to the main stem by short stalks.
<b>Rachis</b>	the axis of an inflorescence or of a compound leaf
<b>Ray</b>	An outer ring of strap-like florets in the head of Asteraceae (daisy) flowers.
<b>Re-introduction</b>	Translocating wild or cultivated individuals to sites where the taxon has been known to occur in the past, but from which it has disappeared.
<b>Recurved</b>	Curved backward.
<b>Reflexed</b>	Bent back on itself
<b>Reniform</b>	Kidney shaped.
<b>Repend</b>	With a slightly wavy margin.
<b>Replum</b>	The outer structure of a pod in which the valves have dehisced (persists after the opening of the fruit)
<b>Restiad</b>	Area dominated by rush-like plants (collectively known as restiads) of the family Restionaceae. Includes Chatham Island and North Island Sporodanthus and oiioi ( <i>Apodasmia similis</i> )
<b>Retorse</b>	Pointing backward.
<b>Retuse</b>	A shallow notch at the rounded or blunt apex of a leaf.
<b>Rhizoid</b>	Any of various slender filaments that function as roots in mosses and ferns and fungi.
<b>Rhizomatous</b>	With underground creeping stems.
<b>Rhizome</b>	An underground stem (usually spreading horizontally or creeping) or short and erect.
<b>Rhombic</b>	Diamond-shaped.
<b>Rhomboid</b>	Diamond shaped, nearly rhombic.
<b>Riparian</b>	Relating to or living or located on the bank of a natural watercourse (as a river) or sometimes of a lake or a tidewater.
<b>Riparian margin</b>	Refers to the edges of streams, rivers, lakes or other waterways.
<b>Riparian plants</b>	Refers to plants found growing near the edges of streams, rivers or other waterways.
<b>Riparian zone</b>	A strip of land next to streams, rivers, and lakes where there is a transition from terrestrial (land vegetation) to aquatic (water) vegetation. Also known as "berm".
<b>Riverine</b>	Pertaining to rivers, streams and such like flowing water systems.
<b>Rootstock</b>	A short, erect, underground stem.
<b>Rosette</b>	A radiating cluster of leaves.
<b>Rostellum</b>	In orchids, a modified stigma that prevents self-fertilisation.
<b>Rosulate</b>	A dense radiating cluster of leaves.
<b>Rugose</b>	Wrinkled.
<b>Rugulose</b>	Having small wrinkles.
<b>Runcinate</b>	Sharply pinnatifid or cleft, the segments directed downward.
<b>Runner</b>	A trailing stem that roots at the nodes.
<b>Rupestral</b>	Growing on rocks.
<b>Rushes</b>	A group of distinctive wetland plants. They have solid stems (grasses have hollow stems), true rushes <i>Juncus</i> sp. have rounded leaves.
<b>Sagittate</b>	Shaped like the head of an arrow; narrow and pointed but gradually enlarged at base into two straight lobes directed downwards; may refer only to the base of a leaf with such lobes; cf. hastate.
<b>Salt marsh</b>	A coastal wetland, with specialized salt tolerant plants (halophytes).
<b>Sapling</b>	A juvenile tree that has reached the stage of 1 or 2 main stems but is still in the shrub layer.
<b>Saprophyte</b>	A plant lacking chlorophyll and living on dead organic matter.
<b>Saprophytic</b>	Lacking chlorophyll and living on dead organic matter.
<b>Sarcotesta</b>	The fleshy, often highly coloured outer layer of the seed coat in some species, e.g., titoki ( <i>Alectryon excelsus</i> ).
<b>Scabrid</b>	Roughened or rough with delicate and irregular projections.
<b>Scale</b>	Any thin, flat, membranous structure.
<b>Scape</b>	A leafless flower stem.
<b>schizocarp</b>	A fruit which splits when dry, from the Greek <i>skhizein</i> 'split' and <i>karpos</i> 'fruit'
<b>schizocarps</b>	Plural of schizocarp, a fruit which splits when dry, from the Greek <i>skhizein</i> 'split' and <i>karpos</i> 'fruit'
<b>Scutiform</b>	Shield-shaped.
<b>Sedges</b>	A group of grass-like or rush-like herbaceous plants belonging to the family Cyperaceae. Many species are found in wetlands some are forest floor plants. Leaves are usually angular. Hence the saying "rushes are round and sedges have edges".



<b>Term</b>	<b>Definition</b>
<b>Seedling</b>	A newly germinated plant.
<b>Self sustaining</b>	Able to sustain itself, or replace itself, independently of management i.e. regenerate naturally
<b>Self thinning</b>	Natural tree death in a crowded, even-aged forest or shrubland.
<b>Semi-deciduous</b>	Partial leaflessness in winter, and greater than 50% leaves lost by the beginning of spring flush.
<b>Sepal</b>	Outer part of flower; usually green.
<b>Serrate</b>	Sharply toothed with teeth pointing forwards towards apex.
<b>Serrulate</b>	Finely serrate, i.e., finely toothed with asymmetrical teeth pointing forward; like the cutting edge of a saw.
<b>Sessile</b>	Attached by the base without a stalk or stem.
<b>Seta</b>	The stalk of a fruiting moss capsule
<b>Sheath</b>	A portion of an organ that surrounds (at least partly) another organ (e.g., the tubular envelope enclosing the stem in grasses and sedges).
<b>Silicles</b>	The flattened usually circular capsule – compared with the narrow, elongated fruit (silique) – containing the seed/seeds. A term used almost exclusively for plants within the cabbage family (Brassicaceae)
<b>Silique</b>	A capsule, usually 2-celled, with 2 valves falling away from a frame (replum) bearing
<b>Simple</b>	Of one part; undivided (cf compound).
<b>Sinuate</b>	With a wavy margin.
<b>Sinus</b>	The space or recess between lobes; in hebes a gap between the margins of two leaves of an opposite pair that may be present in the bud before the pair of leaves separate.
<b>Sorus</b>	A cluster of two or more sporangia on the margin or underside of the lamina of a fern, sometimes protected by an indusium.
<b>Spathulate</b>	Spatula or spoon-shaped, a rounded blade tapering gradually to the base.
<b>Spheroidal</b>	Almost spherical but elliptic in cross section.
<b>Spicate</b>	Arranged in a spike.
<b>Spike</b>	Flowers attached to main stem without stalks.
<b>Spikelet</b>	Collection of individual grass florets borne at the end of the smallest branch of the inflorescence.
<b>Sporangia</b>	Plural of sporangium. Structures in which spores are produced.
<b>Sporangium</b>	Structure in which spores are produced.
<b>Spore</b>	A single-celled reproductive unit similar in function to that of the seed in a flowering plant.
<b>sporophyte</b>	The spore producing plant in ferns that is usually the visible part.
<b>Stamen</b>	The male reproductive organ of a flower where pollen is produced. Consists of an anther and its stalk.
<b>Stamens</b>	The male, pollen bearing organ of a flower.
<b>Standing water</b>	Where water lies above the soil surface for much of the year.
<b>Stellate</b>	Irregularly branched or star shaped.
<b>Stigma</b>	Female part of the flower that is receptive to pollen, usually found at or near the tip (apical end) of the style where deposited pollen enters the pistil.
<b>Stipe</b>	The stalk of a frond.
<b>Stipitate</b>	Borne on a stipe or stalk.
<b>Stipulate</b>	A leaf with stipules.
<b>Stipule</b>	A scale-like or leaf-like appendage at the base of a petiole, usually paired.
<b>Stolon</b>	A stem which creeps along the ground, or even underground.
<b>Stoloniferous</b>	Producing stolons
<b>Stramineous</b>	Chaffy, like straw or straw-colored.
<b>Stria</b>	A fine line or groove.
<b>Striae</b>	Fine lines or grooves.
<b>Striate</b>	Fine longitudinal lines or minute ridges
<b>Style</b>	The elongated part of the flower between the ovary and the stigma.
<b>Sub-</b>	A prefix meaning under, somewhat or almost.
<b>Subglabrous</b>	Very slightly, but persistently, hairy.
<b>Suborbicular</b>	Slightly rounded in outline
<b>Substrate</b>	The surface upon which an orchid grows.
<b>Subtended</b>	Immediately beneath, occupying a position immediately beneath a structure, i.e., flower subtended by bract
<b>Subulate</b>	Slender and tapering to a point.
<b>Succession</b>	Progressive replacement of one species or plant community type by another in an ecosystem.
<b>Successional</b>	Referring to species, plant communities or habitats that tend to be progressively replaced by another.
<b>Succulent</b>	Fleshy and juicy.
<b>Summer-green</b>	Used in New Zealand to indicate herbs or sub-shrubs that die down to a root stock or rhizomatous network.
<b>Supplementary planting</b>	Returning to a revegetation site and creating gaps, or filling existing gaps, with different plants of plants, usually later successional plants which may not have survived being planted in the first phases of the project.

<b>Term</b>	<b>Definition</b>
<b>Surface water</b>	Water present above the substrate or soil surface.
<b>Surveillance</b>	Regular survey for pests inside operational and managed areas e.g. nurseries, standout areas on parks.
<b>Survey</b>	Collection of observations on the spatial distribution or presence or absence of species using standardised procedures.
<b>Sustainable Land Management</b>	The use of farming practices which are sustainable both financially and environmentally including management of nutrient runoff, waste disposal or stock effluent, reducing impacts of nutrients on waterways, preventing erosion and soil loss, and protecting native forest and wetland habitats from stock damage.
<b>Swamp</b>	Low land that is seasonally flooded; has more woody plants than a marsh and better drainage than a bog. They are more fertile and less acidic than bogs because inflowing water brings silt, clay and organic matter. Typical swamp plants include raupo, purei and harakeke (flax). Zonation and succession often leads through manuka to kahikatea swamp forest as soil builds up and drainage improves.
<b>Symbiote</b>	An organism that has an association with organisms of another species whereby the metabolic dependence of the two associates is mutual.
<b>Symbiotic</b>	The relation between two different species of organisms that are interdependent; each gains benefits from the other (see also symbiosis).
<b>Sympatric</b>	Occupying the same geographical region.
<b>Synangia</b>	Structures made up of fused sporangia
<b>Synonym</b>	A botanical name that also applies to the same taxon.
<b>Systematics</b>	The study of taxonomy, phylogenetics, and taxagenetics.
<b>Tabular</b>	Shaped like a rectangular tablet.
<b>Taxa</b>	Taxonomic groups. Used to refer to a group at any level e.g., genus, species or subspecies.
<b>Taxon</b>	A taxonomic group. Used to refer to a group at any level e.g., genus, species or subspecies.
<b>Taxonomy</b>	The process or science of classifying, naming, and describing organisms
<b>Tepal</b>	An individual member of the perianth.
<b>Terete</b>	Cylindrical and tapering.
<b>Terminal</b>	At the tip or apex.
<b>Ternatifid</b>	Leaflets In threes,
<b>Tetrad</b>	A group of four.
<b>Tomentum</b>	A hairy covering of short closely matted hairs.
<b>Translocation</b>	The movement of living organisms from one area to another.
<b>Trifid</b>	Divided into three.
<b>Trifoliate</b>	Having three leaflets.
<b>Trigonus</b>	Three-angled
<b>Tripinnate</b>	With each secondary pinna divided to the midrib into tertiary pinnae
<b>Triquetrous</b>	Triangular in cross section and acutely angled.
<b>Truncate</b>	With the apex or base squared at the end as if cut off.
<b>Tuberculate</b>	Bearing small swellings.
<b>Tubular</b>	Tube-shaped.
<b>turbinate</b>	Top-shaped.
<b>Turgid</b>	Distended through internal pressure
<b>Type locality</b>	The place or source where a holotype or type specimen was found for a species.
<b>Ultramafic</b>	A type of dark, usually igneous, rock that is chemically dominated by magnesium and iron-rich minerals, the partially metamorphosed form of which is serpentinite.
<b>Umbel</b>	Umbrella like; the flower stalks arise from one point at the stem.
<b>Undulate</b>	Wavy edged.
<b>Undulose</b>	Wavy edged.
<b>Unitubular</b>	A tube partitioned once – literally one tube (compare – multitubular – many tubes)
<b>Utricle</b>	A thin loose cover enveloping some fruits (eg., Carex, Uncinia)
<b>Valvate</b>	Opening by valves.
<b>Vascular plant</b>	A plant that possesses specialised conducting tissue (xylem and phloem). This includes flowering plants, conifers and ferns but excludes mosses, algae, lichens and liverworts.
<b>Velutinous</b>	Thickly covered with delicate hairs; velvety.
<b>Ventral</b>	Of the front or inner (adaxial) surface relative to the axis. (cf. dorsal)
<b>Vermiform</b>	Worm-shaped.
<b>Vernicose</b>	Glossy, literally as if varnished, e.g., Hebe vernicosa has leaves that appear as if varnished
<b>Verrucose</b>	Having small rounded warts.
<b>Verticillium</b>	A fungus disease that will cause wilting and death.
<b>Villous</b>	Covered with long, soft, fine hairs.
<b>Water table</b>	The level at which water stays in a soil profile. The zone of saturation at the highest average depth during the wettest season.
<b>Wetland</b>	A site that regularly has areas of open water for part or all of the year, or has a water table within 10 cm of the surface for at least 3 months of the year. Wetland ecosystems support a range of plant and animal species adapted to an aquatic or semi-aquatic environment.

<b>Term</b>	<b>Definition</b>
-------------	-------------------

<b>Whipcord</b>	A shrub in which the leaves are reduced to scales that are close-set and pressed against the stem.
-----------------	--

<b>Whorl</b>	A ring of branches or leaves arising at the same level around the stem of a plant.
--------------	--