

# Chionochloa acicularis

## COMMON NAME

needle snow tussock

## SYNONYMS

None

## FAMILY

Poaceae

## AUTHORITY

*Chionochloa acicularis* Zotov

## FLORA CATEGORY

Vascular – Native

## ENDEMIC TAXON

Yes

## ENDEMIC GENUS

No

## ENDEMIC FAMILY

No

## STRUCTURAL CLASS

Grasses

## NVS CODE

CHIACI

## CHROMOSOME NUMBER

2n = 42

## CURRENT CONSERVATION STATUS

2017 | Not Threatened | Qualifiers: DP

## PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Range Restricted

## DISTRIBUTION

Endemic. New Zealand: South Island (South Westland and Fiordland).

## HABITAT

Lowland to upper montane. On peaty soils in grasslands, and scattered through scrub

## WETLAND PLANT INDICATOR STATUS RATING

FAC: Facultative

Commonly occurs as either a hydrophyte or non-hydrophyte (non-wetlands).



Fiordland. Photographer: Kelvin Lloyd, Licence: All rights reserved.



Franklin Mountains, Fiordland. Photographer: John Barkla, Date taken: 13/11/2012, Licence: CC BY.

## DETAILED DESCRIPTION

Tall, slender, erect, pedicelled tussock, with sharp-pointed, glabrous, deciduous leaves. **Leaf-sheath** to 120 mm, dark above, pale shining below, persistent, becoming chartaceous, keeled, clothed with long (3 mm) abundant deciduous hairs, apical tuft of hairs to 7 mm. **Ligule** to 0.5 mm. Leaf-blade to 500 × 1 mm diameter, acicular rush-like, thickened at midrib, disarticulating at ligule, abaxially glabrous, adaxially with dense very short stiff hairs below, papillae and occasionally some prickle-teeth above; margin smooth. **Culm** to 600 mm, long and short hairs below inflorescence otherwise glabrous. **Inflorescence** to 100 mm, spikelets few; rachis, branches and pedicels abundantly long hairy especially at branch axils. **Spikelets** of up to 7 golden florets. **Glumes** becoming acute or mucronate, glabrous, > adjacent lemma lobes; lower to 11 mm, 3-nerved, upper to 13 mm, 5-nerved. **Lemma** to 5 mm; dense hairs at margin and erect hairs aside central nerve or rarely between all nerves, ± reaching sinus; lateral lobes to 4 mm, triangular-acute; central awn to 14 mm from divergent flat column to 2 mm. **Palea** to 7 mm. **Callus** to 1 mm, hairs to 2.5 mm. **Rachilla** to 0.5 mm. **Lodicules** to 1 mm. **Anthers** to 4 mm. **Ovary** to 0.8 mm; stigma-styles to 2 mm. **Seeds** to 3 mm.

## MANAAKI WHENUA ONLINE INTERACTIVE KEY

[Key to the grasses of New Zealand](#)

## FLOWERING

October–December

## FRUITING

November–February

## LIFE CYCLE

Florets are wind dispersed (Thorsen et al., 2009).

## PROPAGATION TECHNIQUE

Easily grown from fresh seed and rooted pieces. Prefers a permanently damp, acidic soil and semi-shade. Plants must never be allowed to dry out.

## THREATS

A naturally uncommon, regional endemic.

## ETYMOLOGY

**chionochloa**: Snow grass

**acicularis**: From the Latin acicula 'small pin' refers to things which are needle-like, needle-shaped, stiff or pointed

## WHERE TO BUY

Not commercially available

## ATTRIBUTION

Description modified from Edgar and Connor (200)

## REFERENCES AND FURTHER READING

Edgar E, Connor HE. 2000. Flora of New Zealand. Vol. V. Grasses. Christchurch, Manaaki Whenua Press. 650 p.

Thorsen MJ, Dickinson KJM, Seddon PJ. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285–309.

## MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/chionochloa-acicularis/>