

Oreobolus pectinatus

COMMON NAMES

comb sedge, cushion sedge, flat-leaved comb sedge

SYNONYMS

Oreobolus serrulatus Col.; *Oreobolus pumilio* var. *pectinatus* (Hook.f.)
C.B.Clarke

FAMILY

Cyperaceae

AUTHORITY

Oreobolus pectinatus Hook.f.

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Sedges

NVS CODE

OREPEC

CHROMOSOME NUMBER

$2n = c.40$

CURRENT CONSERVATION STATUS

2017 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

DISTRIBUTION

Endemic. North Island, South Island and Stewart Island/Rakiura, Auckland Islands and Campbell Island/Motu Ihupuku.

HABITAT

Coastal to alpine (up to 1500 m a.s.l.). Mostly alpine descending to sea level only in the southern South Island, Stewart Island/Rakiura, Auckland Islands and Campbell Islands. A common species of cushion bogs and alpine seepages and mires and also favouring poorly drained open ground. Very rarely found growing in damp peaty ground under low subalpine scrub.

WETLAND PLANT INDICATOR STATUS RATING

OBL: Obligate Wetland

Almost always is a hydrophyte, rarely in uplands (non-wetlands).



Ruahine range, March. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.



Oreobolus pectinatus, Hawkdun Range. Photographer: John Barkla, Licence: CC BY.

DETAILED DESCRIPTION

Perennial sedge forming dense grey-green to green cushions 10–100 mm tall. **Stems** densely packed, much branched at base, leafy. **Leaves** dark green, green to grey-green, mostly distichous; lamina 0.5–1.0 mm wide, abaxial surface convex, median nerve and two lateral nerves visible at widest part of lamina, adaxial surface channelled, only the median nerve prominent, both surfaces with abundant stomata; sheath 5–7-nerved, not lobed at apex. **Spikelets** usually solitary; mature peduncle usually > leaves. **Glumes** usually 3–(4), usually light green with broad, colourless, membranous margins, the outermost longer, leaf-like, the two inner more or less equal, membranous, the occasional fourth glume, smaller. **Hypogynous scales** < nut, lanceolate, initially colourless, maturing red-brown to almost black. **Nut** slightly > 1 mm long, < 1 mm diameter, light brown or often dark brown with a triangular, pubescent depression at apex.

SIMILAR TAXA

Distinguished from the other endemic species *Oreobolus impar* Edgar and *O. strictus* Bergg. by the leaves which are mostly distichous; by the median nerve and 2 lateral nerves visible on the lower surface at the widest part of the leaf; and by the usually light green glumes.

FLOWERING

October–December

FRUITING

November–April

PROPAGATION TECHNIQUE

Easily grown from rooted pieces and probably from fresh seed. However, this species is best grown in a cooler climate or kept in a pot within an alpine house. It is very slow growing.

ETYMOLOGY

oreobolus: Mountain clump

pectinatus: Like a comb

WHERE TO BUY

Not commercially available.

ATTRIBUTION

Description adapted from Moore and Edgar (1970).

Some of this factsheet information is derived from [Flora of New Zealand Online](#) and is used under a [Creative Commons Attribution 3.0 New Zealand](#) licence.

REFERENCES AND FURTHER READING

Moore LB, Edgar E. 1970. Flora of New Zealand, Volume II. Indigenous Tracheophyta: Monocotyledones except Gramineae. Government Printer, Wellington, NZ. 354 p.

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/oreobolus-pectinatus/>

PDF DATE

17 September 2024