

Potamogeton suboblongus

COMMON NAMES

mud pondweed

FAMILY

Potamogetonaceae

AUTHORITY

Potamogeton suboblongus Hagström

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Herbs - Monocots

NVS CODE

POTSUB

CHROMOSOME NUMBER

2n = 28

CURRENT CONSERVATION STATUS

2017 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

DISTRIBUTION

Endemic. New Zealand: North, South and Stewart Islands


HABITAT

Coastal to subalpine, being mostly found in upper montane and subalpine areas in the northern part of its range. *Potamogeton suboblongus* is more commonly found in shallow, muddy hollows in forest, and colonising tarns and alpine soaks and pools which may partially dry out in summer.

WETLAND PLANT INDICATOR STATUS RATING

OBL: Obligate Wetland

Almost always is a hydrophyte, rarely in uplands (non-wetlands).



In cultivation ex Pureora. Photographer: Jeremy R. Rolfe, Date taken: 29/10/2007, Licence: CC BY.



Central Southland. Photographer: Jesse Bythell, Date taken: 26/02/2016, Licence: CC BY-NC.

DETAILED DESCRIPTION

Aquatic, submerged or floating, or ± semi-terrestrial on muddy ground and seepages, rhizomatous little-branched, perennial herb. Rhizomes usually buried in firm mud; leafy branches erect, usually simple, long or short, often rooted at leaf-bearing nodes. Stipules c.20–30 mm long, free, open, membranous and often conspicuous. Leaves weakly dimorphic, entire; lower leaves apparently not regularly submersed, similar to upper leaves, being somewhat more narrowly elliptic and with fewer, more obscure nerves than emergent leaves; upper and emergent leaves long- or short-petiolate, lamina 15–60 × 8–35 mm, broad-elliptic, rarely oval, subacute and sometimes shortly attenuate at tip, firm and opaque, longitudinal nerves 11–21 in all, usually several laterals emerging from midrib. Inflorescence a densely flowered spike, 10–20 mm long. Peduncles arising in axils of upper leaves, simple, stout, erect, standing well above leaves. Achenes c.2.0 × 1.5 mm, reddish when ripe, neither flattened nor strongly keeled, beak short but slightly curved.

SIMILAR TAXA

Distinguished from *Potamogeton cheesemanii* by the scarcely dimorphic foliage (i.e. only slight differences between submersed and emergent leaves), by the near absence of submersed leaves (these if present of similar size and shape to emergent foliage), by the broadly elliptic, subacute floating leaves, which have usually 6 or more nerves on either side of midrib; and by the swollen, reddish, not or only weakly keeled achenes.

FLOWERING

December - March

FRUITING

January - April

PROPAGATION TECHNIQUE

Easily grown from rooted pieces and fresh seed, An attractive plant for a small, shallow pond or damp, shaded hollow. In warmer, lowland areas this species prefers a cool site

ETYMOLOGY

potamogeton: River dweller

suboblongus: Nearly oblong

ATTRIBUTION

Fact sheet prepared for NZPCN by P.J. de Lange (16 February 2012). Description adapted from Moore & Edgar (1970).

Some of this factsheet information is derived from [Flora of New Zealand Online](#) and is used under a [Creative Commons Attribution 3.0 New Zealand](#) licence.

REFERENCES AND FURTHER READING

Moore, L.B.; Edgar, E. 1970: Flora of New Zealand. Vol. II. Government Printer, Wellington.

NZPCN FACT SHEET CITATION

Please cite as: de Lange, P.J. (Year at time of access): *Potamogeton suboblongus* Fact Sheet (content continuously updated). New Zealand Plant Conservation Network.

<https://www.nzpcn.org.nz/flora/species/potamogeton-suboblongus/> (Date website was queried)

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/potamogeton-suboblongus/>

PDF DATE

17 September 2024