

# Azorella cockaynei

## SYNONYMS

Schizeilema cockaynei (Diels) Cheeseman

## FAMILY

Apiaceae

## AUTHORITY

Azorella cockaynei Diels

## FLORA CATEGORY

Vascular – Native

## ENDEMIC TAXON

Yes

## ENDEMIC GENUS

No

## ENDEMIC FAMILY

No

## STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

## NVS CODE

SCHCOC

## CHROMOSOME NUMBER

2n = 48

## CURRENT CONSERVATION STATUS

2017 | Not Threatened

## PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

## DISTRIBUTION

Endemic. South and Stewart Islands, local from Otago south throughout Southland and on Stewart island

## WETLAND PLANT INDICATOR STATUS RATING

FACW: Facultative Wetland

Usually is a hydrophyte but occasionally found in uplands (non-wetlands).

## FLOWERING

November - December

## FRUITING

December - March

## PROPAGATION TECHNIQUE

Easily grown in a damp soil. Best within a partially submerged pot.

## ETYMOLOGY

**cockaynei**: Named after Leonard Cockayne FRS (7 April 1855 - 8 July 1934) who is regarded as New Zealand's greatest botanist and a founder of modern science in New Zealand



Azorella cockaynei in alpine streamside herbfield, Old Man Range. Photographer: John Barkla, Date taken: 21/03/2020, Licence: CC BY.



Azorella cockaynei in coastal turf with Samolus repens, Riverton, Southland. Photographer: John Barkla, Date taken: 25/12/2017, Licence: CC BY.

**WHERE TO BUY**

Not commercially available.

**MORE INFORMATION**

<https://www.nzpcn.org.nz/flora/species/azorella-cockaynei/>

**PDF DATE**

17 September 2024