

Carex maorica

COMMON NAMES

Māori sedge

SYNONYMS

Carex fascicularis Boott var. *minor* Boott; *Carex forsteri* Wahl. var. *minor* (Boott) Hook.f.

FAMILY

Cyperaceae

AUTHORITY

Carex maorica Hamlin

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Sedges

NVS CODE

CARMAO

CHROMOSOME NUMBER

2n = c.72-76

CURRENT CONSERVATION STATUS

2017 | Not Threatened

PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

DISTRIBUTION

Endemic. New Zealand: North and South Islands. In the North Island uncommon in the east from East Cape to the Wairarapa otherwise widespread. In the South Island apparently absent from Southland and Fiordland

HABITAT

Coastal to lowland in freshwater wetlands, under willow in gully systems, along river and stream banks, lake margins, and in damp seepages, pond margins and clearings within forest. Preferring fertile to mid-fertile wetlands.

WETLAND PLANT INDICATOR STATUS RATING

OBL: Obligate Wetland

Almost always is a hydrophyte, rarely in uplands (non-wetlands).



Taken in Coromandel, January. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.



Atiwhakatu Valley, Wairarapa. Photographer: Jeremy R. Rolfe, Date taken: 02/02/2008, Licence: CC BY.

DETAILED DESCRIPTION

Light green to yellow-green tufted sedge. **Culms** 150–700 × 1.0–2.5 mm., trigonous, smooth or faintly scabrid below inflorescence; basal sheaths light to dark grey, often red-tinged. **Leaves** > culms, to 1 m long, 2–7 mm wide, double-folded, cross-veinlets ± prominent, keel and margins minutely scabrid. **Inflorescence** of 2–5 close-set, sessile, usually erect, occasionally spreading spikes, or the lowest 1–2 rather distant and shortly pedunculate; terminal spike male; remaining spikes female, usually > and overtopping male spike, 20–60 × 7–12 mm, crowded at same level round base of male spike. **Glumes** usually much < utricles, 1–2 mm long, narrowly ovate-lanceolate, margin of upper part fimbriate or lacerate, hyaline with a green midrib produced to a scabrid awn 1–3 mm long, up to 6 mm long in lowermost glumes. **Utricles** 4.0–5.5–(6.0) × 1.0–1.5 mm., plano-convex or biconvex, turgid, ovate or lanceolate, spreading when ripe, shining light green to light brown with numerous distinct white nerves, very slightly tapered above to a beak 1.5–2.0 mm long, with bifid, glabrous orifice, crura c. ⅓ length of entire beak; stipe narrow, c. 0.5 mm long. **Stigmas** 3. **Nut** c. 1.5 mm. long, triquetrous, ellipsoid, cream or light brown.

SIMILAR TAXA

Carex maorica is easily recognised by its slender culms; pale green to yellow green wide leaves; prominent cross-veinlets on sheaths and leaves; and mostly clustered spikelets, and short glumes with lacerate or fimbriate margins; and by the submembranous, strongly nerved, glabrous, spreading utricles. There are only two other carices in New Zealand with cross-veinlets, the indigenous *C. fascicularis* Boott and naturalised *C. lurida* Wahlenberg. *Carex fascicularis* differs from *C. maorica* by the female spikes mostly distant to ± approximate (but then never clustered at one level round base of male spike). *Carex lurida* is a much larger sedge than either species (up to 2.5 m tall) and has utricles 6–9 × 2–4 rather than 4–6 × 1.0–1.5 mm

FLOWERING

October—December

FRUITING

November—May

LIFE CYCLE

Nuts surrounded by inflated utricles are dispersed by granivory and wind (Thorsen et al., 2009).

PROPAGATION TECHNIQUE

Easily grown from fresh seed and by the division of established plants. Although a wetland species *C. maorica* will grow well in most soils and moisture regimes. Does best in full sun.

ETYMOLOGY

carex: Latin name for a species of sedge, now applied to the whole group.

ATTRIBUTION

Fact sheet prepared by P.J. de Lange (12 August 2006). Description adapted from Moore and Edgar (1970). Some of this factsheet information is derived from [Flora of New Zealand Online](#) and is used under a [Creative Commons Attribution 3.0 New Zealand](#) licence.

REFERENCES AND FURTHER READING

Moore LB, Edgar E. 1970. Flora of New Zealand, Volume II. Indigenous Tracheophyta: Monocotyledones except Gramineae. Government Printer, Wellington, NZ. 354 p.
Thorsen MJ, Dickinson KJM, Seddon PJ. 2009. Seed dispersal systems in the New Zealand flora. *Perspectives in Plant Ecology, Evolution and Systematics* 11: 285–309.

NZPCN FACT SHEET CITATION

Please cite as: de Lange, P.J. (Year at time of access): *Carex maorica* Fact Sheet (content continuously updated). New Zealand Plant Conservation Network. <https://www.nzpcn.org.nz/flora/species/carex-maorica/> (Date website was queried)

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/carex-maorica/>

PDF DATE

17 September 2024