

Acaena microphylla var. microphylla

COMMON NAMES

bidibid, pipiriri

FAMILY

Rosaceae

AUTHORITY

Acaena microphylla Hook.f. var. *microphylla*

FLORA CATEGORY

Vascular – Native

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No

ENDEMIC FAMILY

No

STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

CHROMOSOME NUMBER

2n = c.42

CURRENT CONSERVATION STATUS

2017 | Not Threatened | Qualifiers: SO

PREVIOUS CONSERVATION STATUSES

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

FLOWER COLOURS

Red/Pink

LIFE CYCLE

Spiny hypanthia are dispersed by attaching to fur, feathers and clothing and possibly also dispersed by wind and granivory (Thorsen et al., 2009)

ETYMOLOGY

acaena: From the Greek 'akanthos' thorn, referring to the spiny calyx that many species have

microphylla: Small leaf

REFERENCES AND FURTHER READING

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 2009 Vol. 11 No. 4 pp. 285-309

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/acaena-microphylla-var-microphylla/>

PDF DATE

27 September 2024



Tongariro, January. Photographer: John Smith-Dodsworth, Licence: CC BY-NC.