

Aciphylla hookeri

BIOSTATUS

Native

CATEGORY

Vascular

STRUCTURAL CLASS

Herbs - Dicotyledons other than Composites

DETAILED DESCRIPTION

Tufted sparse-leaved herb up to 30 cm tall, stock usually simple, taproot stout. **Leaves** up to approximately 20 cm long, 2-1 pinnate. Sheath very thin, approximately 70 x 10 mm, narrowing gradually to apex; stipules approximately 10 mm long, simple or sometimes with accessory segments; petiole crenulate-serrulate, approximately 40 mm long; lower internodes rather distant. **Primary pinnae** up to 20 mm long, with 1-(2) pairs of pinnules; all segments keeled, grooved, pungent, giving a squarrose appearance. **Flowering stems** sometimes leafy below, stout, grooved, up to 30 cm long including narrow paniculate inflorescence approximately 150 mm long. **Bracts** usually many, leaflike, up to approximately 100 mm long; sheaths roughly 50 mm long; stipules up to 10 mm long, unequal, acicular; lamina pinnate, approximately 50 mm long, segments pungent. **Umbels** several, on 5-10 unequal slender rays less than bracts. Umbellules small, on short slender rays. **Fruit** approximately 4-5 mm long, mericarps with 3-5 wings.

SIMILAR TAXA

Aciphylla anomala is similar in habit, but differs by longer thicker petioles (80 x 2 mm) and leaflets (100 x 2.5 mm); flower bract stems have shorter stipules (5 mm).

DISTRIBUTION

Endemic. South Island, North-west Nelson to Aurthur's Pass mostly on western side.

HABITAT

Low-alpine (900-1400 m.a.s.l.) uncommon, snow tussock-herbfield.

CURRENT CONSERVATION STATUS

2023 | Not Threatened | Qualifiers: DPS, DPT

DETAILED TAXONOMY

FAMILY

Apiaceae

AUTHORITY

Aciphylla hookeri Kirk

TAXONOMIC NOTES

Dawson & LeComte suggest this species is part of the group with clear juice. They note it may also be seen to hybridise; *A. hookeri* x *A. indurata* seen at Mt. Bovis, reported by W. R. B. Oliver; *A. hookeri* x *A. townsonii* seen in the Lyell Range, reported J. W. Dawson (In the last case the probable hybrids were growing in large patches of granite boulders and were so varied that they must include back crosses).

ENDEMIC TAXON

Yes

ENDEMIC GENUS

No



Croesus Track. Photographer: Melissa Hutchison, Licence: CC BY-NC.

ENDEMIC FAMILY

No

ECOLOGY

FLOWERING

December – February (-March)

FRUITING

January – April

LIFE CYCLE

Winged schizocarps are dispersed primarily by wind (Thorsen et al., 2009).

OTHER INFORMATION

ETYMOLOGY

aciphylla: From the Latin acicula 'needle' and the Greek phyllum 'leaf', meaning needle-leaf.

hookeri: Named after Sir Joseph Dalton Hooker (born 1817) - a world famous botanist who travelled on the Antarctic expedition of 1839 under the command of Sir James Ross and wrote "Handbook of New Zealand Flora" published in 1864-67 describing many specimens sent to Kew by collectors. He died in 1911 and has a memorial stone at Westminster Abbey London.

NVS CODE

ACIHOO

PREVIOUS CONSERVATION STATUSES

2017 | Not Threatened | Qualifiers: DP

2012 | Not Threatened

2009 | Not Threatened

2004 | Not Threatened

REFERENCING AND CITATIONS

REFERENCES AND FURTHER READING

Allan, H. H. 1961. Flora of New Zealand. Vol. 1. Wellington: Government Printer. pg. 476-477.

Dawson, J.W. LeComte, J.R. 1978. Research on Aciphylla - a progress report. Tuatara 23: pg. 49-67.

Mark, A. F. 2012. Above the Treeline: A Nature Guide to Alpine New Zealand. Craig Potton Publishing, Nelson. pg. 148.

Thorsen, M. J.; Dickinson, K. J. M.; Seddon, P. J. 2009. Seed dispersal systems in the New Zealand flora. Perspectives in Plant Ecology, Evolution and Systematics 2009 Vol. 11 No. 4 pp. 285-309

ATTRIBUTION

Description adapted by M. Ward from Allan (1961).

Some of this factsheet information is derived from [Flora of New Zealand Online](#) and is used under a [Creative Commons Attribution 3.0 New Zealand](#) licence.

MORE INFORMATION

<https://www.nzpcn.org.nz/flora/species/aciphylla-hookeri/>

PDF DATE

03 October 2024